



SRM
INSTITUTE OF SCIENCE & TECHNOLOGY
Deemed to be University u/s 3 of UGC Act, 1956



**SOUTHERN INDIA
SCIENTIFIC CORPORATION**



Lawrence & Mayo
ESTD 1977



PROCEEDINGS OF

NATIONAL SYMPOSIUM ON INNOVATIVE APPROACHES FOR INTEGRATED PEST MANAGEMENT (IPM NEXUS 25)

OCTOBER 10, 2025



**Organized by
Department of Entomology
SRM College of Agricultural Sciences
SRM Institute of Science and Technology
Baburayanpettai, Chengalpattu Dist. - 603 201**





SRM COLLEGE OF AGRICULTURAL SCIENCES
SRM INSTITUTE OF SCIENCE AND TECHNOLOGY
(Deemed to be University u/s 3 of UGC Act, 1956)

Proceedings of
NATIONAL SYMPOSIUM ON INNOVATIVE APPROACHES FOR
INTEGRATED PEST MANAGEMENT (IPM NEXUS 25)
10th October, 2025

SPONSORS



SOUTHERN INDIA
SCIENTIFIC CORPORATION



Lawrence & Mayo
ESTD 1877



Organized by

Department of Entomology
SRM College of Agricultural Sciences, SRMIST
Baburayanpettai, Chengalpattu-603 201
Tamil Nadu
October 2025

Citation

Proceedings of National Symposium on Innovative Approaches for Integrated Pest Management (IPM Nexus 25) held at SRM College of Agricultural Sciences, SRMIST, Baburayanpettai, Chengalpattu, Tamil Nadu, India on 10th October, 2025.

Editors

Dr. R. Nisha

Dr. M. Muthukumar

Dr. L. Ramazeame

Dr. N. Murugan

Mr. D. Kamaraj

Dr. M. Jawaharlal



DOI: <http://dx.doi.org/10.22573/spg.025.evro/S/040>

Published By,

Skyfox Publishing Group

Skyfox Press

#1789, Medical College Road

Thanjavur-613004

Tamil Nadu, India.

ACKNOWLEDGEMENT

We are thanking SRMIST for providing funding support for organizing this **National Symposium on Innovative Approaches for Integrated Pest Management (IPM Nexus 25)** on October 10, 2025 at SRM College of Agricultural Sciences, Acharapakkam Campus.

We are thankful for **Southern Indian Scientific Corporation (Chennai), “LYNX” Lawrence & Mayo (Chennai) and Krishnaaram Laboratories (Chennai)** for their financial grant for organizing this National Symposium.

**NATIONAL SYMPOSIUM ON INNOVATIVE APPROACHES FOR
INTEGRATED PEST MANAGEMENT (IPM NEXUS 25)
EVENT COMMITTEE**

Chief Patron

Dr. T. R. Parivendhar

Chancellor

SRMIST

Dr. P. Ravipachamuthu

Pro-Chancellor (Admin)

SRMIST.

Dr. P. Sathyanaryanan

Pro-Chancellor (Academics)

SRMIST.

Dr. R. Shivakumar

Vice President

SRMCAS.

Dr. Muthamizhchelvan

Vice Chancellor

SRMIST

Dr. S. Ponnusamy

Registrar

SRMIST

Convenor

Dr. M. Jawaharlal

Dean, SRM CAS

Head of The Department

Dr. M. Muthukumar, Asst. prof. (Entomology) (Sr. Grade)

Organizing Secretary

Dr. R. Nisha, Asst. prof. (Entomology)

Co-organizing Secretaries

Dr. L. Ramazeame, Asst. prof. (Entomology)

Dr. N. Murugan, Asst. prof. (Entomology) (Sr. Grade)

Mr. D. Kamaraj, Asst. prof. (Entomology)

PG Scholars, Department of Entomology (2024 & 2025 Batch)

Dr. M. JAWAHARLAL, Ph.D.,
Dean

Date: 06.10.2025



MESSAGE

It gives me immense pleasure to convey my warm greetings to the Department of Entomology, SRM College of Agricultural Sciences, SRMIST, for organizing the *National Symposium on Innovative Approaches for Integrated Pest Management (IPM Nexus 25)* on 10th October 2025.

Integrated Pest Management (IPM) has emerged as a vital approach in ensuring sustainable crop production, safeguarding the environment, and reducing reliance on chemical pesticides. This symposium provides a timely and meaningful platform for researchers, academicians, students, industry professionals, and policymakers to share innovative ideas, technologies, and field experiences that will contribute to the advancement of IPM strategies.

I am glad that the proceedings of abstracts have been compiled with an ISBN, which will serve as a valuable reference for scholars and practitioners across the country. Such documentation not only reflects the academic rigor of the symposium but also strengthens the visibility and impact of the knowledge shared.

I congratulate the organizing team for their dedicated efforts and extend my best wishes for the grand success of the symposium. I am confident that the deliberations and discussions will inspire new collaborations and contribute significantly to sustainable agricultural development.



Dr. M. Jawaharlal
Convener- IPM Nexus 25

IPM NEXUS 25: INDEX OF ABSTRACTS & FULL LENGTH PAPERS

S.NO	TITLE	PAGE NO
1.	Nanotechnologies for plant health management <i>Dr. K. S. Subramanian</i>	1
2.	Role of plant products in integrated pest management <i>Dr. S. Sridharan</i>	4
3.	Innovative ecological approaches for enhancing natural enemy diversity to strengthen conservation biological control <i>Dr. S. Jeyarani, and J.N.Prithiva</i>	5
4.	Emerging pests and their management: A case study with <i>Phenacoccus saccharifolii</i> of sugarcane <i>Dr. T. Ramasubramanian</i>	17
5.	Smart pest management through IoT and AI approaches <i>Dr. K. Rameash</i>	19
6.	Cyanobacterial extract as biopesticides against <i>Spodoptera frugiperda</i> in sustainable pest management <i>Sandhya Lakshmi P, Namirtha A P, Sandra Binu</i>	22
7.	Towards sustainable pest management: Integrating ecological engineering, biological control and agroecological strategies <i>Yogeshwaran B, Sandra Binu, Penugonda Bhavana</i>	23
8.	Sustainable crop protection: The future of eco-friendly pesticides <i>Sandra Binu, Yogeshwaran B, Namritha A P</i>	24
9.	Entomopathogenic fungi against wheat aphids <i>Namirtha A, Penugonda Bhavana and Sandhya Lakshmi</i>	25
10.	Eco-efficient pest management: Harmonizing environmental and economic goals in farming <i>Penugonda Bhavana, Sandra Binu, Yogeswaran B</i>	26
11.	Smart technologies for pollinator-safe integrated pest management <i>Berjin Viju M G, Muthukumar M and Melvin Joe M</i>	27
12.	Smart IPM: Integrating AI, IoT and remote sensing for real- time precision pest monitoring and decision support <i>Santhoshraj N and Murugan N</i>	28

S.NO	TITLE	PAGE NO
13.	Phyto-rhizosphere interception for nematode exclusion (P.R.I.N.E) A living bio-barrier for root pest management <i>Bhuvaneshwaran S and Murugan N</i>	29
14.	Ecological innovations for sustainable pest management <i>Kirubakaran G and Nisha R</i>	30
15.	Harnessing the black soldier fly (<i>Hermetia illucens</i>) for sustainable waste-to-wealth conversion <i>Ramazeame L, Athira Rajan, Karthik and Berjin Viju M G</i>	31
16.	Evaluation of bio-intensive IPM modules for management of insect-pest-pathogen complex in bhendi ecosystem <i>Rex B</i>	32
17.	Eco-friendly IPM strategies for major pests of okra <i>Ravanachandar A, Devika A and Dhilip Chakkaravarthy T</i>	33
18.	Study on physico-biochemical attributes of brinjal influencing resistance to shoot and fruit borer (<i>Leucinodes orbonalis</i> Guenee) <i>Rameshkumar D, Naveena E, Swarna Priya R and Savitha B K</i>	34
19.	Bioefficacy of novel insecticides against Banana scarring beetle, <i>Basilepta subcostata</i> (Jacoby) through laboratory bioassay <i>Kancharla Ratna Jyothi, Marri Keerthana and Sai Reddy M S</i>	35
20.	Less expensive, DIY method in reflected light microscopy to document insects: Concepts and procedures <i>Vinod Kumar S</i>	36
21.	Oil based formulation of <i>Metarhizium anisopliae</i> (Metchnikoff) Sorokin against <i>Helicoverpa armigera</i> (Lepidoptera: Noctuidae), <i>Plutella xylostella</i> (Lepidoptera: Plutellidae) and <i>Spodoptera frugiperda</i> (Lepidoptera: Noctuidae) <i>Nandha S and Jeyarani S</i>	37
22.	Entomopathogenic potential of soil-isolated entomopathogenic fungi <i>Metarhizium robertsii</i> strain TNAU ENTMR GYU 1 against Ash weevil (<i>Myloccerus subfasciatus</i>) <i>Monisha U and Shanmugam P S</i>	39

S.NO	TITLE	PAGE NO
23.	Evaluating aphid resistance in bread wheat (<i>Triticum aestivum</i> L.) genotypes under field and growth chamber conditions <i>Marri Keerthana, Priya, Kancharla Ratna Jyothi, and Sai Reddy M S</i>	40
24.	Genomic insights into the antifungal and plant growth promoting traits of <i>Pseudomonas plecoglossicida</i> NAN2 isolated from the rice rhizosphere <i>Nandhini Kuberan and Iyappan sellamuthu</i>	41
25.	Fruit fly diversity and resistance screening in mango and custard apple cultivars of Rayalaseema for sustainable pest management <i>Tirumala Geethika G, Sarada G, Ramaiah M, Nischala A and Rajeswari G</i>	42
26.	Advancements in ecological engineering for insect pest control <i>Sharulatha S and Muthukumar M</i>	43
27.	Ecological innovations in pest management: Entomological pathways to sustainable agriculture <i>Karthik S and Ramazeame L</i>	44
28.	Ecological pest management: Insects as catalysts for sustainable agriculture <i>Athira Rajan and Ramazeame L</i>	45
29.	Ecological innovations for sustainable pest management in Bhendi <i>Arun Prasad V and Nisha R</i>	46
30.	Eco-Friendly and climate-smart strategies for managing coffee berry borer in Andhra Pradesh <i>Urati Mahesh M, Venkateswara Reddy, Chaitanya K, Rajini D, Harika M and Lavanya A V N</i>	47
31.	Integrated ecological and digital approaches for managing tomato pinworm in Telangana <i>Urati Mahesh K, Chaitanya, A V N, Lavanya M, Venkateswara Reddy, Harika M and Rajini D</i>	48
32.	Bioefficacy of new Spirotetramat 150 OD against Aphid, <i>Aphis gossypii</i> in brinjal <i>Gajalakshmi M and Gunasekaran K</i>	49
33.	Advances in integrated pest management of San jose scale in mulberry: Synergistic role of biopesticides, predators, and cultural practices <i>Murugan Nagarajan and Vairam Namachivayam</i>	50

S.NO	TITLE	PAGE NO
34.	Smart horticultural production – A novel approach <i>Dhivya M, Usha G and Muruganantham M</i>	51
35.	Role of crop diversification in reducing pest outbreaks in cereal based cropping Systems <i>Jeyajothi R</i>	52
36.	IPM significance in safeguarding fresh water resources <i>Beslin Joshi J</i>	53
37.	Bio-digital pest management: Integrating living sensors (insects, microbes) with AI networks <i>Anitha D and Baradhan G</i>	54
38.	Exploration and cataloging of bee-attractive flora <i>Devi M</i>	55
39.	Development of a comprehensive floral calendar <i>Devi M</i>	56
40.	Digital traceability systems in poultry production using block chain <i>Preethi V</i>	57
41.	Digital pollinator–pest balance systems: AI models that optimize pest control without harming pollinators <i>Reshma M S and Baradhan G</i>	58
42.	Harnessing insect biodiversity for sustainable crop protection <i>Shukla Saurabh and Vishakha Sharma</i>	59
43.	<i>Momordica cymbalaria</i> leaf water extractives against the wrigglers of <i>Culex quinquefasciatus</i> Say <i>Yogapriya Adaikkan and Vikaash M</i>	60
44.	Digital frontiers in IPM: Precision management of Tea mosquito bug (<i>Helopeltis</i> spp.) in cashew <i>Gopu Sushma, Urati Mahesh, Kota Sahithi Chowdary and Repaka Harika</i>	61
45.	Ecological innovations for sustainable management of red palm weevil <i>Rhynchophorus ferrugineus</i> in coconut <i>Gopu Sushma and Urati Mahesh</i>	62

S.NO	TITLE	PAGE NO
46.	Soil microbiome engineering for resistance against rice brown planthopper (<i>Nilaparvata lugens</i>) <i>Repaka Harika, Kota Sahithi Chowdary and Gopu Sushma</i>	63
47.	IoT-Enabled smart traps for precision surveillance of tomato leaf miner (<i>Tuta absoluta</i>) <i>Kota Sahithi Chowdary, Repaka Harika and Gopu Sushma</i>	64
48.	Biocontrol potential of <i>Heterorhabditis bacteriophora</i> (Poinar) against rhinoceros beetle, <i>Oryctes rhinoceros</i> (L.) <i>Anjaly P V, Manu C R, Reji Rani O P, Gavvas R and Nisha M S</i>	65
49.	Semiochemical highways: Using plant–insect chemical signals for sustainable pest management <i>Dhilipan K S and Baradhan G</i>	66
50.	Trapping trouble: Sustainable control of rhinoceros beetle in coconut <i>Abishekkumar S and Selvanarayanan V</i>	67
51.	Synergistic effects of microbial consortia in integrated pest management <i>Kiruthiga N and Sharvesh S</i>	68
52.	Evaluating the dual action efficacy of lemongrass oil against <i>Galleria mellonella</i> <i>Vikaash M, Kanagarajan R, Manikandan E, Mohammad Ikram, Nishanthini K, and Yoga Priya A</i>	69
53.	Habitat manipulation strategies for sustainable management of rugose spiraling whitefly <i>Aleurodicus rugioperculatus</i> in coconut <i>Sumalatha B V, Selvaraj K and Kandan A</i>	70
54.	From harassing nature to harnessing nature – Towards ecofriendly pest management <i>Sriram K</i>	71
55.	RNAi-based biopesticide: A targeted approach for controlling the invasive giant african snail <i>Anjala Leelu Anil</i>	72
56.	Ecological innovations for sustainable management of fruit fly (<i>Bactrocera cucurbitae</i>) in cucurbits <i>Rajani D, Urati Mahesh, Suresh Kumar K and Harika M</i>	73

S.NO	TITLE	PAGE NO
57.	Bio inspired robotic model (Ornithoptera) <i>Ilakkiya V</i>	74
58.	Eco-friendly botanicals treatment and smart packaging: Sustainable management of <i>Rhyzopertha dominica</i> on stored wheat <i>Mekha M Prasad and Ghelani M K</i>	75
59.	Management of fruitflies (<i>Bactrocera dorsalis</i>) in mango ecosystem of A.P <i>Sravanthi G, Lakshmi B K M and Radha Rani K</i>	77
60.	Cold plasma seed treatment: An innovative approach for sustainable pest management in rice <i>Malini B and Vanitha J</i>	79
61.	<i>Thrips parvispinus</i> Karny (Thysanoptera: Thripidae): An emerging pest of chilli in Kerala <i>Anjana S A, Thania Sara Varghese, Santhosh Kumar</i>	80
62.	Resistance studies in green gram (<i>Vigna radiata</i>) parental lines and F1 hybrids against <i>Aphis craccivora</i> Koch <i>Darlin Prakash, R and Selvanarayanan V</i>	81
63.	Nanocarrier-mediated RNAi for sustainable fruit flies management <i>Manojkumar S, Kiruthiga N, Sharvesh S and Vigneswari G</i>	82
64.	Bioefficacy of entomopathogens against the sucking pests of okra <i>Dharani V and Kumar K</i>	83
65.	Harnessing nature for sunflower security: Eco-innovative IPM approaches against capitulum borer (<i>Helicoverpa armigera</i>) <i>Kiruthika M and Selvamuthukumaran T</i>	85
66.	Adjusting planting dates as an ecological innovation for sustainable pest management in rice <i>Yogapriya G and Kandibane Muthusamy</i>	86
67.	Microbiome-driven approaches to eco-friendly insect pest management <i>Arthi Mohan and Hemapriya J</i>	87
68.	Digital decision support tools and real time alert in integrated pest management <i>Rajaganapathy V and Raja Gopal V</i>	88

S.NO	TITLE	PAGE NO
69.	Computer vision and deep learning for pest counting and classification - A review <i>Raja Gopal V and Rajaganapathi V</i>	89
70.	Botanical bullets: Methanol and petroleum ether extracts of <i>Anisomeles malabarica</i> in the battle against tobacco leaf caterpillar – Eco friendly pest management <i>Sakthivel S and Kandibane M</i>	90
71.	Alkannin: A potential biopesticide for sucking pests <i>Vaishnavi Sri K T, Asha S and Nagajothi R</i>	92
72.	Bioefficacy of botanicals against rugose spiralling whitefly in guava: An ecological innovation for sustainable pest management <i>Navadeepchidambaram S and Kumar K</i>	93
73.	Eco-friendly integrated pest management of <i>Noorda moringae</i> and <i>Noorda blitealis</i> in moringa (<i>Moringa oleifera</i> Lam.) <i>Duvaraga Devi M and Muthuswami M</i>	94
74.	IPM and the future of food preservation: A sustainable approach <i>Chandraprabha S</i>	95
75.	Next-gen IPM in post-harvest management: Bridging science and trade <i>Prakash K</i>	96
76.	Bacterial endophytes as biocontrol agents: A sustainable approach to insect pest management <i>Sneha R And Rex B</i>	97
77.	Harnessing nature: Ecological innovations for sustainable pest management <i>Shopiya K and Rageshwari S</i>	98
78.	RNAi-mediated gene silencing in BPH – A review <i>Arul Jerlin J</i>	99
79.	Cold plasma treatment for storage pest management <i>Ambika S</i>	100
80.	A review of sooty mold on mango (<i>Mangifera indica</i> L.): Etiology, impact, and integrated management <i>Aruna G</i>	103

S.NO	TITLE	PAGE NO
81.	Socioeconomic determinants and behavioral drivers of digital IPM adoption tools among farming communities <i>Rajasekaran R, Jeevapriya A and Indumathy K</i>	104
82.	Host plant susceptibility index of okra genotypes against leafhopper and its association with nutrient content <i>Prithiva J N, Ganapathy N, Jeyarani S, Premalatha K and Rajesh R</i>	105
83.	Nature's shield for eco-friendly seed storage pest control <i>Vinothini Nedunchezhiyan</i>	106
84.	Influence of weather parameters on natural parasitization of <i>Helicoverpa armigera</i> by <i>Campoletis chloridiae</i> Uchida in chickpea ecosystem of the new alluvial zone of West Bengal <i>Sathees Kumar K and Meenambigai C</i>	107
85.	Bacteriophages – A novel biopreservative for vegetables <i>Ponvizhi Ramya V and Gunasekaran S</i>	108
86.	Bioacoustic surveillance systems for detecting insect for sustainable agriculture <i>Deepika D, Jeyajothi R and Thamizharasu T</i>	111
87.	The microbial shield: harnessing plant-microbiome interactions for pest resilience <i>Thamizharasu T, Jeyajothi R and Deepika D</i>	112
88.	Impact of spides on slaughter characteristics of broiler chickens from long-term stored eggs <i>Kowsalya N, Anandhi M, Jayanthi D, Shamsudeen P and Karthikeyan G</i>	114
89.	Influence of dietary azolla on growth and performance parameters in commercial broilers chicken <i>Kowsalya N, Sreyass K S and Karthikeyan G</i>	115
90.	Digital transformation in the pest management of fruit crops <i>Deepika V, Abishek and Akshaya P</i>	116
91.	Zero-residue integrated pest management (ZR-IPM) <i>Abinesh and Marimuthu S</i>	117
92.	Advancing integrated pest management through AI and machine learning innovations <i>Sathya Devi S</i>	118

S.NO	TITLE	PAGE NO
93.	Harnessing the insect microbiome: A new frontier in sustainable pest management <i>Swetha K, Sweeta Annie Mary A and Balasubramanian P</i>	119
94.	Capsaicin: A pungent molecule with potent insecticidal potential <i>Ilakiya T</i>	120
95.	Sun-powered guardians: Reducing pesticides, saving beneficial insects <i>Sanchana Dhas A K, Ramadass S and Priyanjena Chacko J</i>	121
96.	Harnessing nature: Ecological innovations for sustainable pest management <i>Yuvaraj N and Akshaya S B</i>	122
97.	Role of methyl eugenol-based pheromone traps in sustainable control of mango fruit fly <i>Mathiazhagan V</i>	123
98.	Innovative approaches for integrated pest management - A review <i>Nisha R</i>	124
99.	Integrated approaches for the control of insect in spices and cereals <i>Saran M S, Ashmitha K, Vinitha E and Sheela P</i>	128
100.	Association of endosymbionts with <i>Bemisia tabaci</i> of phylogenetically different groups <i>Rageshwari S, Buvaneswari S and Deivamani M</i>	129
101.	Studies on nutritional optimization for mycelial growth and biomass enhancement of entomopathogenic fungi <i>Ophiocordyceps sinensis</i> <i>Akshaya S B</i>	130
102.	Influence of light sources on sporulation of <i>Colletotrichum capsici</i> <i>Karpagavalli S</i>	131
103.	Eco-friendly suppression of guava whitefly through biological control agents within an IPM strategy <i>Vinothkumar G</i>	132

S.NO	TITLE	PAGE NO
104.	Field evaluation of <i>Trichogramma chilonis</i> and <i>Chrysoperla carnea</i> for the biological control of <i>Helicoverpa armigera</i> in tomato <i>Ramya P</i>	133
105.	Enhancing antixenosis-mediated resistance to yellow stem borer (<i>Scirpophaga incertulas</i> Walker) through intercropping and border crop strategies in irrigated rice ecosystems <i>Vairam Namachivayam and Murugan Nagarajan</i>	134
106.	Eco-friendly pest management through drip irrigation: A sustainable approach <i>Ramadass S, Rajeshkumar A, Saravana Kumar M, Rajasekar M and Muthukumar M</i>	141
107.	Biology and development of the black soldier fly (<i>Hermetia illucens</i> L.) reared on organic waste <i>Ramazeame L</i>	142
108.	Image processing for high-density mango farming: Detection of plant disease and fruit grading <i>Arunkumar J and Rajasekar P</i>	147
109.	Eco-intelligent trap cropping: A sustainable approach to smart pest management <i>Marimuthu Subramani, Sivakumar Kaliyanan and Jeyajothi Raman</i>	148
110.	Smart farming frontiers: Precision agriculture meets integrated pest management <i>Devi Vaishnavi V</i>	149
111.	Advancing pest control through RNAi gene silencing <i>Naresh J and Karpagavalli S</i>	150
112.	Plant-derived biofumigants: A safer and sustainable alternative for stored pest management <i>Sharulatha S, Muthukumar M and Anandhi S</i>	152

S.NO	TITLE	PAGE NO
113.	Black soldier fly larvae meal: Alternative dietary protein sources for the chicken <i>Sreyass K S and Karthikeyan G</i>	153
114.	Aphid mechanism of resistant in groundnut genotypes <i>Vanitha J and Mahendran R</i>	154
115.	Morphological and Molecular Characterization of Root-Knot Nematode, <i>Meloidogyne enterolobii</i>, infecting Guava (<i>Psidium guajava</i> L.) <i>Ashokkumar Natarajan, Sarrvesh lakshman, Rageshwari S and Gopu B</i>	155

NANOTECHNOLOGIES FOR PLANT HEALTH MANAGEMENT

Dr. K. S. Subramanian, Ph.D. (Canada)

Scientific Advisor, Coromandel International Ltd.,
Fmr. Director of Research & Founder Head, Center for Agricultural Nanotechnology,
Tamil Nadu Agricultural University, Coimbatore 641 003
Corresponding author Email: kss@tnau.ac.in / Phn: 98940 65449

ABSTRACT

Nanotechnology is a fascinating field of science dealing with manipulation of atom by atom and thus products and processes evolved from nano science are the most précised ones that are impossible to achieve by the conventional systems. Nano particles are very tiny, measuring a dimension of one-billionth of a metre (10⁻⁹ m) or one in 100 Crores, giving extensive surface area for nanomaterials to react and exhibit the reactions of the pathogens at the atomic scale. Nanotechnology is being exploited in a wide spectrum of agricultural sciences of which early detection of plant diseases, smart delivery of inputs and quarantine of plant diseases are quite significant.

Early detection of plant diseases is utmost essential to save the crop well before the symptoms exhibited. Quick on-site detection of plant pathogens using nano based kits such as Lateral Flow Immunoassay Devices (LFIDs) using gold nanoparticles, loop mediated isothermal amplification (LAMP) combined with LFD, Quantum dots (QDs), Nano barcode and Bio-imaging. These methods have high sensitivity, specificity, precision and speed of the pathogen detection that assist in high- throughput analysis and rapid response time to take suitable remedial measures.

Nano-Formulations are developed using active ingredients that are unequivocally demonstrated to possess pesticide value against devastating diseases. Persistence of agro chemicals in the initial stage of crop growth helps in bringing down the spread of the disease below economic threshold level and to have an effective control for a longer period. Hence, the persistence of pesticides is one of the most cost-effective and versatile means of controlling insect pests. In order to protect the active ingredient from the environmental conditions and to promote persistence, a nanotechnology approach “encapsulation (nano / micro)” can be used to improve the insecticidal value. Microencapsulation comprises nano-sized particles of the active ingredients being sealed by a thin-walled sac or shell (protective coating). Nanoencapsulation of pesticides allows for proper absorption of the chemical into the plants unlike the case of conventional formulations. Nano-encapsuaton of insecticides, fungicides or nematicides will help in producing nano-formulations which offer effective control of pests

while preventing residues in soil.

In addition to the encapsulated forms of insecticides, some of the nanoparticles are being used as effective strategy to protect the crops from damage by pests and diseases. Surface modified hydrophobic nanosilica has been successfully used to control a range of agricultural pests. The successful use of silver nanoparticles (Ag NPs) in diverse medical streams as antifungal and antibacterial agents has led to their applications in controlling phytopathogens. Ag NPs with broad spectrum of antimicrobial activity reduces various plant diseases caused by spore producing fungal pathogens. The effectiveness of Ag NPs can be improved by applying them well before the penetration and colonization of fungi within the plant tissues. The small size of the active ingredient effectively controls fungal diseases like powdery mildew. However, it was also observed that a very high concentration of nano-silica silver produced chemical injuries on the cucumber. The use of Ag NPs as an alternative to fungicides for the control of sclerotium forming phytopathogenic fungi was also investigated. Exposure of fungal hyphae to Ag NPs caused severe damage by the separation of layers of hyphal wall and collapse of hyphae. The efficacy of Ag NPs in extending the vase life of gerbera flowers was also studied and the results show inhibited microbial growth and reduced vascular blockage which increased the water uptake and maintained the turgidity of gerbera flowers. Apparently, the use of biocide containing polymeric nanoparticles for introducing organic wood preservatives and fungicides into wood products thereby reducing the wood decay was also studied. Among the nanoparticles, Ag NPs are widely used accounting for more than 30 per cent of the nano-based commercial products in the world.

“Smart Delivery Systems” for agriculture can possess timely controlled, spatially targeted, self regulated, remotely regulated, pre-programmed, or multi-functional characteristics to avoid biological barriers to successful targeting. Smart delivery systems can monitor the effects of delivery of nutrients or bioactive molecules or any pesticide molecules. In the smart delivery system, a small, sealed package carries the drug which opens up only when the desirable location or infection site of the human or animal system is reached. This would allow judicious use of antibiotics than otherwise would be possible. A molecular-coded “address label” on the outside of the package could allow the package to be delivered to the correct site in the body. Similarly, implanting nanoparticles in the plants could determine the nutrient status in plants and take up suitable remedial measures well before the malady causes yield reduction in crops. The fertilizer or irrigation requirement of crops can be scouted by nanotechnology. The exciting possibility of combining agricultural science and nanoscale technology into sensors holds the potential of increased sensitivity and therefore a significantly

reduced response- time to sense field problems. “Smart” delivery systems could contain on board chemical detection and decision-making capability for self-regulation that could deliver active chemical molecules or nutrients as needed. Remote activation and monitoring of intelligent delivery systems can assist agricultural growers of the future to minimize antibiotic and pesticide use.

Keywords: Nanoproducts, Ag NPs, Diseases, Sustainable pest management.

ROLE OF PLANT PRODUCTS IN INTEGRATED PEST MANAGEMENT

Dr. S. Sridharan, M.Sc., (Agri.) Ph.D.

Professor (Retd), Dept. of Agrl. Entomology, TNAU, Coimbatore – 641 003

Former Dean, PGPCAS, Namakkal; Former Principal, ACAR, Krishnagiri

Corresponding author Email: raji4sridharan@gmail.com

ABSTRACT

Plant-derived compounds are alternatives to synthetic pesticides with varied modes of action and contribute to sustainable agriculture. They can be insect antifeedants, repellents, deterrents, oviposition inhibitors showing direct insecticidal effects, hormonal imbalance causing growth abnormalities, besides, altering insect behaviour and physiology while generally being safer to nontarget organisms and the environment. The active phytochemical classes include alkaloids, terpenes, phenols, coumarins, flavonoids, quinone, glycosides, quassinoids, tannins, lignans, steroids, polyacetylene, and the like. For example, the members of Rutales contain limonoid, whereas the plants of Brassicaceae contain mustard oil in general, but a few species have cardenolides or cucurbitacins. Other families may diversify their deterrents, eg, non-protein amino acids (l-canavanine in jack bean), alkaloids, cyanogens, and iso flavonoids in Leguminosae.

The common botanical compounds with pesticidal activity that have been successfully isolated and commercialized include azadirachtin from neem (*Azadirachta indica*, *A. juss*), and pyrethrin from pyrethrum, *Chrysanthemum cinerariifolium* (Trevir). Other plants used in pest management include garlic *Allium sativum* L., turmeric *Curcuma longa* L., rosemary *Salvia rosemarinus* *spenn*, ginger *Zingiber officinale* Roscoe, and thyme *Thymus vulgaris* L.

With recent advances in research, the development of model tobacco and tomato plants with genetically engineered pyrethrin genes expressing higher levels of resistance is made possible. Likewise, engineering genes Bowman-Birk type protease inhibitors (BBIs) of cowpea or blackgram into plants like tobacco, rice, and wheat enhanced resistance against coleopteran and lepidopteran pests. The development of nanoemulsions of compounds of herbal plants and pesticidal plants makes pest management more effective and sustainable. The advent of drone technology offers a solution to pesticide drifting with precise application to site-specific fields.

Hence, integration of botanicals in integrated pest management ensures major benefits to the farmers, including food safety, reduced pest levels, improved quality of produce, realizing premium prices, and guaranteed market access.

Keywords: Pesticidal plants, Phytochemicals, non-target organisms, Environment, Sustainable agriculture.

**INNOVATIVE ECOLOGICAL APPROACHES FOR ENHANCING NATURAL
ENEMY DIVERSITY TO STRENGTHEN CONSERVATION BIOLOGICAL
CONTROL**

Dr. S. Jeyarani^{1*} and J N Prithiva²

¹Professor of Entomology, Department of Rice, Tamil Nadu Agricultural University,
Coimbatore 641 003

²Research Associate, Department of Agricultural Entomology, Tamil Nadu Agricultural
University, Coimbatore 641 003

*Corresponding author Email: jeyarani.s@tnau.ac.in

Abstract

Conservation Biological Control (CBC) is a cornerstone of sustainable pest management, emphasizing the protection and enhancement of naturally occurring predators, parasitoids, and entomopathogens. Innovative ecological approaches that enhance natural enemy diversity have emerged as effective and eco-friendly alternatives to chemical pesticides, offering stability and resilience to agroecosystems. Ecological engineering and habitat manipulation are among the most widely adopted strategies. Planting flowering strips and nectar-rich plants such as buckwheat (*Fagopyrum esculentum*), coriander (*Coriandrum sativum*), and marigold (*Tagetes* spp.) enhances the longevity and fecundity of parasitoids like *Trichogramma* and *Aphidius* species. In rice ecosystems, bund cropping with cowpea and sesame attracts coccinellids and chrysopids that suppress planthopper populations. Similarly, cotton intercropped with alfalfa enhances spider and lady beetle abundance, reducing whitefly and aphid infestations. Landscape diversification also plays a critical role in strengthening CBC. Hedgerows, non- crop habitats, and refuge patches provide shelter and alternative prey, ensuring the persistence of natural enemies during off-seasons. Push–pull strategies in maize, employing Desmodium as an intercrop and Napier grass as a border crop, not only suppress stem borers but also enhance parasitoid activity.

Besides, semiochemicals, particularly herbivore-induced plant volatiles (HIPVs) and synthetic kairomones, represent a rapidly advancing frontier in enhancing natural enemy efficiency. Compounds such as methyl salicylate (MeSA) and (Z)-3-hexenyl acetate have been shown to attract lady beetles, lacewings, and hoverflies to crop fields. Recent developments include slow-release dispensers and nanotechnology-based formulations that provide sustained emission of attractants, improving field- level efficacy. For instance, controlled release of MeSA has successfully enhanced predator recruitment in apple and soybean systems, leading to improved suppression of aphids and mites. Combining floral resources with semiochemical lures has been demonstrated to synergistically increase parasitoid foraging efficiency, creating multifunctional habitats that stabilize natural enemy populations.

The integration of ecological approaches with semiochemical tools represents a novel strategy for precision conservation biological control. These approaches reduce pesticide dependency, enhance biodiversity, and provide ecosystem services beyond pest regulation, including soil fertility and pollinator safety. The future of CBC lies in combining habitat manipulation, semiochemicals, and selective bio-rational inputs to develop climate-resilient, scalable, and sustainable pest management systems.

Introduction


Biological control is a cornerstone of sustainable pest management, offering an alternative to chemical insecticides by harnessing the potential of natural enemies and ecological processes. The use of biological control agents is one approach to reduce the undesirable ecological and health problems associated with overuse of chemical insecticides in agro ecosystems (Thoming, 2024). Biological control is a naturally occurring phenomenon and the biocontrol agents act on the crop pests and maintain them at certain densities. However, loss of biodiversity in intensive farming systems has undermined natural enemy communities and weakened pest suppression. To address these issues, historically, two primary directions have shaped biological control: conservation of naturally occurring beneficial organisms and augmentation through mass rearing and release (Gurr *et al.*, 2012). Over the last four decades, research has expanded beyond these ecological frameworks to include tritrophic interactions - linkages between plants, herbivores, and natural enemies mediated by infochemicals. These advances have illuminated how plants can “recruit bodyguards” by emitting volatile cues upon herbivore attack, enhancing the effectiveness of predators and parasitoids. Either it is conservation and enhancement or mass culture and release, a thorough knowledge of the crop ecosystem and the pests occurring is essential for successful implementation of the biocontrol programmes. Conservation biological control has emerged as a cornerstone of sustainable pest management. Rather than relying solely on external releases of natural enemies, it emphasizes the protection, enhancement and diversification of naturally occurring predators and parasitoids in agroecosystems.

This review integrates classical ecological approaches (habitat and cultural management, conservation strategies) with modern advances in infochemical-mediated pest regulation. By merging these perspectives, we highlight a hybrid approach that builds resilient agroecosystems capable of sustaining crop health with minimal chemical input.

Conservation and Enhancement of Natural enemies

Many avoidance strategies forming foundation of the management pyramid remains untapped. Integration of “avoidance” strategies into overall management programs improve

the efficiency of IPM. Agricultural intensification efforts should be designed based on ecological principles (Bommarco *et al.*, 2013) and the ecological intensification should be in favour of ecosystem services like natural enemies.



1. Selective application of chemical insecticides

Insecticides available are all potent and even in minute quantities (far less than that required against the target pest) is sufficient to eliminate many parasitoids, particularly micro-hymenopterans (Wilby & Thomas, 2002). If they are persistent, then the damage will be more. Hence, when a chemical insecticide has been used in a crop, the release of the natural enemies should be postponed until the toxicity has dissipated. Pest based selective use of insecticides like systemic / acaricides etc rather than broad spectrum can spare natural enemies. Also, use of chemicals should be avoided, when the activity of the natural enemy is high in a crop. However the decision has to be made taking into account other aspects of management.

2. Preservation of inactive stages of natural enemies

The preservation of inactive stages is most critical when there is a small reservoir or none at all of natural enemies outside the cropped area. At the initial establishment site of the parasitoids of the cereal leaf beetle, *Oulema melanopus* in Michigan, a high percentage of the major parasitoid, *Tetrastichus julis* overwintered in the soil of oat stubble. Hence a portion of the oat stubble was not ploughed until the parasitoids had emerged in the spring, resulting in a rapid build up of the parasitoid (Stehr and Haynes, 1972).

3. Crop Management Practices

Field-level management strongly affects trophic interactions. Reduced tillage preserves soil-dwelling predators, while intercropping and delayed sowing alter pest phenology to favour natural control (Bengtsson *et al.*, 2005). Crop rotation and diversified planting enhances habitat

heterogeneity, indirectly benefiting predator populations (Altieri, 1999). Conversely, pesticide use and nitrogen fertilization reduce natural enemy abundance (Koss *et al.*, 2005; Swift *et al.*, 1996). Thus, crop management should integrate ecological considerations alongside agronomic goals.

4. Avoidance of harmful cultural practices

Cultural control is a valuable part of managing pests but ploughing, hoeing, mowing, burning etc., can be equally harmful to natural enemies. The effects of these should be fully evaluated before such operations and harmful ones eliminated or modified. The strip harvesting of alfalfa in California has resulted in much greater survival of natural enemies and the hosts necessary to keep them active. In sunflower, early sowing favours the build up of important predators like *Chrysopa* and ladybird beetles (Arora *et al.*, 1998).

5. Maintenance of diversity and necessary hosts

Biodiversity enhances pest suppression through complementarity, whereby different natural enemies exploit pests at different times, life stages, or microhabitats (Straub *et al.*, 2008; Snyder, 2019). The maintenance of diversity is frequently a necessary part of many other measures since it may provide needed hosts, sources of food, over wintering sites, refuges etc. Diversity in itself is not essential but most natural enemies have evolved in communities much more diverse than crop communities, hence it is reasonable to expect the right kinds of diversity to be essential. Hambleton (1944) illustrated the value of crop diversity in the irrigated Canete Valley in Peru where the tobacco budworm *Heliothis virescens* reduced cotton yields by 30-40% following the planting of nearly entire valley to cotton. Arsenical insecticides had no effect and hence farmers were encouraged to plant winter crops such as corn, flax, beans and sweet potatoes which supported non-economic populations of various lepidoptera including *H. virescens*. These species provided the hosts necessary to maintain through the winter the population of predators and parasitoids that controlled *H. virescens* during the long summer growing season.

Whitman and Norduland (1994) found that parasitization of *Helicoverpa zea* (Boddie) eggs by *Trichogramma pretiosum* females was relatively high in plots of tomato but almost non-existent in adjacent plots of corn. It was found that the presence of volatile stimuli from tomato plants in corn fields resulted in increased rates of parasitization by *T. pretiosum*. Seasonal and diurnal differences among predators (e.g., day-active vs. nocturnal species) further extend pest suppression across temporal niches. However, risks such as intraguild predation and redundancy may reduce effectiveness (Rosenheim *et al.*, 1993; Straub and Snyder, 2006).

Structurally complex habitats and abundant alternative prey minimize these risks by dampening interference (Finke and Denno, 2002; Halaj and Wise, 2002).

6. Alternate hosts

The right life stages of the host may not be available at the right time or the right place to ensure the survival of natural enemies throughout the year. Manipulating the crop, planting different crops nearby, leaving some weeds or providing diversity in other ways is usually essential in providing for alternate hosts.

Alternative or non-pest hosts provide a continuous food source for natural enemies during pest-scarce periods, ensuring their survival and availability when pest outbreaks occur. These hosts can act as ecological reservoirs, supporting parasitoid and predator populations across cropping cycles. For example, parasitoids often rely on alternative lepidopteran or aphid hosts that persist in nearby vegetation, thereby stabilizing their populations. Such ecological engineering ensures that beneficial are not entirely dependent on target pests, making them more reliable allies in pest suppression. Besides, provision of artificial food also conserves beneficial fauna at times. For example, many parasitoids and predators require foods (nectar, pollen and honey dew) frequently not available in monocultures. In irrigated monoculture desert areas of California artificial honeydew and pollen in the form of food sprays have induced the early oviposition of naturally present Chrysopids and Coccinellids in treated fields resulting in significantly lower populations of aphids in alfalfa and of bollworms in cotton (Hagen *et al.*, 1970). The abundance of the predatory mite *Amblyseius hibisci* in avocado orchards in California was related to the availability of pollen, even when prey mites were absent. Although promising, the practical deployment of alternative hosts requires careful management to avoid creating refuges for secondary pests.

Few practices that could preserve natural enemies' population are tabulated below.

Crop	Natural enemy	Pest	Practice
Sweet pepper, ornamental crops	Syrphids, lacewings Predatory mites, <i>Orius laevigatus</i> , <i>O. majusculus</i> <i>O. insidiosus</i>	White flies, thrips, aphids	Plants providing pollen and plant sap as food sources for natural enemies like sweet alyssum, coriander, <i>Ricinus communis</i> and flowering ornamental
Cucumber, chrysanthemum	predatory mites <i>Amblyseius swirskii</i> & <i>Euseius scutalis</i> ,	Thrips, whitefly	Spraying or dusting artificial or natural food supplements onto the crop. i.e. corn pollen, apple pollen, <i>Typha latifolia</i> pollen

Cereal crops	Aphid parasitoids	Cereal aphids	Introducing non-crop plants harbouring the prey species
Chrysanthemum	Phytoseiid predatory mites	Spider mite	Applying yeast and sugars for astigmatic mites that are suitable prey for phytoseiid predatory mites
Rose plants	predatory mites	Spider mites	Providing oviposition sites and shelters: Sweet pepper was used by predatory mites for oviposition.
Cotton, wheat, tomato	<i>Orius</i> spp., lacewings, lady beetles	Aphids, thrips, leaf-feeders	Induced plant responses that attract and/or retain natural enemies
Wheat	<i>Orius</i> spp., lacewings, lady beetles	Aphids, thrips, leaf-feeders	Mitigation of pesticide side-effects by selecting pesticides that are compatible with natural enemies

1. Artificial shelters

Provision of artificial shelters in the crop helps the natural enemies to exert their influence on the insect pest. In North Carolina, substantial reduction of tobacco hornworms was achieved by predacious *Polistes* wasps following the erection of nesting shelters near field margins.

2. Reduction of undesirable predators

In some cases the desirable natural enemies are selectively fed by predators thereby affecting the efficiency of the parasitoid. The biological control of the walnut aphid, *Chromaphis juglandicola* by a different ecotype of *Trioxys pallidus* from Iran has been affected in some areas of California by selective predation of parasitized aphids in preference to unparasitized aphids by the Argentine ant, *Iridomyrmex humilis* (Frazer and van den Bosch, 1973).

3. Influence of Plant Traits and Breeding

Plant morphology and phytochemistry influence natural enemy performance. High trichome density in tomato, for example, increases egg parasitism by *Trichogramma* spp. Advances in plant breeding and genetic engineering now enable the incorporation of natural enemy-enhancing traits, such as rapid emission of herbivore-induced volatiles (Gould, 1998).

4. Control of honeydew feeding ants

Honeydew-feeding ants can make the biological control of honeydew-producing

species such as scales, mealybugs and aphids unsatisfactory because of interference with natural enemies (Styrsky and Eubanks, 2007)

5. Favourable temperature

Among the weather factors, temperature plays a major role in the effectiveness of parasitoids. The effect is more on parasitoids than on predators. Higher/extreme temperature usually affects the survival of certain delicate hymenopteran parasitoids. Besides this, beyond optima, the sex ratio of the progenies is also affected. In case of the egg parasitoid, *Trichogramma chilonis* beyond 31°C only males are produced (Prasad and Prasad, 2023).

6. Role of Semi-natural Habitats

Semi-natural habitats—hedgerows, meadows, fallows, and woodlots—provide stable resources such as pollen, nectar, and alternative prey, supporting diverse enemy communities (Marshall, 2004). These habitats act as refuges and overwintering sites, enabling enemy populations to persist when crops are unavailable (Pickett *et al.*, 2000; Rusch *et al.*, 2010). Studies showed that proximity and connectivity of these habitats enhance enemy spillover into crops, thereby stabilizing pest suppression (Bianchi *et al.*, 2006).

7. Avoidance of dust

Dust created in the farms by farm operations affect the efficiency of the natural enemies. Road dust and other dusts have been shown to drastically reduce the effectiveness of some natural enemies in California while having little effect on the hosts.

Innovations in Conservation Biological Control (CBC)

Recent innovations combine ecological knowledge with practical farming strategies. CBC efforts increasingly aim to deliver multiple ecosystem services like pollination, soil fertility, and climate resilience alongside pest suppression (Gurr *et al.*, 2016; Karp *et al.*, 2018). Plant volatiles, pheromones, and synthetic attractants can selectively enhance natural enemy foraging efficiency (Pickett and Bugg, 1998; Gontijo, 2019). Refuge strips combining bunch grasses for ground predators and floral resources for parasitoids exemplify ecological engineering that fosters complementary natural enemies (Snyder, 2019). Leaving crop residues increases habitat heterogeneity, reduces intraguild predation and enhances prey availability (Rowen *et al.*, 2019).

Chemical Ecology / Semiochemicals in Conservation Biological Control

Chemical ecology explores how chemical signals mediate interactions between organisms, particularly between plants, pests, and natural enemies. Semiochemicals such as

kairomones, allomones, and synomones play a vital role in enhancing host-finding efficiency of parasitoids and predators. For instance, herbivore-induced plant volatiles (HIPVs) can attract parasitoids like *Trichogramma* spp. to infested plants, thereby improving natural enemy recruitment. Understanding these complex chemical- mediated interactions has enabled researchers to manipulate agroecosystems in ways that boost biological control while reducing pesticide dependency.

Natural enemies may also respond to odours like sex pheromones or alarm pheromones produced by their prey / host species. Compounds such as methyl salicylate (MeSA) and (Z)-3-hexenyl acetate have been shown to attract lady beetles, lacewings, and hoverflies to crop fields. Recent developments include slow-release dispensers and nanotechnology-based formulations that provide sustained emission of attractants, improving field-level efficacy. For instance, controlled release of MeSA has successfully enhanced predator recruitment in apple and soybean systems, leading to improved suppression of aphids and mites. Some of the examples are tabulated below.

Crop	Semio-chemicals	Effect	Reference
Cotton	Pink bollworm mating- disruption chemicals	Predators increased by 3-fold	(McVeigh <i>et al.</i> , 1983)
Apple	Iiridodial-methyl salicylate	<i>Chrysopa oculata</i> (8-145 nos.) & <i>C. nigricornis</i> (86-446 nos.) than control trees (0-3 & 0-7nos.)	(Jones <i>et al.</i> , 2011)
Chinese cabbage	E-β-farnesene lures	9-41 parasitised aphids/20 plants compared to 5-19 parasitised aphids in controls	(Cui <i>et al.</i> , 2012)
Turnip	Methyl salicylate lure	More <i>Diadegma semiclausum</i> (1.6-7.2 wasps) than control (1.4-6.4 wasps)	(Orre <i>et al.</i> , 2010)
Citrus	Lavandulyl senecioate lures	19-51 <i>Anagyrus</i> sp. from citrus mealybug, <i>Planococcus citri</i> than in control with 0-20 wasps	(Franco <i>et al.</i> , 2011)
Broccoli	Treated with allyl isothiocyanate	Parasitism in aphid increased from 8.5 to 22.5%	(Titayavan & Altieri 1990)

Challenges and Research Gaps

While laboratory studies have elucidated mechanisms of tritrophic signaling, translating them to complex field environments remains challenging. Interactions among multiple natural enemies, variability in volatile release across cultivars, and climatic influences demand long-term, landscape- scale studies (Turlings *et al.*, 1995).

Innovative Hybrid Approaches

Overall, integrating ecological and infochemical approaches offers a synergistic framework for biological control. Key hybrid strategies include:

1.Habitat Manipulation + Volatile Recruitment: Designing diverse crop landscapes while deploying synthetic HIPVs or natural plant volatiles to attract natural enemies.

2.Breeding for Multi-Trait Resistance: Developing crop cultivars with both direct resistance (secondary metabolites, trichomes) and indirect resistance (enhanced volatile emission).

3.Landscape-Level Push–Pull Systems: Combining cultural diversity with semiochemical traps to manage pests across heterogeneous fields.

4.Biotechnology and Smart Delivery: Engineering plants to release volatiles at low infestation levels, coupled with slow-release dispensers of synthetic kairomones. Such integrative models not only strengthen natural enemy efficacy but also build resilience against pest adaptation and climate change.

Conclusion

When we decide to employ biocontrol as a component of IPM, a thorough knowledge of the crop ecosystem, weather conditions, pests occurring on the plants, their life cycle, already occurring natural enemies, crop damage level, etc. is necessary. Only after careful consideration of the prevailing crop condition and also the market situation we can employ appropriate biocontrol agent for obtaining maximum benefits.

Conservation biological control represents an ecologically sound and sustainable alternative to chemical-intensive pest management. The combined use of alternative hosts, chemical ecology, and semiochemicals has demonstrated significant potential in enhancing the abundance and diversity of natural enemies. By integrating these innovative approaches, agricultural systems can become more resilient, productive, and environmentally sustainable. The success of these strategies will ultimately depend on interdisciplinary research, farmer engagement, and policy support aimed at reducing chemical reliance and promoting ecological intensification.

Future Perspectives

Despite promising advances, several challenges remain in strengthening conservation biological control. The impact of climate change on multitrophic interactions is poorly understood, and the long- term effectiveness of alternative host strategies in different

agroecological contexts requires further study. There is also a need for molecular tools to trace trophic linkages and identify cryptic species of natural enemies. Farmer adoption can be limited by a lack of awareness or short-term economic considerations. Future research should focus on developing scalable, farmer-friendly technologies that combine ecological engineering with innovative chemical tools.

Selected References

- Altieri, M.A. 1999. The ecological role of biodiversity in agroecosystems. *Agriculture, Ecosystems & Environment*, **74**: 19–31.
- Arora, R., Kaur, S. and Dhillon, A.S. 1998. Manipulation of date of sowing for natural control of insect pests in sunflower agroecosystem. In: G.S. Dhaliwal, M.S. Randhawa, R. Arora and A.K. Dhawan (eds.). *Ecological Agriculture and Sustainable Development*. Vol.1. Indian Ecological Society and Centre for Research in Rural and Industrial Development, Chandigarh. pp. 449- 456.
- Bengtsson, J., Ahnstrom, J. and Weibull, A.C. 2005. The effects of organic farming on biodiversity and abundance: a meta-analysis. *Journal of Applied Ecology*, **42**: 261–269.
- Bianchi, F.J.J.A., Booij, C.J.H. and Tscharntke, T. 2006. Sustainable pest regulation in agricultural landscapes: a review on landscape composition, biodiversity and natural pest control. *Proceedings of the Royal Society B*, **273**: 1715–1727.
- Bommarco, R., Kleijn, D. and Potts, S.G. 2013. Ecological intensification: harnessing ecosystem services for food security: A Review. *Trends in Ecology and Evolution*, **28**(4): 230-238.
- Finke, D.L. and Denno, R.F. 2002. Intraguild predation diminished in complex-structured vegetation: implications for prey suppression. *Ecology*, **83**: 643–652.
- Frazer, B. D. and van den Bosch, R. 1973. Biological control of the walnut aphid in California: The interrelationship of the aphid and its parasite. *Environmental Entomology*, **2**(4): 561–568.
- Gould, F. 1998. Sustainability of transgenic insecticidal cultivars: integrating pest genetics and ecology. *Annu. Rev. Entomol.*, **43**: 701–726.
- Gurr, G.M., Wratten, S.D. and Snyder, W.E. 2012. *Biodiversity and Insect Pests: Key Issues for Sustainable Management*. Wiley-Blackwell, 360pp
- Hagen, K.S., Sawall, Jr., E.F. and Tassan, R.L. 1970. The use of food sprays to increase effectiveness of entomogenous insects. *Proc. Tall Timbers Conf. Ecol. Anim. Control Habitat Manage.*, **2**: 59-82.

Halaj, J. and Wise, D.H. 2002. Impact of a detrital subsidy on trophic cascades in a terrestrial food web.

Ecology, **83**: 3141–3151.

Hambleton, E.J. 1944. *Heliothis virescens* as a Pest of Cotton, with Notes on Host Plants in Peru. *Journal of Economic Entomology*, **37**(5): 660–666.

Koss, A.M., Jensen, A.S., Schreiber, A., Pike, K.S. and Snyder, W.E. 2005. Comparison of predator and pest communities in Washington potato fields treated with broad-spectrum, selective, or organic insecticides. *Environmental Entomology*, **34**: 87–95.

Marshall, E.J.P. 2004. Agricultural landscapes: field margin habitats and their interaction with crop production. *Journal of Crop Improvement*, **12**: 365–404.

Pickett, C.H. and Bugg, R.L. 1998. Enhancing Biological Control: Habitat Management to Promote Natural Enemies of Agricultural Pests. University of California Press.

Prasad, R.P. and Prasad, Y.S. 2023. Biotic and abiotic factors that affect parasitism in *Trichogramma*

spp. *Egyptian Journal of Biological Pest Control*, **33**(1): 56.

Rosenheim, J.A., Kaya, H.K., Ehler, L.E., Marois, J.J. and Jaffee, B.A. 1993. Intraguild predation among biological control agents: theory and evidence. *Biological Control*, **3**: 303–335.

Rusch, A., Valantin-Morison, M., Sarthou, J.P. and Roger-Estrade, J. 2010. Biological control of insect pests in agroecosystems: effects of crop management, farming systems and seminatural habitats at the landscape scale. *Agronomy for Sustainable Development*, **30**: 367–389.

Snyder, W.E. 2019. Give predators a complement: Conserving natural enemy biodiversity to improve biocontrol. *Biological Control*, **135**: 73–82.

Stehr, F.W. and Haynes, D.L. 1972. Establishment in the United States of *Diaparsis carnifer*, a larval parasite of the cereal leaf beetle. *J. Econ. Entomol.*, **65**: 405–417.

Straub, C.S. and Snyder, W.E. 2006. Species identity dominates the relationship between predator biodiversity and herbivore suppression. *Ecology*, **87**: 277–282.

Straub, C.S., Finke, D.L. and Snyder, W.E. 2008. Are the conservation of natural enemy biodiversity and biological control compatible goals? *Biological Control*, **45**: 225–237.

Styrsky, J.D. and Eubanks, M.D. 2007. Ecological consequences of interactions between ants and honeydew-producing insects. *Proceedings of the Royal Society B: Biological Sciences*, **274**(1607): 151–164.

- Swift, M.J., Izac, A.M.N. and van Noordwijk, M. 1996. Biodiversity and ecosystem services in agricultural landscapes—are we asking the right questions? *Agriculture, Ecosystems & Environment*, **58**: 5–17.
- Thoming, G. 2024. Conservation biological control on-farm, using VOCs combined with maintained natural border vegetation for a more unerring and resilient biological control of aphids in cereals. *Frontiers in Horticulture*, **3**: 1440410.
- Whitman, D.W. and Nordlund, D.A. 1994. Plant chemicals and natural enemies of herbivorous arthropods. In: *Functional Dynamics of Phytophagous Insects*. Oxford & IBH.
- Whitman, D.W. and Nordlund, D.A. 1994. Plant chemicals and the location of herbivorous arthropods by their natural enemies. In: T.N. Ananthakrishnan (ed.) *Functional Dynamics of Phytophagous insects*. Oxford & IBH Publishing Co. Pvt. Ltd., New Delhi. pp.133-159.
- Wilby, A. and Thomas, M.B. 2002. Natural enemy diversity and pest control: patterns of pest emergence with agricultural intensification. *Ecology Letters*, **5**: 353–360.

**EMERGING PESTS AND THEIR MANAGEMENT: A CASE STUDY WITH
Phenacoccus saccharifolii OF SUGARCANE**

Dr. T. Ramasubramanian

Principal Scientist (Entomology), Division of Crop Protection, ICAR-Sugarcane Breeding
Institute, Coimbatore, Tamil Nadu, India

Corresponding author Email: tramasubbu@gmail.com

ABSTRACT

Nowadays minor pests have become major ones and threaten the cultivation of economically important crops worldwide. It is very challenging to manage such emerging pests as one has to study almost all the entomological aspects of the pests including, but not limited to their basic taxonomy, biology, ecology, host plant resistance, biological control and insecticide toxicology. Further, appropriate management strategies have to be developed within no time as the new pests are more devastating in nature. The mealybug, *Phenacoccus saccharifolii* (Green) (Hemiptera: Pseudococcidae), a lesser-known pest, has become a major threat to cane cultivation in India and Tamil Nadu, in particular. Though this species has been reported to occur on sugarcane in as early as 1942 in the Bihar State of India, it had never been considered as a pest of sugarcane until 2021. It has now emerged as a major pest of sugarcane warranting insecticidal interference to protect the crop from being perished. Since the infestation by *P. saccharifolii* at initial stage is generally restricted between -1 and -3 leaves of sugarcane, it is often unnoticed by the farmers and cane developmental personnel. If unnoticed at early stage, the central leaf whorl becomes rotten and the apical growth gets arrested leading to stunted growth. A novel and reliable method of estimating yield loss due to this pest has been developed and the same has been employed to assess the loss in the ratoon crop of cv. Co 11015. The yield loss due to this pest in cv. Co 11015 has been estimated to be 7.1 tonnes/ha at 10% level of incidence. Almost all the entomological aspects of this pest have been studied at the ICAR-SBI and the pest is now manageable with the adoption of appropriate plant protection measures. A study on the detailed biology of this pest revealed that the male and female individuals could complete their life cycle in 21.9 ± 1.2 and 39.2 ± 1.6 days, respectively. Like other Pseudococcids, males undergo complete metamorphosis with pupal stage, while females undergo incomplete metamorphosis without a pupal stage. There were two and three instars in male and female sexes, respectively. The mean fecundity was observed to be 281 ± 90 eggs/female and the eggs were laid in an ovisac during the oviposition period of 6.2 ± 1.1 days. Parthenogenetic reproduction with amphitoky was also observed in this species. A three years extensive and intensive survey indicated that the plant crop suffers more as compared to the ratoon crop. Among the germplasm screened for their reaction against the

mealybug, cv. CoV 09356 and Co 86032 were observed to be the most susceptible and least susceptible varieties, respectively. The solitary parasitoid, *Leptomastix sylvae* (Encyrtidae: Hymenoptera) was found to regulate the pest population to a considerable extent in the surveyed fields. The parasitism by *Aenasius hayati* has been found to be suppressed to a greater extent by the hyper parasitoid, *Promuscidea unfasciativentris* (Eriaporidae: Hymenoptera). *L. sylvae* was also found to reproduce parthenogenetically with the sex ratio skewed to male. Thiamethoxam was observed to be more toxic to this mealybug species. It was also observed to be harmful to the parasitoid, *L. sylvae*. All the granular insecticides evaluated for their efficacy against *P. saccharifolii* were found to be effective with more than 80% reduction in the population after two applications at their respective recommended doses. An integrated approach with the cultivation of less susceptible varieties like Co 86032, promotion of natural parasitism by dominant species like *L. sylvae* and need-based application of effective insecticides like thiamethoxam is being suggested for successful and sustainable management *P. saccharifolii* in sugarcane crop ecosystem.

Keywords: Sustainable pest management, Integrated approaches.

SMART PEST MANAGEMENT THROUGH IOT AND AI APPROACHES

Dr. K. Rameash

Principal Scientist (Agricultural Entomology), ICAR - Central Institute for Cotton Research,
Regional Station, Coimbatore

Corresponding author Email: krameash@gmail.com

ABSTRACT

The agricultural sector faces unprecedented challenges due to the rising global population, erratic climate fluctuations, and the growing pressure to ensure food security in an environmentally responsible manner. In recent years, remarkable advances in smart pest management have been driven by the convergence of Internet of Things (IoT), artificial intelligence (AI), and remote sensing (RS) technologies. Modern pest monitoring increasingly relies on robust sensor networks, unmanned aerial vehicles (UAVs), GPS-enabled robotics, and cloud-based analytics, which collectively enable early detection of insect and disease outbreaks.

AI-powered computer vision and deep learning algorithms, in particular, provide highly accurate image-based identification and forecasting of key pests and diseases, moving beyond conventional manual scouting. Recent commercial deployments—such as precision spray systems that leverage AI for targeted herbicide application—demonstrate significant reductions in chemical inputs and environmental impact. Additionally, public–private initiatives have begun creating large, accessible datasets and scalable IoT infrastructures to support real-time, field-level pest detection and adaptive management decisions. Despite challenges related to data quality and limited digital access in rural regions, the rapid adoption of AI-enabled solutions in crop protection is already transforming integrated pest management (IPM) strategies by enhancing prediction, optimizing input use, and improving sustainability on a global scale (Nyakuri *et al.*, 2025). These developments, reinforced by scalable IoT frameworks and real-time analytics, are fostering data-driven advances in pest monitoring, forecasting, and management, thereby ensuring higher productivity while minimizing input costs and safeguarding agro-ecosystem health.

Advances in micro-fabrication and embedded systems have also enabled the development of miniature electronic devices for recording plant health parameters. Radar, thermal infrared imaging, and high-resolution RGB (red, green, blue) and hyperspectral cameras are increasingly being deployed for real-time crop monitoring, which indirectly supports the detection of pest- or disease-induced stress. A notable example is the AI-based smart pheromone trap developed by the ICAR – Central Institute for Cotton Research to

overcome the limitations of conventional pheromone traps for monitoring the pink bollworm (PBW). In this system, trap catches of PBW adult moths are recorded as digital images and transmitted to a remote server via the GSM communication network, along with weather parameters recorded at fixed intervals. The remote server stores the data in cloud storage, where a machine learning algorithm (YOLO – You Only Look Once) identifies PBW moths and counts them in each time-stamped image. The processed information, including insect counts, trap images, and corresponding weather data, is then transmitted to end users via Android-based mobile and Windows-based desktop applications. By integrating real-time trap data with weather parameters, this system enables a better understanding of pest dynamics across wider areas, contributing to more reliable pest forewarning systems and improved management practices in cotton (Rameash, 2024).

AI tools are also being widely employed for insect pest identification, population counting, and the prediction of pest spread in major crop ecosystems. Machine learning algorithms and computer vision techniques play a central role in these applications, particularly for the classification and detection of plant diseases. Key image features such as color, texture, and edges are extracted from diseased leaf or insect pest samples, and the segmented images are processed through classification algorithms such as Support Vector Machines (SVM), Convolutional Neural Networks (CNN), and Artificial Neural Networks (ANN) (Ebrahimi *et al.*, 2017).

Notable applications include disease identification in rice and cassava (Hassan and Maji, 2022), citrus disease classification using leaf images (Syed-AbRahman *et al.*, 2022), and viral disease diagnosis in pulses (Joshi *et al.*, 2021), all of which have demonstrated promising results using CNN- based object detection techniques. In cotton, CNN models have been developed to classify 13 insect pests, including *Helicoverpa armigera*, *Aphis gossypii*, and *Anthonomus grandis*. Similarly, diseases such as cercospora leaf spot (SVM), bacterial leaf spot (ANN), fusarium wilt (SVM), and ascochyta blight (CNN) have been detected with accuracies ranging from 80% to 96% (Toscano-Miranda *et al.*, 2022).

The integration of IoT and AI holds considerable potential to address critical challenges faced by farmers, such as the judicious use of pesticides, reducing dependence on manual labour for pest monitoring, and enhancing crop productivity while safeguarding environmental and soil health. Nevertheless, several constraints remain. AI systems require extensive spatial and temporal datasets for model training and precise prediction. Developing location-specific algorithms demands substantial initial investment, concentrated research and development, and significant operational and maintenance costs. Moreover, poor internet connectivity and

bandwidth limitations in rural and remote regions may restrict the deployment and adoption of IoT- and AI-based solutions. Despite these hurdles, rapid technological progress and growing investment by agricultural enterprises are accelerating the adoption of AI-driven tools for plant health management, offering promising support for progressive farming practices.

Keywords: AI technology, Sustainable pest management, pest and diseases.

References:

- Ebrahimi M A, Khoshtaghaza M H, Minaei S, Jamshidi B 2017. Vision-based pest detection based on SVM classification method. *Computers and Electronics in Agriculture* 137: 52-58.
- Hassan S M and Maji A K 2022. Plant Disease Identification Using a Novel Convolutional Neural Network. *IEEE Access* 10: 5390-5401
- Joshi R C, Manoj K, Dutta M K, Srivastava A, Choudhary, N 2021. VirLeafNet: Automatic analysis and viral disease diagnosis using deep-learning in Vigna mungo plant. *Ecological Informatics* 61 : 190-197.
- Rameash K and Prasad, Y G. 2024. Artificial Intelligence and Area Wide Approaches in Tackling the Dreaded Pest of Cotton. *Cotton Statistics and News* 34 (1): 1-4.
- Nyakuri J P, Nkundineza C, Gatera O. 2025. AI and IoT-powered edge device optimized for crop pest and disease detection. *Sci Rep* 15, 22905 <https://doi.org/10.1038/s41598-025-06452-5>.
- Syed-Ab-Rahman S F, Hesamian M H, Prasad M 2022. Citrus disease1 detection and classification using endto-end anchor-based deep learning model. *Applied Intelligence* 52 : 927-938
- Toscano-Miranda R, Toro M, Aguilar J, Caro M, Marulanda A, Trebilcok A 2022. Artificial-intelligence and sensing techniques for the management of insect pests and diseases in cotton: A systematic literature review. *The Journal of Agricultural Science* 160 : 16–31

CYANOBACTERIAL EXTRACT AS BIOPESTICIDES AGAINST *Spodoptera frugiperda* IN SUSTAINABLE PEST MANAGEMENT

Sandhya Lakshmi P*, Namirtha A P and Sandra Binu

PG Scholar, Department of Entomology, SRM College of Agricultural Sciences,
Baburayanpettai, Chengalpattu, Tamil Nadu - 603 201.

*Corresponding author Email: sp6007@srmist.edu.in

ABSTRACT

In intensified agriculture, the long-term use of chemical pesticides to control pests has led to ecological imbalance, pesticide resistance. Ecological solution being viable alternative to manage pest in a sustainable way. Fall armyworms, scientifically called *Spodoptera frugiperda*, are polyphagous pests causing severe damage on more than eighty crop species worldwide and known to have surprising resilience to chemical approaches. In this study, food overlay bioassay is conducted to evaluate the effectiveness of cyanobacterial extracts, from *Spirulina* sp. and *Nostoc muscorum*, against the second instar larvae of *S. frugiperda*. An assessment was carried out on the extracts of *N. muscorum* obtained using hexane, petroleum ether, ethanol, and methanol. These extracts were subjected to evaluation for their lethal concentration (LC50) values and their effectiveness on biological parameters. It was noted that the hexane and petroleum ether extracts of *N. muscorum* showed the highest toxic effect, which was known by the lowest LC50 values also significant negative impact on larval and pupal duration, percent pupation, pupal and adult malformation, pupal weight, and further reduced adult emergence, fecundity, and longevity. Similar effects were exerted by *Spirulina* extracts, though the changes were not so. The results impart metabolites of cyanobacteria as a considerable as eco friendly alternatives for the green control of *S. frugiperda*, also providing an approach for the design and the utility of cyanobacteria-based biopesticides in integrated pest management approaches.

Keywords: Ecological pest control, Integrated pest management, *Spodoptera frugiperda*, Biopesticides, Cyanobacterial extracts

**TOWARDS SUSTAINABLE PEST MANAGEMENT: INTEGRATING
ECOLOGICAL ENGINEERING, BIOLOGICAL CONTROL AND
AGROECOLOGICAL STRATEGIES**

Yogeshwaran B*, Sandra Binu and Penugonda Bhavana

PG Scholar, Department of Entomology, SRM College of Agricultural Sciences, SRM
Institute of Science and Technology, Baburayanpettai, Chengalpattu,
Tamil Nadu - 603 201.

*Corresponding author Email: yb3697@srmist.edu.in

ABSTRACT

The pursuit of sustainable agriculture has highlighted the urgent need to reduce reliance on synthetic pesticides and strengthen ecologically based pest management approaches. Across disciplines, ecological engineering, integrated pest management (IPM), and biologically based strategies converge on enhancing natural regulations of pest populations while safeguarding ecosystem stability, food security and farmer livelihoods. Push-Pull systems, conservation of natural enemies, and the strategic use of biological control agents represent promising pathways, yet their is challenged by inconsistent IPM implementation, limited farmer participation, and the uncertainties imposed by climate change. Recent advances in agroecological crop protection emphasize the integration of ecological principles at the field, farm and landscape levels, where biological control, habitat management and judicious chemical use can operate synergistically. Future progress requires adaptive management under changing climates, stronger farmer-centered approaches and interdisciplinary collaboration to achieve resilient, sustainable and scalable pest management systems.

Keywords: Ecological Engineering, Integrated Pest Management, Biological control, Agroecology, Climate change, Sustainable crop protection.

SUSTAINABLE CROP PROTECTION: THE FUTURE OF ECO-FRIENDLY PESTICIDES

Sandra Binu*, Yogeshwaran B and Namritha A P

PG Scholar, Department of Entomology, SRM College of Agricultural Sciences,
Baburayanpettai, Chengalpattu, Tamil Nadu - 603 201.

*Corresponding author Email: sb3950@srmist.edu.in

ABSTRACT

Modern agriculture's heavy reliance on synthetic pesticides has led to a range of serious problems—soil and air pollution, disruption of ecosystems, health concerns for humans, and the growing issue of pest resistance. These challenges highlight the pressing need for safer, more sustainable alternatives. This study explores the potential of eco-friendly pesticides made from gadung tubers (*Dioscorea hispida*), a plant long valued for its natural bioactive compounds. Using a combination of literature review, expert interviews, and field trials in Gowa, Indonesia, the research found that gadung-based pesticides effectively controlled key agricultural pests such as brown planthoppers and cocoa pod suckers, with mortality rates above 70%. Encouragingly, these biopesticides caused little harm to beneficial organisms or the surrounding environment. Their effectiveness lies in naturally occurring compounds like diosgenin and alkaloids, which disrupt the nervous systems of target pests, leading to paralysis and death. The study also points to the importance of developing improved formulations for greater stability and ease of use, along with farmer training programs to encourage adoption. When paired with organic farming practices, such eco-friendly innovations can significantly reduce dependence on harmful chemicals, protect biodiversity, and contribute to building a more sustainable, resilient, and productive agricultural future.

Keywords: *Dioscorea hispida* (Gadung tubers), Biopesticides, Paralysis, Diosgenin, Alkaloids

ENTOMOPATHOGENIC FUNGI AGAINST WHEAT APHIDS

Namirtha A P*, Penugonda Bhavana and Sandhya Lakshmi

PG Scholar, Department of Entomology, SRM College of Agricultural Sciences,
Baburayanpettai, Chengalpattu, Tamil Nadu - 603 201.

*Corresponding author Email: na0634@srmist.edu.in

ABSTRACT

Pests making farmers often rely on chemical pesticides. Even though they are effective initially, these inputs have created long-term problems like degradation, ecological imbalance, resistance. To tackle this, ecological innovations offer sustainable alternatives that harness natural processes for pest regulation. Aphids such as *Rhopalosiphum padi* and *Schizaphis graminum* are the most destructive pests of wheat across the world, these pest are alternatives to chemical insecticides. By this study, it is clear that the pathogenic potential of two entomopathogenic fungi, *Beauveria bassiana* and *Metarhizium anisopliae*, under both lab and field conditions. In Lab, bioassays demonstrated that *B. bassiana* at a concentration of 1.05×10^{13} conidia/ml induced mortality rates of 83.4% in *R. padi* and 79.4% in *S. graminum* after seven days, showing outperforming *M. anisopliae*. In Field trials, involving three foliar applications at 15-day intervals, confirmed the superior performance of *B. bassiana*, achieving 69.3% and 66.3% mortality in *R. padi* and *S. graminum*, respectively. These findings are very strong biocontrol potential of *B. bassiana* against wheat aphids and support its integration into eco-friendly pest management strategies and a good alternative to synthetic insecticides.

Key words: Entomopathogenic fungi, Biological control, Pest management, Sustainable, Eco-friendly.

**ECO-EFFICIENT PEST MANAGEMENT: HARMONIZING ENVIRONMENTAL
AND ECONOMIC GOALS IN FARMING**

Penugonda Bhavana*, Sandra Binu and Yogeswaran B

PG Scholar, Department of Entomology, SRM College of Agricultural Sciences,
Baburayanpettai, Chengalpattu, Tamil Nadu - 603 201.

*Corresponding author Email: pb4685@srmist.edu.in

ABSTRACT

Sustainable pest management plays a vital role in modern agriculture by protecting crops while minimizing environmental harm and maintaining economic viability. Conventional pest control methods, largely dependent on chemical pesticides, have raised concerns due to their negative impacts on ecosystems, human health, and biodiversity. This review examines sustainable pest management approaches that incorporate ecological principles and economic factors. It explores methods such as Integrated Pest Management (IPM), biological control, and the application of environmentally friendly pesticides, evaluating their effectiveness in controlling pest populations and preserving ecological balance. The review highlights the importance of employing diverse strategies tailored to specific agricultural conditions, including crop types, pest species, and local environments. Furthermore, it discusses the economic advantages of sustainable pest management, such as cost reduction, enhanced crop yields, and improved long-term soil health. The paper concludes by stressing the need for ongoing research and innovation to develop pest management systems that are both ecologically responsible and economically practical, ensuring the sustainability of agricultural practices over time.

Keywords: Sustainable pest management, Integrated Pest Management (IPM), Biological control, Eco-friendly pesticides, Agricultural sustainability

SMART TECHNOLOGIES FOR POLLINATOR-SAFE INTEGRATED PEST MANAGEMENT

Berjin Viju M G^{1*}, Muthukumar M² and Melvin Joe M³

¹PG Scholar, Department of Entomology, SRM College of Agricultural Sciences, SRMIST, Baburayanpettai, Chengalpattu District, Tamil Nadu– 603201

²Asst. Professor and Head, Department of Entomology, SRMCAS, SRMIST, Baburayanpettai, Chengalpattu District, Tamil Nadu– 603201

³Associate Professor (Microbiology) Department of Soil Science & Agricultural Chemistry, SRMCAS, SRMIST Baburayanpettai, Chengalpattu District, Tamil Nadu– 603201

*Corresponding author Email: bm5736@srmist.edu.in

ABSTRACT

Honeybees play a critical role in agriculture as pollinators, yet their survival is increasingly threatened by pests, diseases, pesticide exposure, and environmental stress. Advances in digital technologies are transforming Integrated Pest Management (IPM) into a more precise, predictive, and pollinator-friendly approach. Smart hive monitoring systems using IoT sensors enable early detection of colony stress and pest infestations, while remote sensing and GIS tools help identify pest hotspots and pesticide drift risks in cropping systems. Artificial intelligence and machine learning provide powerful tools for pest recognition, disease diagnosis, and predictive modelling to align pest control with pollinator safety. Decision support systems and mobile platforms further support farmers and beekeepers by integrating real-time data into safe and effective pest management practices. Robotics and precision spraying technologies also minimize chemical exposure to foraging bees. Together, these innovations represent a digital frontier in IPM, offering opportunities to protect honeybees, reduce chemical inputs, and enhance agricultural sustainability.

Keywords: Honeybee Conservation, Digital Integrated Pest Management (IPM), IoT Hive Monitoring, AI-based Pest and Disease Detection, Precision Agriculture

SMART IPM: INTEGRATING AI, IOT, AND REMOTE SENSING FOR REAL-TIME PRECISION PEST MONITORING AND DECISION SUPPORT

Santhoshraj N^{1*} and Murugan N²

¹PG Scholar, Department of Entomology, SRM College of Agricultural Sciences, SRM Institute of Science and Technology, Baburayanpettai, Chengalpattu – 603 201

²Asst. Professor (Sr. Grade), Department of Entomology, SRMCAS, SRMIST, Baburayanpettai, Chengalpattu District, Tamil Nadu– 603201

*Corresponding author Email: sn1734@srmist.edu.in

ABSTRACT

Digital innovations are transforming Integrated Pest Management (IPM) by enabling precise, data driven, and environmentally responsible pest control. This study presents a comprehensive smart IPM framework that integrates artificial intelligence (AI), Internet of Things (IoT) devices, and remote sensing technologies to enhance pest monitoring, forecasting, and decision-making in real time. The system combines multispectral satellite imagery with in-field data from smart traps and microclimate sensors to monitor pest populations and crop health indicators. Machine learning algorithms analyze these data streams to identify pest hotspots, model infestation risks, and generate timely, location specific management enabled early detection of pest outbreaks, leading to proactive rather than reactive pest control measures. This not only resulted in better crop protection and yield stability but also promoted sustainable practices aligned with climate-smart agriculture. By integrating cutting-edge technology with ecological understanding, smart IPM empowers farmers, extension agents, and policymakers to make informed, sustainable pest management decisions, and underscores the potential of digital agriculture to scale precision IPM solutions and reduce agriculture's environmental footprint in the face of evolving pest dynamics and climate challenges.

Keywords: Precision agriculture, smart farming, artificial intelligence, remote sensing, pest forecasting, decision support systems

**PHYTO-RHIZOSPHERE INTERCEPTION FOR NEMATODE EXCLUSION
(P.R.I.N.E.) A LIVING BIO-BARRIER FOR ROOT PEST MANAGEMENT**

Bhuvaneshwaran S^{1*} and Murugan N²

¹PG Scholar, Department of Entomology, SRM College of Agricultural Sciences, SRM Institute of Science and Technology, Baburayanpettai, Chengalpattu – 603 201

²Asst. Professor (Sr. Grade), Department of Entomology, SRMCAS, SRMIST, Baburayanpettai, Chengalpattu District, Tamil Nadu– 603201

*Corresponding author Email: bc0074@srmist.edu.in

ABSTRACT

Phyto-Rhizosphere Interception for Nematode Exclusion (P.R.I.N.E.) introduces a transformative approach to sustainable nematode management by deploying a perimeter of genetically selected "interceptor plants" as a living, self-sustaining bio-barrier. This system specifically targets sedentary endoparasitic nematodes e.g., *Meloidogyne* spp. through a "lure and trap" mechanism. Interceptor plants are engineered to exude root-specific biochemical cues that mimic vulnerable hosts, acting as potent chemo-attractants to divert nematodes from primary crops. Upon root penetration, a rapid hypersensitive response is triggered within interceptor tissues, inducing localized cell death that encapsulates and eliminates juvenile nematodes before they establish feeding sites or reproduce. This method offers significant advantages over conventional strategies. It eliminates reliance on synthetic nematicides, reducing chemical runoff, soil toxicity, and non-target harm to beneficial soil organisms. By leveraging ecological deception rather than chemical toxicity, P.R.I.N.E. circumvents pest resistance development. The system operates continuously throughout the growing season, requiring minimal human intervention once established. Interceptor plants also contribute to soil health—post-harvest, their biomass can be tilled into soil as green manure, releasing natural biofumigants that further suppress nematode populations while improving soil organic matter. Compatible with diverse cropping systems, P.R.I.N.E. is scalable for both smallholder and industrial farming. It provides an economically viable, eco-friendly solution that aligns with agroecological principles, enhancing crop resilience and supporting biodiversity. Crops protected by P.R.I.N.E. also offer market advantages as sustainably produced goods, appealing to environmentally conscious consumers. This innovation represents a proactive, systems-based strategy for long-term root pest management.

Keywords: Sustainable agriculture, nematode management, root exudates, bio-barrier, hypersensitive response, agroecology, soil health, green manure, resistance management.

ECOLOGICAL INNOVATIONS FOR SUSTAINABLE PEST MANAGEMENT

Kirubakaran G^{*1} and Nisha R²

¹PG Scholar, Department of Entomology, SRM College of Agricultural Sciences, SRMIST, Baburayanpettai, Chengalpattu – 603 201

²Assistant Professor, Department of Entomology, SRM College of Agricultural Sciences, SRMIST, Baburayanpettai, Chengalpattu – 603 201

*Corresponding author Email: kg3301@srmist.edu.in

ABSTRACT

India's agriculture has distinct challenges due to its extensive crop diversity, the preponderance of small landholdings, and the existence of diverse agroecological zones. Ecological innovation in pest control offers a lot of promise to make food more secure, help rural economies, and protect the environment. The new developments in sustainable pest control in India on rice, cotton, vegetables, and pulses were seen. It also looks at the ecological, economic, and social aspects of these developments. Significant progress includes the increased use of biopesticides and biological agents like *Bacillus thuringiensis*, *Trichoderma spp.*, *Beauveria bassiana*, neem extracts, and specificity-based natural antagonists. These agents lower the need for synthetic pesticides and lessen their effects on non- target organisms. Changes to crop ecosystems, such as intercropping, trap crops, cover crops, hedgerows, and refuges and to keep predator and parasitoid populations alive in cotton and vegetable systems. Precision agriculture tools, such as AI and drone-based early detection systems for diseases and insect outbreaks in crops like cashew, rice, and cotton, are reducing the time it takes to act and enabling the more effective use of control chemicals. Comparative field trials in bio-intensive pest management, such as those conducted on cowpea in the sub-Himalayan region, demonstrate that while traditional chemical-based methods may occasionally achieve superior pest suppression or crop yield, eco-friendly approaches more effectively conserve natural predators and pose diminished risks to human health and the environment. An economic study reveals that ecological systems exhibit better cost-benefit ratios, particularly when considering factors such as health hazards, environmental damage, and waste. Challenges to adoption include the diversity of small farms, insufficient farmer education or extension assistance, regulatory and market access difficulties for biopesticides, and the initial capital required for monitoring and data systems. Ecological innovation has significant potential for sustainable pest control in India; the integration of biological, cultural, and technological approaches may provide resilient, productive, and ecologically sustainable agricultural systems.

Keywords: Ecology, predator, parasitoid, sustainable agriculture

HARNESSING THE BLACK SOLDIER FLY (*Hermetia illucens*) FOR SUSTAINABLE WASTE-TO-WEALTH CONVERSION

Ramazeame L¹, Athira Rajan², Karthik² and Berjin Viju M G²

¹Assistant Professor, Department of Entomology, SRM College of Agricultural Sciences
Baburayanpettai, Chengalpattu – 603 201

²PG Scholar, Department of Entomology, SRM College of Agricultural Sciences
Baburayanpettai, Chengalpattu – 603 201

Corresponding author Email: ramazeal@srmist.edu.in

ABSTRACT

Organic waste, particularly food and fruit/vegetable residues, imposes a significant environmental as well as economic burden in urban and agriculture systems. The larvae of the black soldier fly, *Hermetia illucens*, provide a potentially interesting bioconversion process to convert these organic wastes into two valuable materials (insect biomass as animal feed and residual frass as an organic fertilizer). In this review/study, we consider the efficiency and positive and negative aspects of food wastes - fruit waste converting into feed and manure by means of black soldier fly larvae (BSFL). The latter reduces the input weight of 60–85 %, depending on kind of substrate and larval density as well as process conditions. The larvae produced are high in protein and lipids, which make them effective as a supplement or partial replacement in aquaculture, poultry and livestock feed. In the meantime, frass as a by-product contains plenty of macro/micro-nutrients (NPK etc.) and could be used to enhance soil properties, however management of its stability, moisture content and maturity is necessary. Key challenges are substrate heterogeneity, moisture balance, process scale-up, separation and sanitization costs, and regulatory acceptance. Life cycle assessments indicate that when operating under ideal conditions the CO₂ intensity and resource use associated with bioconversion using BSFL can be lower than disposal into alternative methods of waste treatment. It therefore stands out as a promising technology within circularity of organic waste, connecting waste valorization to sustainable feed and fertilizer production—assuming operational, economic and regulatory barriers can be tackled.

Keywords: Black soldier fly, *Hermetia illucens*, waste conversion, manure

EVALUATION OF BIO-INTENSIVE IPM MODULES FOR MANAGEMENT OF INSECT-PEST–PATHOGEN COMPLEX IN BHENDI ECOSYSTEM

Rex B

Assistant Professor, Department of Plant Pathology, SRM College of Agricultural Sciences, SRMIST, Baburayenpettai, Chengalpattu, Tamil Nadu.

*Corresponding author Email: rexpatho@gmail.com

ABSTRACT

Bhendi (*Abelmoschus esculentus* L. Moench) is a high-value vegetable crop widely cultivated in South India, but its productivity is severely constrained by the simultaneous occurrence of major insect pests such as the shoot and fruit borer (*Earias vittella*), jassids (*Amrasca biguttula biguttula*), and whiteflies (*Bemisia tabaci*), along with diseases including yellow vein mosaic virus (YVMV) and Fusarium wilt. Over-reliance on chemical pesticides has resulted in pest resurgence, resistance, and residue hazards. Hence, the present study was undertaken during 2024–2025 in Chengalpattu District to develop an eco- friendly, sustainable management strategy. A field experiment was conducted in a Randomized Block Design (RBD) with six bio-intensive IPM modules and a chemical control check. The modules integrated the use of neem-based formulations, *Beauveria bassiana* and *Trichogramma chilonis* releases, soil application of *Trichoderma harzianum*, intercropping with marigold, and yellow sticky traps. Observations were recorded on pest and disease incidence, yield parameters, and cost-benefit ratio. Preliminary results indicated that the module combining neem oil (3%), *B. bassiana* sprays, and *T. harzianum* soil treatment significantly reduced pest and disease incidence while improving marketable yield by 28.4% over control. The bio-intensive modules also promoted natural enemy abundance and reduced chemical dependency. This study demonstrates the potential of bio-intensive IPM as a viable alternative to synthetic pesticides, contributing to safer produce, environmental sustainability, and enhanced livelihood security for bhendi growers in Tamil Nadu's irrigated ecosystems.

Keywords: Bhendi, *Trichoderma harzianum*, Fusarium wilt, IPM, *Beauveria bassiana*

ECO-FRIENDLY IPM STRATEGIES FOR MAJOR PESTS OF OKRA

Ravanachandar A^{1*}, Devika A² and Dhilip Chakkaravarthy T²

^{1*} Assistant Professor and Head, Department of Vegetable Science, SRM College of Agricultural Sciences, SRM University, Vendhar Nagar, Baburayanpettai – 603 201,

² PG Scholar, Department of Vegetable Science, SRM College of Agricultural Sciences, SRM University, Vendhar Nagar, Baburayanpettai – 603 201, India

*Corresponding author Email: ravanachandar88@gmail.com

ABSTRACT

One of the most widely grown annual vegetable crops grown from seed in India and other tropical and subtropical regions of the world is okra (*Abelmoschus esculentus*), also known as lady's finger and locally called "Bhendi" or "Dherosh." Numerous insect pests target the crop, but the fruit and shoot borer (*Earias spp.*) is the most damaging, resulting in significant damage and production losses of up to 40–50%. Environmental risks, pest resistance, and the extinction of beneficial creatures have resulted from an over-reliance on chemical pesticides for pest management. As a result, in India's agricultural systems, implementing a sustainable Integrated Pest Management (IPM) approach has grown in significance. IPM provides a comprehensive and environmentally responsible strategy by combining several strategies, including the application of resistant cultivars, seed treatment, and successful cultural practices like intercropping, crop rotation, and field sanitation. The use of botanicals or less-toxic pesticides like neem oil and neem seed kernel extract (NSKE), the installation of pheromone traps for pest monitoring, and biological control with natural enemies are important elements of this strategy. In order to promote sustainable pest management and increase okra productivity, farmers must participate in extensive extension activities, demonstrations, and awareness programs in order to guarantee the widespread adoption of IPM.

Keywords: Fruit and shoot borer, okra, Integrated Pest Management (IPM), resistant cultivars, neem oil, NSKE, biological management, fruit and shoot borer, and environmentally friendly pest control

**STUDY ON PHYSICO-BIOCHEMICAL ATTRIBUTES OF BRINJAL
INFLUENCING RESISTANCE TO SHOOT AND FRUIT BORER *Leucinodes
orbonalis* GUENEE**

Rameshkumar D^{1*}, Naveena E², Swarna Priya R³ and Savitha B K³

¹SRM College of Agricultural Sciences, SRM Institute of Science and Technology, Faculty of Agricultural Sciences, Baburayanpettai, Chengalpattu, 603 201, Tamil Nadu,

²St. Joseph University, School of Agricultural Sciences, Tindivanam, Vellore,
Tamil Nadu, India.

³Department of Vegetable Crops, Horticultural College & Research Institute, TNAU,
Coimbatore

*Corresponding author Email: rameshkd@srmist.edu.in

ABSTRACT

An experiment was carried out with six parents and thirty hybrids of brinjal at Tamil Nadu Agricultural University, Coimbatore to find the influence of biophysical and biochemical characters of brinjal on the infestation to shoot and fruit borer. Among the hybrids Sevathampatti local x Spiny local and Sevathampatti local x Seetipulam local was found least attacked by the borers recording minimum percentage of fruit infestation (14.68 percent) with maximum yield per plant of 3.27 kg per plant. Fruit infestation was positively but not significantly correlated with length of calyx ($r = 0.314$) whereas, yield per plant ($r = -0.297$), solasodine ($r = -0.170$) and phenol ($r = -0.572$) showed significantly negative correlation with fruit infestation and positive correlation with shoot infestation ($r = 0.041$), polyphenol oxidase ($r = 0.011$) and total sugars ($r = 0.072$). The cross Manaparai local x Spiny local, Sevathampatti local x Spiny local and Sevathampatti local x Seetipulam local recording maximum yield, less shoot and fruit infestation, lowest calyx length, low sugar, high polyphenol oxidase and high phenol could be used as resistant cultivar for further shoot and fruit borer resistance breeding programme.

Key words: Biophysical, biochemical, *Leucinodes orbonalis*, resistance

**BIOEFFICACY OF NOVEL INSECTICIDES AGAINST BANANA SCARRING
BEETLE *Basilepta subcostata* JACOBY THROUGH LABORATORY BIOASSAY**

Kancharla Ratna Jyothi*, Marri Keerthana and Sai Reddy M S

Department of Entomology, Dr. Rajendra Prasad Central Agricultural
University, Pusa, Samastipur-848125, Bihar, India

*Corresponding author Email: ratnajyothikancharla2001@gmail.com

ABSTRACT

The banana scarring beetle, *Basilepta subcostata* (Jacoby), is a significant pest distributed across major banana-growing regions of India, with historical reports identifying it as a major pest in multiple states including Bihar, Assam, Uttar Pradesh, and West Bengal. It causes substantial economic losses through foliar and fruit damage. To identify effective control strategies, laboratory bioassays were conducted to evaluate the bioefficacy of thirteen insecticidal formulations against adult beetles. Adult mortality and foliar damage were assessed at 4, 6, and 12 hours after treatment (HAT) using a leaf-dip method. Results indicated that Thiamethoxam 25% WG and Fipronil 5% SC were the most effective treatments, achieving 100% mortality across all time intervals and exhibiting the lowest cumulative leaf damage (1.67% and 2.23%, respectively). Imidacloprid 17.80% SL, Chlorpyrifos 20% EC, and Fipronil 0.3% G also demonstrated high efficacy, with mean mortality rates exceeding 95%. In contrast, Azadirachtin 0.03% and Chlorantraniliprole 0.4% G were the least effective, showing the lowest mortality and highest leaf damage among all treatments. Probit analysis of the top-performing insecticides revealed that Thiamethoxam 25% WG had the lowest LC₅₀ value (0.103 ppm) and the steepest dose-response slope, confirming its superior toxicity. The study identifies highly effective chemical and botanical insecticides for the sustainable management of *B. subcostata* in banana cultivation systems.

Keywords: *Basilepta subcostata*, Banana scarring beetle, Insecticide efficacy, Probit analysis, Pest management.

LESS EXPENSIVE, DIY METHOD IN REFLECTED LIGHT MICROSCOPY TO DOCUMENT INSECTS: CONCEPTS AND PROCEDURES

Vinod Kumar S

Department of Mycology and Plant Pathology, Institute of Agricultural Sciences, Banaras
Hindu University, Varanasi - 221005, Uttar Pradesh, India.
Corresponding author Email: vinodfytopathologos@gmail.com

ABSTRACT

The practice of microscopy has been around since the sixteenth century. The field of microscopy has advanced significantly in recent years. Starting with a basic microscope, there are now many different methods, including as electron microscopy, fluorescence microscopy, reflected light microscopy, phase contrast, differential interference contrast, stereo microscopy, and epi lighting. Despite the extreme magnification and high resolution images that electron microscopy provide, we were unable to see the specimen's actual colors. Every type of microscopy requires the sample to be fixed. Delicate structures become crushed, fractured, or deformed as a result. A third eye for seeing the precise morphology of the structures is provided by reflected light microscopy with focus stacking and stitching. More significantly, in contrast to electron microscopy, the sample's original color pattern may be recorded. There are several well-known brands of metallurgical and reflected light microscopes. I'll be talking about the principles and methods of reflected light microscopy. However, not many researchers can afford them due to their high cost. However, they are very expensive and not possible to afford by all the researchers. The DIY technique discussed in my presentation may be useful to amateur photographer, insect enthusiasts and entomologists to photo document insects with less expensive equipment and resulting in a comparably high quality images as that of a standard stereo microscope. The work procedure will be explained with video.

Keywords: Reflected light microscopy, insects, DIY, Concepts and procedures.

OIL BASED FORMULATION OF *Metarhizium anisopliae* (METCHNIKOFF) SOROKIN AGAINST *Helicoverpa armigera* (LEPIDOPTERA: NOCTUIDAE), *Plutella xylostella* (LEPIDOPTERA: PLUTELLIDAE) AND *Spodoptera frugiperda* (LEPIDOPTERA: NOCTUIDAE)

Nandha S ^{*1} and Jeyarani S ²

¹Doctoral student, Tamil Nadu Agricultural University, Coimbatore- 641 003

²Professor, Department of Agricultural Entomology, Tamil Nadu Agricultural University, Coimbatore- 641 003

*Corresponding author Email: nandhasaminathan0808@gmail.com

ABSTRACT

Effective biopesticide formulations are crucial for field-level pest management, ensuring enhanced stability, targeted delivery, and prolonged efficacy under diverse environmental conditions. This study developed and evaluated oil-based formulations of *Metarhizium anisopliae* (TNAU-MA- GDU) for their efficacy against key pests, *Helicoverpa armigera*, *Plutella xylostella*, and *Spodoptera frugiperda*, under laboratory conditions. It also optimized culture media for large-scale conidial production and assessed the physico-chemical properties and safety of the formulations. Potato dextrose agar (PDA) was the most effective medium, achieving a radial growth of 81.30 mm and a conidial yield of 2.16×10^8 spores mL⁻¹. Sabouraud maltose agar with yeast (SMA+Y) and Sabouraud dextrose agar with yeast (SDA+Y) also supported significant growth (65.60 mm and 60.60 mm) and yields of 1.16×10^8 and 1.02×10^8 spores mL⁻¹, respectively, suitable for industrial production. Compatibility tests identified sunflower, soybean, mineral, corn, castor, and palm oils as viable carriers at 2–10% concentrations, with palm oil excelling at 10%, yielding a colony diameter of 81.00 mm and 1.61×10^8 spores mL⁻¹. Skimmed milk and ascorbic acid were optimal UV protectants, preserving 65.33% and 64.67% spore germination after 90 minutes of UV exposure at 5% concentration. Emulsifiable concentrate (EC) and oil dispersion (OD) formulations were developed, with OD showing superior spore viability (86.33% and 85.33% germination on sterile and media-coated slides). Both formulations maintained stability across real-time, refrigerated, and accelerated storage, with OD retaining 168.67×10^8 CFU mL⁻¹ after 28 days under ambient conditions. At 119 days, OD exhibited the highest viability (83.66×10^8 CFU mL⁻¹) compared to EC, wettable powder (WP), and crude suspensions. Pathogenicity assays confirmed OD's superior efficacy, with the lowest LC₅₀ (1.56×10^3 to 3.16×10^5 spores mL⁻¹) and LT₅₀ (94.53–147.33 hours) across pest instars. Safety assessments showed minimal impact on parasitoids *Telenomus remus* and *Trichogramma pretiosum* (>83% emergence) and moderate effects on predator *Chrysoperla zastrowi sillemi* (survival, pupation, and emergence).

59%, 81%, and 81% respectively). The OD formulation's exceptional stability, efficacy, and safety highlight its potential as a robust biopesticide for sustainable field-level pest management.

Keywords : *Metarhizium anisopliae*, UV protectants, oil dispersion

**ENTOMOPATHOGENIC POTENTIAL OF SOIL-ISOLATED
ENTOMOPATHOGENIC FUNGI *Metarhizium robertsii* STRAIN TNAU ENTMR GYU
1 AGAINST ASH WEEVIL (*Myllocerus subfasciatus*)**

Monisha U¹ and Shanmugam P S^{2*}

¹Doctoral student, Department of Agricultural Entomology, AC&RI,
TNAU, Coimbatore -641003, India.

²Associate Professor, Department of Agricultural Entomology, AC&RI, TNAU, Coimbatore -
641003, India.

*Corresponding author Email: Shanmugam.ps@tnau.ac.in

ABSTRACT

The ash weevil *Myllocerus subfasciatus* Guérin-Ménéville (Curculionidae; Coleoptera) poses a significant threat to various crops, particularly in tropical and subtropical regions. The adults are foliage feeders and grub feeds on the roots making them difficult to manage. In search of sustainable and eco- friendly management options, the present study focused on the isolation and characterization of the entomopathogenic fungus *Metarhizium robertsii* (J.F. Bisch., Rehner & Humber) from rhizospheric soil and evaluated its pathogenic potential against *M. subfasciatus*. Soil samples were collected from different districts of Tamil Nadu, Kerala and Karnataka. The fungal isolates were obtained using the insect bait method with *Galleria mellonella* (Linnaeus) (Pyrilidae; Lepidoptera) larvae. The results of isolates through morphological identification confirmed *Metarhizium* spp. The colony was white in colour on vegetative phase and dark green on reproductive phase. Molecular identification was performed using ITS-rDNA sequencing, and BLAST analysis confirmed the identity as *M. anisopliae*. The isolates were submitted in NCBI database and got the accession numbers. Bioassays conducted under laboratory conditions demonstrated significant pathogenicity of the isolate against second instar ash weevil grubs, with mortality rates exceeding 50% within 7-9 days post-inoculation. The sporulation and virulence were studied through SEM and ultramicroscopic studies. TNAU ENTMR GYU 1 demonstrated superior efficacy with lowest LC₅₀ of 2.13x10⁶ conidia/ml respectively in the second instar grub of *M. subfasciatus*. The lethal time (LT₅₀) values were 7.04 days calculated to assess virulence. The findings suggest that among the isolates the *M. anisopliae* TNAU ENTMR GYU 1 possesses promising biocontrol potential and could serve as an effective component in integrated pest management (IPM) strategies for ash weevil control. Further field validation and formulation development are recommended to facilitate its practical application.

Keywords: Ash weevil, TNAU ENTMR GYU 1, *M. robertsii*, histopathology, SEM

EVALUATING APHID RESISTANCE IN BREAD WHEAT *Triticum aestivum* L. GENOTYPES UNDER FIELD AND GROWTH CHAMBER CONDITIONS

Marri Keerthana*¹, Priya ², Kancharla Ratna Jyothi ³ and Sai Reddy M S⁴

Department of Entomology, Dr. Rajendra Prasad Central Agricultural University, Pusa,
Samastipur-848125, Bihar, India

*Corresponding author Email: keerthanamarri1004@gmail.com

ABSTRACT

Aphids (*Sitobion avenae*, *Rhopalosiphum padi*, and *Schizaphis graminum*) are emerging as a serious threat to wheat production in India, with the potential to significantly reduce future crop productivity. To address this challenge, 400 bread wheat genotypes were evaluated for aphid resistance under natural field conditions during the Rabi seasons of 2023-24 and 2024-25, as well as under growth chamber conditions during Rabi season of 2024-25 at Dr. Rajendra Prasad Central Agricultural University, Pusa. The study employed the Aphid Resistance Index (ARI) scored on a 5-point scale. The evaluation revealed substantial variation in resistance levels among genotypes. Based on ARI scores, 82 genotypes were categorized as highly susceptible (21 or more aphids/plant), 283 genotypes as susceptible (11 to 20 aphids/plant) and 55 genotypes as moderately resistant (6 to 10 aphids/plant). No genotypes demonstrated immunity (0 aphids/plant) or resistance (1 to 5 aphids/plant). These findings highlight promising wheat genotypes with partial resistance to aphids, which can serve as valuable genetic resources for breeding programs aimed at enhancing aphid resistance and sustaining wheat productivity under field conditions.

Keywords: Wheat, *Sitobion avenae*, *Rhopalosiphum padi*, *Schizaphis graminum*, Aphid Resistance

GENOMIC INSIGHTS INTO THE ANTIFUNGAL AND PLANT GROWTH PROMOTING TRAITS OF *Pseudomonas plecoglossicida* NAN2 ISOLATED FROM THE RICE RHIZOSPHERE

Nandhini Kuberan and Iyappan Sellamuthu*

Department of Genetic Engineering, School of Bioengineering, Faculty of Engineering and Technology, SRM Institute of Science and Technology, Kattankulathur, Chengalpattu 603 203, Tamil Nadu, India.

*Corresponding author Email: nandhinin049@gmail.com

ABSTRACT

Plant growth promoting rhizobacteria (PGPRs) are beneficial microorganisms that inhabit the rhizosphere and enhance plant growth through various mechanisms. In this study, a PGPR strain designated NAN2 was isolated from the rice rhizosphere and demonstrated multiple plant growth promoting traits, including the production of hydrogen cyanide (HCN), ammonia, indole-3-acetic acid (IAA), phosphate solubilization and antifungal activity against *Magnaporthe oryzae*. Complete genome sequencing and annotation of strain NAN2 revealed a genome size of 5356785 basepairs (bp) with a GC content of 62 %, comprising 227 contigs, 4807 coding sequences (CDSs) and a total of 4960 genes. Notably, the genome contains a nonribosomal peptide synthetase (NRPS) gene cluster associated with the biosynthesis of rhizomides (A, B and C). These results suggest that NAN2 has strong potential as an environmentally resilient biocontrol agent that can protect plants from invasive diseases. To our knowledge, this is the first genomic analysis of *Pseudomonas plecoglossicida* NAN2 isolated from rice fields, providing valuable insights into its biocontrol capabilities and plant growth promoting (PGP) properties.

Keywords: Antifungal, Plant Growth Promoting, Traits *Pseudomonas Plecoglossicida*, Nan2, Rice, Rhizosphere

FRUIT FLY DIVERSITY AND RESISTANCE SCREENING IN MANGO AND CUSTARD APPLE CULTIVARS OF RAYALASEEMA FOR SUSTAINABLE PEST MANAGEMENT

Tirumala Geethika G*, Sarada G, Ramaiah M, Nischala A and Rajeswari G
Department of Entomology Dr. YSR Horticultural University, Andhra Pradesh 516105
*Corresponding author Email: tirumalageethika@gmail.com

ABSTRACT

The study to document fruit fly species diversity in different fruit crops and their sources of resistance were carried out at College of Horticulture, Anantharajupeta during the years 2023-2024. Fruit fly infested fruit samples of mango and custard apple collected through survey from farmer fields of Ananthapuramu, Chittoor, Nandyal and Annamayya districts of the Rayalaseema region and reared in the laboratory. A total of 15 mango varieties were evaluated under field conditions to assess their susceptibility to fruit fly infestation. Among these, four varieties Jahangir×Khader, Neelum, Prabhashankar and Suvarnarekha were classified as moderately resistant, exhibiting infestation levels ranging from 21% to 50%. Ten varieties Bangalora, Neelum, Baneshan, Yeramanda, Ali Pasand, Neeleshan, Royal Special, Pulihora, Proddutur Avakay and Peether were categorized as susceptible, with infestation levels between 51% and 75%. Notably, Dashaheri recorded the highest infestation rate at 75.21%, indicating it as the most preferred variety by fruit flies. Similarly, in custard apple, three varieties Atemoya × Washington, Yengalampalli No.13 and Rayadurg were found to be resistant, with infestation levels between 11% and 20%. Nine cultivars Pythota-5, Arka Sahan, APK-1, Pythota-1, CRIDA, Pythota-3, Jamdugumpala No.7, Sinhan Local and Molkalnur-8 were moderately resistant (21%–50%), while three varieties Balanagar, NMk-Gold and Pythota-2 were classified as susceptible (51%–75%). Importantly, none of the tested mango or custard apple varieties exhibited complete resistance to fruit fly infestation. These findings reflect the natural host preference of fruit flies observed under field conditions.

Keywords: Fruit fly, Mango, Custard apple, Screening, Varieties.

ADVANCEMENTS IN ECOLOGICAL ENGINEERING FOR INSECT PEST CONTROL

Sharulatha S^{1*} and Muthukumar M²

¹PG Scholar, Department of Entomology, SRM College of Agricultural Sciences, SRM Institute of Science and Technology, Baburayanpettai, Chengalpattu – 603201

²Asst. Professor and Head, Department of Entomology, SRMCAS, SRMIST, Baburayanpettai, Chengalpattu District, Tamil Nadu– 603201

*Corresponding author Email: ss6710@srmist.edu.in

ABSTRACT

In agriculture, sustainable pest management is essential because of rising chemical resistance, harm to the environment, and worries about human health. Using ecological innovations that use biodiversity, natural processes, and new technology may help control pests while causing the least amount of harm. Some of the strategies include Integrated Pest Management (IPM), biological management using beneficial insects, entomopathogens, and habitat alteration methods including companion planting, trap cropping, and polyculture. The Sterile Insect Technique (SIT), biofumigation, biosolarization, and smart monitoring utilising IoT sensors or computer vision are all examples of advanced techniques that have even more potential. Case studies—from smallholder farms in Africa using push-pull systems and biopesticides to rice farmers in South-east Asia using adaptive networks— show how important it is to adapt to the local environment, get stakeholders involved, and provide socio-economic incentives. Ecological pest control protects ecosystem services, soil health, and biodiversity. It also helps the economy by lowering input costs and increasing yields over time. The conclusions stress the need for policies that work together, research that involves everyone, and investments that make farming techniques fit with how nature works for long-term sustainability.

Keywords: Sustainable pest management; ecological innovations; Integrated Pest Management; biological control; habitat manipulation; precision monitoring; ecosystem services; yield stability; environmental sustainability

**ECOLOGICAL INNOVATIONS IN PEST MANAGEMENT: ENTOMOLOGICAL
PATHWAYS TO SUSTAINABLE AGRICULTURE**

Karthik S^{1*} and Ramazeame L²

¹PG Scholar, Department of Entomology, SRM College of Agricultural Sciences, SRM
Institute of Science and Technology, Baburayanpettai, Chengalpattu – 603201

²Asst. Professor, Department of Entomology, SRM College of Agricultural Sciences, SRM
Institute of Science and Technology, Baburayanpettai, Chengalpattu – 603201

Corresponding author Email: karthiksanthosh7744@gmail.com

ABSTRACT

Agricultural sustainability requires pest management approaches that harmonize productivity with ecological balance. Harnessing nature through entomological innovations offers eco-friendly alternatives to chemical-intensive practices. Beneficial insects provide vital services as pollinators, predators, parasitoids, decomposers, and bioindicators, while strategies such as Integrated Pest Management (IPM), conservation biocontrol, pheromone-based monitoring, and biopesticides strengthen natural regulation of pests. Recent advances in genetic tools, insect farming, remote sensing, and artificial intelligence further expand opportunities for precision and resilience in pest control. Although barriers such as limited awareness, high initial costs, and climate change impacts remain, ecological pest management presents a transformative pathway to safeguard food security, enhance biodiversity, and promote sustainable farming systems.

Keywords: Sustainable agriculture, Integrated pest management, Artificial intelligence.

**ECOLOGICAL PEST MANAGEMENT: INSECTS AS CATALYSTS FOR
SUSTAINABLE AGRICULTURE**

Athira Rajan^{1*} and Ramazeame L²

¹PG Scholar, Department of Entomology, SRM College of Agricultural Sciences, SRM
Institute of Science and Technology, Baburayanpettai, Chengalpattu – 603201

²Asst. Professor, Department of Entomology, SRM College of Agricultural Sciences, SRM
Institute of Science and Technology, Baburayanpettai, Chengalpattu – 603201

Corresponding author Email: ar1036@srmist.edu.in

ABSTRACT

Sustainable agriculture increasingly depends on ecological innovations that harness the natural roles of insects to reduce reliance on synthetic pesticides and safeguard ecosystem health. Insects serve as pollinators, biological control agents, decomposers, bioindicators, and even alternative protein sources, making them indispensable to resilient farming systems. This paper highlights nature-based pest management strategies such as Integrated Pest Management (IPM), conservation of natural enemies, habitat manipulation, and the use of pheromone traps and biopesticides. Emerging tools—including genetic approaches, sterile insect techniques, and AI-based monitoring—further enhance the ecological regulation of pests while minimizing environmental harm. Despite challenges such as knowledge gaps, economic constraints, and climate-induced shifts in pest dynamics, integrating entomological innovations into agricultural practices provides a pathway toward balanced productivity, biodiversity conservation, and long-term food security. By harnessing nature's own mechanisms, ecological pest management fosters a sustainable future for agriculture that aligns productivity with environmental stewardship.

Keywords: Sustainable agriculture, Integrated pest management, AI based technology.

ECOLOGICAL INNOVATIONS FOR SUSTAINABLE PEST MANAGEMENT IN BHENDI

Arun Prasad V^{1*} and Nisha R²

¹PG Scholar, Department of Entomology, SRM College of Agricultural Sciences, SRMIST, Baburayanpettai, Chengalpattu - 603201

² Assistant Professor, Department of Entomology, SRM College of Agricultural Sciences, SRMIST, Baburayanpettai, Chengalpattu - 603201

*Corresponding author Email: av4779@srmist.edu.in

ABSTRACT

Bhendi (*Abelmoschus esculentus*) is severely impacted by pests like shoot & fruit borer (*Earias spp.*), whitefly, leafhoppers, aphids and mites, which lower both yield and fruit quality. Reliance on chemical insecticides has led to resistance, environmental harm, residue issues and loss of beneficial organisms. Recent ecological innovations offer sustainable alternatives: Integrated Pest Management (IPM) approaches tailored for bhendi combine regular monitoring, threshold-based interventions, and use of biocontrol agents (parasitoids, predators) to reduce dependence on chemicals. Botanicals and biopesticides (e.g. *Bacillus thuringiensis*, *azadirachtin*, neem products, Spinosad) have shown strong reductions in both shoot and fruit infestation, sometimes exceeding 50–80% lower pest load, with improved economic returns. Cultural practices such as intercropping with non-host or repellent plants, use of trap/border crops, removal and destruction of infested shoots/fruits, conservation or augmentation of natural enemies, and varietal selection for partial resistance also contribute significantly to pest suppression. Some IPM modules have demonstrated not only lower pest incidence but also higher yield and benefit-cost ratios compared to conventional pesticide-heavy regimes. Challenges remain: availability of resistant varieties suited to local conditions, consistency and efficacy of biocontrol under field conditions, farmer awareness, labor and cost for monitoring and implementing IPM components, and supportive policy/regulatory frameworks. Nonetheless, the confluence of cultural, biological, and technological innovations provides a pathway for sustainable pest management in bhendi ensuring stable, high yields, reduced environmental footprint, safer produce, and improved livelihood resilience for farmers.

Keywords: Bhendi, Biocontrol, IPM, Sustainable agriculture, Natural Enemies, Pest outbreak.

**ECO-FRIENDLY AND CLIMATE-SMART STRATEGIES FOR MANAGING
COFFEE BERRY BORER IN ANDHRA PRADESH**

**Urati Mahesh*, Venkateswara Reddy M, Chaitanya K, Rajini D, Harika M and
Lavanya A V N**

Department of Horticulture, College of Agriculture, Rajendranagar, PJTAU, Telangana

*Corresponding author Email: u.maheshreddy98@gmail.com

ABSTRACT

The coffee berry borer (*Hypothenemus hampei*), a globally recognized pest of coffee, has recently emerged in Andhra Pradesh, with infestations first reported in the Alluri Sitarama Raju district's Araku Valley and already spreading from 80 to 140 acres within weeks. This development poses a critical threat to more than 2.5 lakh acres of coffee plantations and the livelihoods of 2.45 lakh tribal farmers in the region. Given the pest's cryptic lifecycle inside coffee berries, conventional chemical control measures remain largely ineffective, making eco-friendly strategies the cornerstone of sustainable management. Ecological innovations such as shade tree regulation, berry sanitation, maintenance of on-farm biodiversity and the use of biological control agents like *Beauveria bassiana* can provide effective, environmentally safe suppression. Coupled with climate-smart practices that integrate weather-based forewarning models, soil health management and adaptive cultural operations, these approaches enhance resilience under changing climatic conditions. Moreover, incorporating digital tools for pest surveillance and farmer advisories can complement ecological interventions, enabling timely and precise responses. Addressing such emerging pest challenges requires fostering multidisciplinary collaboration among researchers, extension agents, farmers and policymakers to co-develop integrated solutions that are both socially acceptable and ecologically sustainable. The case of the coffee berry borer in Andhra Pradesh underscores the urgent need to harness nature-based innovations, supported by digital and collaborative frameworks, to secure crop protection and safeguard the future of India's coffee sector.

Keywords: Coffee berry borer, Sustainable management, Climate resilience, Biocontrol, Collaboration.

INTEGRATED ECOLOGICAL AND DIGITAL APPROACHES FOR MANAGING TOMATO PINWORM IN TELANGANA

Urati Mahesh*, Chaitanya K, Lavanya A V N, Venkateswara Reddy M, Harika M and Rajini D

Department of Horticulture, College of Agriculture, Rajendranagar, PJTAU,
Telangana

*Corresponding author Email: u.maheshreddy98@gmail.com

ABSTRACT

The tomato pinworm (*Keiferia lycopersicella*), a highly invasive pest of tomato, has rapidly emerged as a significant challenge in India, with severe outbreaks reported in Telangana's key tomato-growing districts such as Khammam and Adilabad. Yield losses ranging from 20–30% in open fields to over 70% in polyhouse cultivation have been recorded, threatening farmer incomes and regional vegetable supply chains. Conventional chemical control measures are limited due to the pest's cryptic leaf-mining and fruit-boring behaviour and the risk of resistance development. In this context, eco-friendly strategies form the backbone of sustainable pest suppression. Integrated ecological approaches such as the use of pheromone traps for mass trapping and monitoring, regular destruction of infested plant parts, neem-based botanicals (*Azadirachtin*) and biological control agents like *Trichogramma pretiosum* and *Bacillus thuringiensis* have shown significant promise. Climate-smart practices, including crop rotation, resistant varieties, and optimized irrigation regimes, enhance resilience, while incorporating weather-based forewarning models can help predict pest outbreaks. Further, digital tools such as mobile-based advisory systems and sensor-assisted pest surveillance can complement ecological interventions, ensuring timely and precise farmer action. The Telangana experience underscores the urgency of harmonizing ecological innovations, digital technologies and climate-smart practices to safeguard tomato production and strengthen sustainable crop protection frameworks across India.

Keywords: Tomato pinworm, Sustainable management, Climate resilience, Biocontrol, Eco-friendly practices

BIOEFFICACY OF NEW SPIROTETRAMAT 150 OD AGAINST APHID *Aphis gossypii* IN BRINJAL

Gajalakshmi M* and Gunasekaran K

Tamil Nadu Agricultural University, Coimbatore

*Corresponding author Email: gajalakshmiagri@gmail.com

ABSTRACT

An experiment was conducted to assess the efficacy of spirotetramat 150OD against aphids in brinjal ecosystem. Three doses of spirotetramat 150 OD, viz at 60, 75, 90 g a.i. ha⁻¹ along with two standard checks, Thiamethoxam 25WG at 25 g a.i. ha⁻¹ and Dicofol 18.5 EC at 250 g a.i. ha⁻¹ were appraise for their relative efficacy to aphids. After two rounds of application the highest dose of spirotetramat 150 OD at 90 g a.i.ha⁻¹ and 75 g a.i. ha⁻¹ had registered mean reduction of 88.11; 86.72 and 86.28; 85.26 per cent over untreated check in the first and second experiment, respectively on *A. gossypii*. Thiamethoxam 25 WG at 25 g a.i ha⁻¹ registered mean reduction of 65.64 and 75.10 per cent in aphid's population over untreated check. Based on the results acquired from field studies spirotetramat was found to be effective against *A.gossypii*.

Key words: Spirotetramat, brinjal, aphids, efficacy and brinjal

ADVANCES IN INTEGRATED PEST MANAGEMENT OF SAN JOSE SCALE IN MULBERRY: SYNERGISTIC ROLE OF BIOPESTICIDES, PREDATORS, AND CULTURAL PRACTICES

Murugan Nagarajan^{*1} and Vairam Namachivayam²

¹Assistant Professor-Senior Grade, Department of Entomology, SRM College of Agricultural Sciences, Baburayenpettai, Chengalpattu.

²Assistant Professor, Department of Agricultural Engineering, SRM Valliammai Engineering College, Kattankulathur, Chengalpattu.

*Corresponding author Email: murugan.agri@gmail.com

ABSTRACT

San Jose scale (*Quadraspidiotus perniciosus* Comstock) is an economically important pest of mulberry, reducing leaf yield and quality essential for silkworm rearing. To overcome limitations of chemical control, the present research at SRM College of Agricultural Sciences, Baburayenpettai, was conducted on mulberry variety Victory1 and Genotype 4. The study aimed to evaluate advances in integrated pest management (IPM) with a focus on the synergistic role of biopesticides, predators, and cultural practices. Field trials compared the response of Victory1 and Genotype 4 under Integrated Pest Management modules involving neem oil, pungam oil, and fish oil rosin soap sprays, integrated with natural predators (*Chilocorus nigritus* (Fabricius), *Cryptolaemus montrouzieri* (Mulsant)) and parasitoids (*Aphytis melinus*), along with pruning and cultural practices. Results demonstrated that both Victory1 and Genotype 4 recorded significant reduction in pest incidence (65–82%) under Integrated Pest Management modules, though Genotype 4 exhibited comparatively higher tolerance to infestation. Biopesticide–predator combinations produced a synergistic effect, while cultural practices minimized reinfestation. Leaf yield and quality improved notably in Victory1 under IPM, ensuring better feed for silkworm rearing. Economic analysis revealed higher benefit–cost ratios in both varieties under IPM compared to conventional management. The findings highlight that integrating eco-friendly IPM strategies tailored for mulberry varieties like Victory1 and Genotype 4 can sustainably suppress San Jose scale, support mulberry leaf productivity and align with climate-resilient practices.

Keywords: Mulberry, San Jose scale, Biopesticides, Predators, Cultural practices

SMART HORTICULTURAL PRODUCTION – A NOVEL APPROACH

Dhivya M*, Usha G and Muruganantham M

Assistant Professor, Department of Horticulture PGP College of Agricultural Sciences,
Namakkal -637405, Tamil Nadu

*Corresponding author Email: drdhivya26@gmail.com

ABSTRACT

In the realm of intelligent agriculture, the concept of smart horticulture production has been gaining traction. It is revolutionizing the agriculture industry by incorporating advanced technologies to increase efficiency and productivity. From automated irrigation systems to drones for monitoring crops, the possibilities are endless. However, the current horticulture production and quality levels are not sufficient to meet the demands of a growing population, climate change, a decrease in the labour force, and a lack of automation. To fully embrace smart horticulture production, need to invest in cutting-edge technologies such as sensors, drones, and automated machinery. Artificial intelligence, big data, the Internet of Things, and cloud computing can provide valuable data and insights to optimize crop production for sustainable growth. With the integration of smart technologies, farmers can remotely monitor soil moisture levels, detect pest infestations early, and adjust irrigation schedules accordingly. This level of precision allows for more sustainable and efficient farming practices. Over the next 30 years, it is anticipated that the industry will evolve towards a smart industry, utilizing mechanical, automated, and information-based processes for production. Smart horticulture production is transforming the way we grow crops. By harnessing the power of technology, farmers can improve yields, conserve resources, and ultimately, feed a growing population. The future of agriculture is smart, and the possibilities are endless.

Keywords: Sustainable pest management, Smart agriculture, Artificial intelligence.

ROLE OF CROP DIVERSIFICATION IN REDUCING PEST OUTBREAKS IN CEREAL-BASED CROPPING SYSTEMS

Jeyajothi R

Assistant Professor, Department of Agronomy, SRM College of Agricultural Sciences, SRM Institute of Science and Technology, Baburayanpettai, Chengalpattu District - 603 201, Tamil Nadu, India.

Corresponding author Email: jeyajotr@srmist.edu.in

ABSTRACT

Crop diversification is an ecologically sustainable approach to pest management that enhances the stability and resilience of cereal-based agroecosystems. Monocropping systems, which dominate modern agriculture, often lead to pest outbreaks due to the continuous availability of host plants and the disruption of natural enemy habitats. In contrast, diversified cropping systems through intercropping, crop rotation, relay cropping, and the inclusion of legumes or cover crops promote ecological balance and suppress pest populations by disrupting pest life cycles and enhancing biological control. The increased plant diversity modifies microclimatic conditions, alters pest colonization behavior, and supports a wider range of predators and parasitoids. Furthermore, diversified systems improve soil health and microbial activity, which indirectly contributes to pest resistance and crop vigor. Empirical studies in rice–wheat and maize sorghum systems have shown that crop diversification can significantly reduce the incidence of major insect pests such as stem borers, aphids, and leafhoppers, while improving yield stability and ecosystem services. Therefore, integrating crop diversification within cereal-based farming not only reduces the dependency on chemical pesticides but also strengthens the ecological integrity of agricultural landscapes. The present review highlights the mechanisms, benefits, and management strategies of crop diversification as a cornerstone of ecological pest management in sustainable cereal production systems.

Key words: Cropping system, relay cropping, cover crops, diversified cropping system

IPM SIGNIFICANCE IN SAFEGUARDING FRESH WATER RESOURCES

Beslin Joshi J

Scientist, Centre for Water Resources Development and Management,

Kozhikode – 673 571

Corresponding author Email: beslinjoshi@cwrwm.org

ABSTRACT

Pesticides contaminate water bodies by running off agricultural land or urban areas and entering the surface water and groundwater. These toxic chemicals can harm aquatic ecosystems, causing oxidative stress in organisms and leading to bioaccumulation through the food chain. This contamination threatens biodiversity and poses risks to human health through the consumption of contaminated fish and water. Integrated Pest Management (IPM) safeguards freshwater resources by significantly reducing the overuse of chemical pesticides that contaminate groundwater. IPM emphasizes preventing pest problems through better crop management practices, such as selecting resilient crop varieties, proper soil health management, and good field hygiene, which lessens the need for chemical intervention. IPM employs a combination of cultural, biological, and physical methods to control pests, and only resorts to targeted chemical use as a last resort. This multi-faceted approach ensures that only necessary, low-risk chemical applications are used, and only when justified by economic thresholds, further limiting water pollution. With advances in digital technology, farmers today can remotely monitor environmental conditions (e.g., temperature, humidity, soil moisture) and detect pests or disease outbreaks at their earliest onset. Drones for crop monitoring and aerial surveillance are transforming the fight against pests across large fields or in inaccessible areas. Water management can be linked to pest incidence. For instance, practices like drip irrigation or planting on raised beds can control weeds and use water more efficiently, thereby reducing the amount of water needed for irrigation and preventing potential pest issues that could lead to more pesticide use. This reduces pesticide runoff and leaching, thereby protecting water quality and promoting healthier ecosystems.

Keywords: Integrated Pest Management, Water Resources, Reduce Pollution, Sustainability

**BIO-DIGITAL PEST MANAGEMENT: INTEGRATING LIVING SENSORS
(INSECTS, MICROBES) WITH AI NETWORKS**

Anitha D ^{*1} and Baradhan G ²

¹ Research Scholar, Faculty of Agriculture, Annamalai University

² Associate Professor, Agricultural College and Research Institute, Vazhavachanur

*Corresponding author Email: durairajanjanitha584@gmail.com

ABSTRACT

The rapid escalation of global pest challenges in agriculture has underscored the need for innovative, sustainable and precise management strategies. Even if conventional chemical-based control techniques are short-term successful, they endanger ecosystems, biodiversity and human health in the long run. Bio-Digital Pest Management is a paradigm shift that combines sophisticated artificial intelligence (AI) networks with biological living sensors, including insects, microbial ecosystems and bioengineered creatures to overcome these constraints. Real-time AI- driven analytics can digitize and understand their bio-signals. Predictive pest surveillance, early outbreak detection, and dynamic ecosystem monitoring are made possible by this bio-digital interface with previously unheard-of accuracy. Pest management systems can transition from reactive interventions to proactive, adaptive tactics by integrating biological sensing, Internet of Things (IoT) sensors and machine learning. In addition to increasing pest control effectiveness, this integration lessens reliance on pesticides, promoting ecological resilience and sustainable food production. Additionally, AI-enabled feedback loops enable ongoing learning from biological data streams, resulting in suggestions for farmers and stakeholders that are more context- specific. Scaling the use of biosensors, guaranteeing the reliability of biological data under changing field settings, and resolving ethical issues with bio-digital convergence are still difficult tasks. However, the combination of AI and live sensors could reinterpret pest control as an eco-centric, self- optimizing system, bringing technology development into line with agroecological principles. This multidisciplinary strategy promotes a balance between production, environmental stewardship, and global food security while providing a fresh route to precision agriculture.

Keywords: Bio-Digital Pest Management, biosensors, living sensors, precision agriculture and Resilience

EXPLORATION AND CATALOGING OF BEE-ATTRACTIVE FLORA

Devi M

Associate Professor, Agriculture Entomology, MIT College of Agriculture and
Technology, Musiri.

*Corresponding author Email: deviagri84@gmail.com

ABSTRACT

Bees play a crucial role in maintaining ecosystem balance and enhancing agricultural productivity through pollination. Understanding the diversity of flora that attracts bees is essential for conserving pollinator populations and improving habitat management. This study focuses on the exploration and cataloging of bee-attractive flora in a selected geographic region, aiming to document the variety of flowering plant species utilized by bees for nectar and pollen foraging. Field surveys were conducted across different habitats, including gardens, grasslands, forest edges, and agricultural zones, during peak flowering seasons. Flowering plant species were identified and recorded alongside bee visitation frequency, time of activity, and floral characteristics such as color, scent, and morphology. The study revealed a high diversity of plant species across families such as Asteraceae, Fabaceae, and Lamiaceae, which consistently attracted various bee taxa, including honeybees (*Apis spp.*), carpenter bees (*Xylocopa spp.*), and solitary bees. Seasonal variation in floral abundance and bee activity was also observed, highlighting the importance of continuous floral resources throughout the year. The findings provide valuable insights for pollinator conservation strategies, especially in designing pollinator-friendly landscapes and promoting native plant species that support bee diversity. The catalog developed through this research serves as a foundational resource for ecologists, conservationists, and agricultural planners aiming to enhance pollination services and biodiversity in both natural and managed ecosystems.

Keywords: Bee-attractive plants, pollinators, floral diversity, foraging behavior, native flora, biodiversity conservation

DEVELOPMENT OF A COMPREHENSIVE FLORAL CALENDAR

Devi M

Associate Professor, Agriculture Entomology, MIT College of Agriculture and
Technology, Musiri.

*Corresponding author Email: deviagri84@gmail.com

ABSTRACT

A comprehensive floral calendar is a vital tool for understanding the seasonal dynamics of flowering plant species and their availability to pollinators, particularly bees. This study aimed to develop a detailed floral calendar by documenting the phenology of bee-attractive plant species across different seasons in a selected region. Field surveys were conducted monthly over the course of one year to record the flowering patterns, peak blooming periods and abundance of nectar and pollen resources. Data were collected from various habitats, including agricultural fields, forest margins, home gardens, and natural grasslands. Each plant species was identified, and its flowering duration was mapped to create a month- by-month schedule of floral availability. The findings revealed distinct seasonal variations, with the highest diversity of flowering species observed during spring and late summer. Key plant families contributing to year-round bee foraging included Asteraceae, Fabaceae, Lamiaceae, and Malvaceae. The resulting calendar provides a visual representation of continuous floral resources, aiding in habitat management, crop pollination planning, and conservation efforts. It also helps identify floral gaps— periods with limited flower availability—allowing for targeted planting of supplementary species to support pollinators year-round. This comprehensive floral calendar can serve as a practical guide for farmers, ecologists and conservationists to enhance pollinator health and ensure sustainable agricultural productivity.

Keywords: Floral calendar, phenology, pollinators, bee forage, flowering plants, biodiversity conservation

DIGITAL TRACEABILITY SYSTEMS IN POULTRY PRODUCTION USING BLOCK CHAIN

Preethi V

M. Tech (Poultry Technology) Student, College of Poultry Production and Management,
Hosur, Tamil Nadu Veterinary and Animal Sciences University

*Corresponding author Email: preethibanu685@gmsil.com

ABSTRACT

Blockchain technology, built on the principles of decentralization, security, immutability, and smart contracts, is emerging as a powerful digital traceability system in poultry production. Its distributed data structure enables transparent, tamper-proof, and real-time recording of activities across the supply chain. Although challenges such as block size, scalability, latency, security, and privacy remain, blockchain has demonstrated wide potential in diverse sectors including supply chain management, medical records, finance, IoT, and food traceability. In poultry production, blockchain integrated with IoT devices captures data on feed quality, temperature, humidity, and bird health, which is securely stored on decentralized ledgers. This ensures accurate farm-to-fork traceability, enhancing food safety, fraud prevention, and consumer trust. Smart contracts further automate compliance, quality monitoring, and financial transactions, minimizing human error and disputes. Applications in the poultry, egg, and meat industries highlight its promise in rapid recall management, accountability, and sustainable practices. Despite challenges related to technical complexity, high implementation costs, regulatory frameworks, and adoption barriers, blockchain continues to advance rapidly and offers immense potential to transform poultry production. By ensuring transparency, accountability, and quality assurance, blockchain-driven traceability systems can significantly strengthen confidence among stakeholders and consumers.

Keywords: Block chain, Traceability, Poultry Supply Chain, Smart Contracts, Transparency, and Consumer Trust.

DIGITAL POLLINATOR–PEST BALANCE SYSTEMS: AI MODELS THAT OPTIMIZE PEST CONTROL WITHOUT HARMING POLLINATORS

Reshma M S ¹ and Baradhan G ²

¹ Research Scholar, Department of Agronomy, Faculty of Agriculture, Annamalai University

² Associate Professors, Agricultural College and Research Institute, Vazhavachanur

*Corresponding author Email: msreshmathirunallar@gmail.com

ABSTRACT

Agriculture today stands at a crossroads, where the urgent need to suppress pest populations collides with the equally critical necessity of conserving pollinators. The tension between these two objectives threatens both food security and ecological stability. The proposed Digital Pollinator–Pest Balance System (DPPBS) offers a transformative solution by integrating artificial intelligence (AI), real-time monitoring, and predictive analytics to manage insect populations in a balanced manner. Using computer vision and deep learning models, the system differentiates between pest and pollinator species with high accuracy, while Internet of Things (IoT) sensors track environmental variables influencing their activity. AI-driven simulations then forecast pest infestation risks alongside pollinator foraging schedules, thereby optimizing the timing and type of interventions. Rather than relying solely on chemical pesticides, the system can recommend targeted biological control agents, eco-friendly formulations, or precision-applied chemicals during pollinator-inactive periods. This minimizes non-target impacts while sustaining pollination services. Moreover, the system incorporates adaptive learning, where continuous field data refine the accuracy of species recognition and predictive thresholds over time. By combining ecological intelligence with digital technology, DPPBS promotes a resilient agricultural framework that enhances yield stability, protects biodiversity, reduces chemical dependency, and strengthens climate-smart farming practices. In essence, this model redefines Integrated Pest Management as a dynamic, pollinator-sensitive strategy, ensuring that food production systems remain both profitable and ecologically sustainable.

Keywords: Pollinator protection, AI-driven pest management, Digital agriculture, Precision ecology, Sustainable food systems.

HARNESSING INSECT BIODIVERSITY FOR SUSTAINABLE CROP PROTECTION

Shukla Saurab*¹ and Vishakha Sharma²

¹Research Scholar, Department of Agricultural Economics, Institute of Agricultural Science, Bundelkhand University, Jhansi, Uttar Pradesh, India

²Junior Research Assistant, Rani Lakshmi Bai Central Agricultural University, Jhansi, Uttar Pradesh, India

*Corresponding author Email: saurabhanilshukla@gmail.com

ABSTRACT

The idea of sustainable pest management has gained more attention in entomology and agricultural sciences. This is due to increasing evidence of pesticide resistance, ecosystem damage, and negative effects on helpful organisms. Many studies show that using chemical pesticides without care disrupts ecological balance, speeds up pest resurgence, and seriously harms pollinators and natural enemies. To address these problems, approaches that utilize natural processes and insect diversity are being promoted widely. Methods like biological control using predators, parasitoids, and entomopathogens; conserving and boosting natural enemies through habitat management; developing botanical and microbial pesticides; and including these in Integrated Pest Management (IPM) systems have shown successful results in various cropping systems. In addition to reducing reliance on synthetic products, these strategies improve essential ecosystem services such as pollination, soil health, and natural pest control. This, in turn, strengthens overall agro-ecosystem resilience. Moreover, combining traditional farming knowledge with modern entomological research offers solutions that are adaptable, environmentally friendly, and economically sound for farming communities. These comprehensive approaches support biodiversity, enhance food security, and promote agriculture that can withstand climate changes. By restoring ecological balance and reducing harm to non-target species, these advancements place entomology at the forefront of sustainable agricultural change. Tapping into nature's potential not only provides a way to control pests effectively but also lays the groundwork for future agricultural development and long-term sustainability.

Keywords: sustainable pest management, ecological innovations, biological control, integrated pest management, insect biodiversity, climate resilience

***Momordica cymbalaria* LEAF WATER EXTRACTIVES AGAINST THE WRIGGLERS
OF *Culex quinquefasciatus* SAY**

Yogapriya Adaikkan*¹ and Vikaash M²

¹ Guest faculty – Entomology, School of Agriculture and Animal Sciences, The Gandhigram
Rural Institute – DTBU, Gandhigram, Dindigul.

² Ph. D Scholar, Department of Entomology, Annamalai University

*Corresponding author Email: yogaento96@gmail.com

ABSTRACT

Mosquitoes are a serious threat to public health as they act as vectors helping on transmitting many diseases that can be morbid and fatal. Mosquito control is the major menace that rely to a maximum on chemical/ synthetic materials which are widely used in public and household areas where it creates a cause for several serious health issues. The effects are eerie as well. We planned a plant-based material for mosquito control and stepped in with a simple fresh water extractive of *Momordica cymbalaria* leaves. Mosquito management using fresh leaf extracts of *M. cymbalaria* against the wrigglers of *Culex quinquefasciatus* was experimented and the effects of the leaves extract on wrigglers and their corresponding sub-lethal and lethal doses were calculated along with their midgut effect on wrigglers. The sub-lethal (LC 50) and lethal (LC 95) concentrations were 1.96 and 4.24 percent. The midgut histology has shown complete loss of columnar cells, and complete break-down of plasma membrane, which shows potential insecticidal properties against mosquitoes.

Keywords: Mosquitoes; *Momordica cymbalaria*; Bio pesticides; Larvicidal; Midgut histology.

DIGITAL FRONTIERS IN IPM: PRECISION MANAGEMENT OF TEA MOSQUITO BUG (*Helopeltis* Spp.) IN CASHEW

Gopu Sushma^{*1}, Urati Mahesh², Kota Sahithi Chowdary¹ and Repaka Harika³

¹Department of Entomology, College of Agriculture, Rajendranagar, PJTAU

²Department of Horticulture, College of Agriculture, Rajendranagar, PJTAU Department of Entomology, IGKV, Raipur

^{*}Corresponding author Email: sushmareddygopu@gmail.com

ABSTRACT

The tea mosquito bug (*Helopeltis spp.*) is a major pest in cashew production, causing substantial yield losses of 40–60 per cent by feeding on young shoots, inflorescence and nuts leading to necrosis, nut drop and reduced quality. Conventional management relying on repeated chemical sprays is increasingly unsustainable due to the pest's rapid population growth, cryptic feeding behaviour and associated ecological and health risks. Precision driven Integrated Pest Management (IPM) approaches offer transformative solutions for sustainable control. Digital monitoring tools, including sensor- enabled sticky traps and image-based pest detection systems facilitate real time surveillance of pest populations and hotspot mapping. Decision support systems, integrating predictive analytics with weather based forewarning models enable timely and targeted interventions. Drone assisted application of biopesticides and beneficial natural enemies enhances field level efficiency while reducing labour and chemical inputs. Ecological strategies such as release of egg and nymphal parasitoids, neem based formulations and systematic removal of infested plant parts further strengthen pest suppression. Mobile enabled advisories and farmer focused digital platforms bridge knowledge gaps, promoting adoption of climate smart and environmentally sustainable practices. By integrating technological innovations with ecological principles, cashew protection systems can achieve precision, resilience and long term sustainability.

Keywords: *Helopeltis spp.*, Cashew, Precision IPM, Digital surveillance, Biological control and Climate-smart pest management.

ECOLOGICAL INNOVATIONS FOR SUSTAINABLE MANAGEMENT OF RED PALM WEEVIL *Rhynchophorus ferrugineus* IN COCONUT

Gopu Sushma*¹ and Urati Mahesh²

¹Department of Entomology, College of Agriculture, Rajendranagar, PJTAU

²Department of Horticulture, College of Agriculture, Rajendranagar, PJTAU

*Corresponding author Email: sushmareddygopu@gmail.com

ABSTRACT

The red palm weevil (*Rhynchophorus ferrugineus*), a devastating pest of coconut has emerged as a major threat to coconut production across tropical regions, causing severe damage to palms and yield losses that may exceed 50–60 per cent under favourable conditions. Conventional chemical control strategies are largely constrained by the pest's concealed larval stage within the trunk and fronds, rapid resistance development and associated environmental hazards. Ecological innovations provide promising avenues for sustainable management of this pest. Integrated pest management (IPM) approaches combining pheromone traps for mass trapping and monitoring, mechanical removal of infested tissues and application of botanicals such as neem extracts have demonstrated notable effectiveness. Biological control agents, including entomopathogenic nematodes, fungi and natural predators contribute to eco-friendly regulation of weevil populations. Climate-resilient strategies such as staggered planting, tolerant coconut varieties and regular field sanitation enhance system level sustainability, while predictive models based on weather and phenology enable timely interventions. Digital tools, including mobile based advisories and sensor assisted surveillance further support precision management. By integrating natural processes with climate-smart and digitally enabled strategies, coconut cultivation can transition towards a more resilient and environmentally sustainable pest management paradigm.

Keywords: *Rhynchophorus ferrugineus*, Coconut, Ecological pest management, Integrated Pest Management, Biological control, Climate-smart agriculture.

SOIL MICROBIOME ENGINEERING FOR RESISTANCE AGAINST RICE BROWN PLANTHOPPER (*Nilaparvata lugens*)

Repaka Harika^{1*}, Kota Sahithi Chowdary¹ and Gopu Sushma²

¹Department of Entomology, College of Agriculture, IGKV, Raipur

²Department of Entomology, College of Agriculture, Rajendranagar, PJTAU

*Corresponding author Email: harikarepaka20@gmail.com

ABSTRACT

The brown planthopper (BPH), *Nilaparvata lugens* is one of the most serious insect pests of rice, causing hopper burn, stunting and transmitting grassy stunt and ragged stunt viruses. Overuse of insecticides has led to resistance development, ecological disruption and resurgence of BPH, necessitating sustainable alternatives. Recent advances in soil microbiome engineering demonstrate its potential as an ecological tool to strengthen plant defenses against BPH. Beneficial microbes such as *Trichoderma harzianum*, *Pseudomonas fluorescens*, *Bacillus subtilis* and arbuscular mycorrhizal fungi colonize rice roots and induce systemic resistance by activating jasmonic acid and salicylic acid signaling. This leads to enhanced production of defense enzymes (polyphenol oxidases, peroxidases) and secondary metabolites that deter insect feeding. In addition, microbial volatile organic compounds (VOCs) disrupt BPH probing and oviposition, while certain rhizobacteria enhance silicon uptake, reinforcing plant cell walls. Innovations such as microbial consortia, seed bio-priming and biochar-based inoculants improve the survival and persistence of beneficial microbes under field conditions. Studies have shown that rice plants treated with *Trichoderma* and *Pseudomonas* exhibit up to 40–60 per cent reductions in BPH populations along with improved nutrient use efficiency and plant vigor. Metagenomic studies now identify keystone taxa linked to insect resistance, enabling the design of synthetic microbiomes tailored to rice ecosystems. Field evidence further demonstrates that microbiome-based interventions conserve natural enemies and reduce pesticide reliance, making them vital for regenerative rice production. In conclusion, soil microbiome engineering offers a frontier in sustainable pest management by coupling belowground biodiversity with aboveground pest suppression. This ecological approach can minimize insecticide dependence, enhance soil health, and ensure stable rice yields under intensive cultivation systems.

Keywords: Soil microbiome, *Nilaparvata lugens*, induced systemic resistance, beneficial microbes, rice IPM

**IOT-ENABLED SMART TRAPS FOR PRECISION SURVEILLANCE OF TOMATO
LEAF MINER (*Tuta absoluta*)**

Kota Sahithi Chowdary*¹, Repaka Harika¹ and Gopu Sushma²

¹Department of Entomology, College of Agriculture, IGKV, Raipur

²Department of Entomology, College of Agriculture, Rajendranagar, PJTAU

*Corresponding author Email: sahithikota1999@gmail.com

ABSTRACT

The tomato leaf miner (*Tuta absoluta*) is a highly invasive pest causing severe yield losses in tomato cultivation worldwide. Its rapid life cycle, high fecundity and ability to develop resistance to chemical insecticides make timely and accurate pest monitoring essential for effective integrated pest management (IPM). This study focuses on the development and deployment of IoT-enabled smart traps for real time surveillance of *T. absoluta*. These traps combine species specific pheromone lures with IoT sensors to automatically detect and quantify adult moth populations, transmitting data wirelessly to a centralized platform for continuous monitoring. Advanced data analytics and mapping tools enable farmers and researchers to identify pest hotspots, predict infestation trends and implement targeted interventions with minimal environmental impact. Field trials conducted across multiple tomato growing regions showed that smart traps significantly reduced manual monitoring labour, improved early detection accuracy and supported data driven pest management decisions. By integrating IoT technology with conventional pest control strategies, this approach contributes to sustainable agriculture, reduces economic losses and offers a scalable model for precision pest management applicable to other agricultural pests.

Keywords: *Tuta absoluta*, IoT, smart traps, precision pest management, tomato leaf miner, integrated pest management, real-time monitoring, sustainable agriculture.

BIOCONTROL POTENTIAL OF *Heterorhabditis bacteriophora* (POINAR) AGAINST RHINOCEROS BEETLE, *Oryctes rhinoceros* (L.)

Anjaly P V, Manu C R*, Reji Rani O P, Gavas R and Nisha M S

Department of Entomology, College of Agriculture, Vellayani, Kerala Agricultural University, Kerala, India

*Corresponding author Email: manudasc2014@gmail.com

ABSTRACT

The coconut rhinoceros beetle (CRB), *Oryctes rhinoceros* (L.) (Coleoptera: Scarabaeidae), is a major pest of coconut, causing over 10% yield loss. Adult beetles bore into unopened leaves and inflorescences, while grubs feed on decomposing organic matter in manure pits. In this study, the infectivity of the entomopathogenic nematode (EPN) *Heterorhabditis bacteriophora*, multiplied in black soldier fly larvae (BSFL), *Hermetia illucens* L., was evaluated against all three larval instars of CRB using different concentrations of infective juveniles (IJs). All three instars showed dose-dependent mortality, with no mortality at very low doses (100–300 IJs per insect). In the first and second instars, mortality began within 2–4 days and reached 100% by 6 DAT at 20,000 IJs and by 10 DAT at 10,000 IJs, while lower doses (500–5,000 IJs) caused partial mortality of 28–87% and 33–93%, respectively, by 12 DAT. The third instar was less susceptible, with mortality starting at 6–8 days and reaching only 40% at 20,000 IJs by 12 DAT. A t-test comparing nematodes produced from BSFL and wax moth larvae showed no significant difference in infectivity. These results confirm that *H. bacteriophora* is highly effective against early instars of CRB. Even though wax moth larvae (*Galleria mellonella*) are effective for nematode multiplication, BSFL offer a simpler, more cost-effective alternative host for multiplication of these nematodes, which act as a valuable tool for integrated pest management.

Keywords: Coconut rhinoceros beetle, EPN, *Heterorhabditis bacteriophora*, black soldier fly larvae

SEMIOCHEMICAL HIGHWAYS: USING PLANT–INSECT CHEMICAL SIGNALS FOR SUSTAINABLE PEST MANAGEMENT

Dhilipan K S ^{*1} and Baradhan G²

¹Research Scholar, Department of Agronomy, Faculty of Agriculture, Annamalai University

²Associate Professor, Agricultural College and Research Institute, Vazhavachanur

*Corresponding author Email: k.s.dhilipan001@gmail.com

ABSTRACT

Plants and insects communicate extensively through chemical signals, forming a complex network of semiochemicals that regulate ecological interactions. These signals, often termed semiochemical highways, play a pivotal role in mediating attraction, repulsion, and defense within agricultural ecosystems. Harnessing these natural chemical cues offers a promising alternative to pesticide-dependent pest management, aligning with the goals of sustainable and climate-smart agriculture. Semiochemicals such as pheromones, allomones, kairomones, and synomones can be deployed to manipulate insect behavior—either by disrupting pest mating cycles, repelling herbivores, or enhancing the activity of natural enemies. Advances in chemical ecology, coupled with precision application technologies, now make it feasible to exploit these interactions at the field scale. For instance, pheromone traps and attract-and-kill strategies reduce pest populations without harming pollinators or beneficial arthropods. Similarly, plants can be engineered or selected for enhanced volatile organic compound (VOC) emissions to attract parasitoids and predators, strengthening biological control. Integration of semiochemical strategies with digital agriculture such as remote sensing, AI-based insect monitoring, and automated release systems further enhances efficiency, allowing real-time, adaptive pest management. Unlike synthetic pesticides, semiochemical-based approaches leave minimal residues, reduce resistance development, and maintain ecosystem balance. As agriculture seeks resilient solutions to climate variability and pest adaptation, semiochemical highways provide a biologically grounded, environmentally friendly framework. By turning plant–insect communication into a management tool, we can shift from reactive pest suppression to proactive ecological regulation, ensuring both crop productivity and biodiversity conservation.

Keywords: Semiochemicals, Plant–insect interactions, Sustainable pest management, Chemical ecology, Biological control

TRAPPING TROUBLE: SUSTAINABLE CONTROL OF RHINOCEROS BEETLE IN COCONUT

Abishekkumar S ^{1*} and Selvanarayanan V²

¹PG scholar, Department of Entomology, Faculty of Agriculture, Annamalai University, Annamala Nagar, Cuddalore dt, 608002, Tamil Nadu, India.

²Professor & Head, Department of Entomology, Faculty of Agriculture, Annamalai University, Annamalai Nagar, Cuddalore dt, 608002, Tamil Nadu, India.

*Corresponding author Email: abishekkumars776@gmail.com

ABSTRACT

The Rhinoceros beetle (*Oryctes rhinoceros* L.) is a destructive pest of coconut, damaging unopened fronds and palm crowns, which leads to reduced vigour, yield loss, and in severe cases, plant mortality. Conventional insecticidal control poses environmental and health concerns, highlighting the need for eco-friendly alternatives. In the present study, the effectiveness of bucket traps baited with male aggregation pheromone (*Oryctes* lure contains ethyl 4-methyloctanoate) was evaluated as a component of Integrated Pest Management (IPM) against *O.rhinoceros*. Traps were installed within plantations at 4-6 traps per acre, monitored regularly, and combined with sanitation practices and biological control agents. The bucket traps effectively mass-trapped adult beetles, with a significant reduction in pest population observed over the cropping season. Integration with field sanitation (removal of decaying organic matter) and entomopathogenic fungi (*Metarhizium anisopliae*) further enhanced control efficiency. Compared with chemical-only treatments, the IPM approach incorporating bucket traps resulted in reduced infestation, healthier palms, and minimized pesticide use. These findings confirm that bucket traps serve as a practical, low-cost, and environmentally safe tool within an IPM framework for sustainable rhinoceros beetle management. Adoption of this strategy can support long-term productivity and ecological balance in coconut ecosystems.

Keywords: Rhinoceros beetle, Bucket traps, Aggregation pheromone, IPM, Biological control.

SYNERGISTIC EFFECTS OF MICROBIAL CONSORTIA IN INTEGRATED PEST MANAGEMENT

Kiruthiga N^{*1} and Sharvesh S²

¹Research scholar, Department of Agricultural Microbiology, Faculty of Agriculture, Annamalai University, Chidambaram, Tamil Nadu, India

²Research Scholar, Department of Horticulture, Faculty of Agriculture, Annamalai University, Chidambaram, Tamil Nadu, India

*Corresponding author Email: kiruthiga0110@gmail.com

ABSTRACT

Microbial biopesticides have appeared as a promising alternative to synthetically produced chemical pesticides, providing target-specific pest control methods that are environmentally friendly. Although studies have indicated that single strain applications of bacteria, fungi, or viruses have great potential, research is beginning to show the advantages of using microbial consortia. These mixtures contain a variety of organisms that use a variety of compatible mechanisms of action, such as the production of toxins and parasitism, as well as competition and induced systemic resistance to increase pest suppression. Not only do the interactions between these organisms increase efficacy across a wider variety of insects pests and potential diseases, but they also reduce the probability of pests developing resistance. Further, microbial consortia can better adapt to changing environmental conditions, which will help provide longer persistence and stability compared to using a single organism treatment. Developments in both biotechnology and formulation science have also allowed the composition of associated mixtures that can be active, while still maintaining decent shelf life and stability. By introducing these consortia into current pest management practices, we, as a group, can reduce the use of chemical pesticides and promote more sustainable agricultural practices, decrease pesticide impact on soil health, and promote biodiversity. Still challenges exist for the large-scale production or and regulatory approval of multi-microbe biopesticides. Assisting researchers in overcoming these hurdles via permission and policy will encourage researchs further use of microbials; bio pesticidal properties. Thus, the utilization of microbial consortia represents an innovative approach for integrated pest management along with global efforts to achieve food security while minimizing environmental impacts.

Keywords: Microbial consortia, biopesticides, synergistic interactions, integrated pest management, sustainable agriculture, pest resistance.

**EVALUATING THE DUAL ACTION EFFICACY OF LEMONGRASS OIL AGAINST
*Galleria mellonella***

**Vikaash M¹, Kanagarajan R^{1*}, Manikandan E¹, Mohammad Ikram ², Nishanthini K³,
and Yoga Priya A⁴**

¹Department of Entomology, Faculty of Agriculture, Annamalai University,
Chidambaram, Tamil Nadu, India

²Department of Entomology, Tamil Nadu Agricultural University,
Coimbatore, Tamil Nadu, India

³Pandit Jawaharlal Nehru College of Agriculture and Research Institute, Karaikal

⁴School of Agriculture and Animal Sciences, The Gandhigram Rural Institute – DTBU,
Gandhigram, Dindigul

*Corresponding author Email: mvikaash123@gmail.com

ABSTRACT

The greater wax moth, *Galleria mellonella*, is a significant economic pest in apiculture, causing extensive damage to honeybee combs. The reliance on chemical fumigants raises concerns about honey and wax contamination, necessitating the exploration of safe and sustainable alternatives for pest management. This study evaluated the dual action efficacy of *Cymbopogon citratus* oil, a widely available botanical, as both a larvicide and an oviposition repellent against *G. mellonella*. A laboratory study was conducted to assess five treatments, spray emulsions at 2.5%, 5%, and 10%, pure oil vapor treatment and control. Larval mortality and adult oviposition were quantified to establish a dose- response relationship. The results revealed a strong, dose dependent efficacy. The 10% spray achieved 98.0% larval mortality and almost completely inhibited egg-laying. Notably, the pure oil vapor treatment resulted in 99.5% larval mortality and complete repellence, demonstrating the profound potential of fumigation. The 5% concentration also proved highly effective, suggesting a viable threshold for practical control. We conclude that lemongrass oil is a potent biopesticide, and its application as a fumigant presents a highly effective, practical and ecologically sound strategy for beekeepers to protect colonies from wax moth infestation.

Keywords: Greater wax moth, Lemongrass, Botanical Pesticide, Repellent, Fumigant.

**HABITAT MANIPULATION STRATEGIES FOR SUSTAINABLE MANAGEMENT
OF RUGOSE SPIRALING WHITEFLY *Aleurodicus rugioperculatus* IN
COCONUT**

Sumalatha B V^{1,2*}, Selvaraj K¹ and Kandan A¹

¹ Division of Germplasm Conservation and Utilization, ICAR-National Bureau of
Agricultural Insect Resources, Bengaluru-560024, Karnataka, India

² Department of Entomology, Dr. Rajendra Prasad Central Agricultural
University, Pusa, Samastipur - 848 125, Bihar, India

*Corresponding author Email: sumachaaru55@gmail.com

ABSTRACT

Coconut palm (*Cocos nucifera* L.) suffer variety of insect pest infestations, notably the recent invasive rugose spiraling whitefly (RSW), *Aleurodicus rugioperculatus* Martin (Hemiptera: Aleyrodidae), which is an economically significant pest. Further, this pest also coexists with subsequent invasive whitefly species viz., Bondars nesting whitefly, *Paraleyrodes bondari*, nesting whitefly, *P. minei* and palm infesting whitefly *Aleurotrachelus atratus*. Numerous natural enemies have been documented in association with RSW. Among, the most effective are the parasitoid *Encarsia guadeloupae* Viggiani (Hymenoptera: Aphelinidae) and the predator *Apertochrys aastur* (Banks) (Neuroptera: Chrysopidae). However, their use alone to suppress this whitefly may be difficult under severe outbreak conditions; therefore, integration of compatible biocontrol-based management strategies is ideal to develop state specific protocol. Hence, bio-intensive integrated pest management (BIPM) is considered the desirable path for achieving sustainability and which reduces chemical input costs and on-farm and off-farm environmental impacts. Two BIPM modules and conventional agronomic practices were validated under field condition. Results revealed that BIPM modules-I was found superior in managing RSW, which consists of standard recommended agronomic practices; intercropping with banana; installation of yellow sticky traps @ 30 traps /ha; inundative release of *A. astur* @ 1000 eggs /ha; inundative release of *E. guadeloupae* @ 1000 adults /ha and foliar application of *Isaria fumosorosea* @ 5g/L, with the maximum overall pest reduction (75.88- 80.24%). Besides, highest natural parasitism (67.36- 81.45%) and population of *A. astur* (5.89- 6.85 grub/leaf) were maximum in BIPM module-1 than farmers practices and other BIPM modules. Therefore, farmers and other stakeholders were sensitized for extensive promotion of BIPM module to achieve greater adoption for management of RSW in coconut.

Keywords: Invasive whiteflies, Coconut, Habitat manipulation, *Cana indica*, banana, intercrop

**FROM HARASSING NATURE TO HARNESSING NATURE – TOWARDS
ECOFRIENDLY PEST MANAGEMENT**

Sriram K

Undergraduate Scholar (Agriculture)

SRM College of Agricultural Sciences, SRM Institute of Science and Technology,

Chengalpattu – 603 201, Tamil Nadu, India

Corresponding author Email: sk3448@srmist.edu.in

ABSTRACT

The transition “*From Harassing Nature to Harnessing Nature – Towards Ecofriendly Pest Management*” reflects the urgent need to shift human practices from exploitative to sustainable. Over the past decades, the excessive use of synthetic chemicals in agriculture has caused severe ecological imbalance. According to the U.S. FAO Pesticide Use and Trade Report (1990–2020), an average of 30 million tons of pesticides have been dumped into soils globally, leading to soil degradation, contamination of water resources, loss of biodiversity, and adverse effects on human health. These persistent chemicals disrupt soil microbiota, reduce crop productivity, and accumulate in the food chain, posing chronic health hazards. To mitigate these impacts, Integrated Pest Management (IPM) offers an eco-friendly framework combining cultural, mechanical, and biological control methods. Cultural practices such as crop rotation, field sanitation, and resistant varieties reduce pest incidence; mechanical methods like trapping and hand-picking directly remove pests; and biological approaches harness natural enemies like parasitoids, predators, and entomopathogenic microbes. Innovative strategies have further strengthened IPM—such as the use of *Beauveria bassiana* and pheromone trapping for coffee berry borer (*Hypothenemus hampei*) management, and nucleopolyhedrovirus (NPV) and egg parasitoids like *Trichogramma pretiosum* for fall armyworm (*Spodoptera frugiperda*) suppression. These examples illustrate how modern science is aligning with nature’s mechanisms, marking the transformation from harming ecosystems to harnessing their resilience for sustainable agricultural development.

Keywords: Sustainable agriculture, Integrated Pest Management (IPM), pesticide pollution, soil contamination, environmental health, coffee berry borer, fall armyworm, ecosystem resilience.

RNAi-BASED BIOPESTICIDE: A TARGETED APPROACH FOR CONTROLLING THE INVASIVE GIANT AFRICAN SNAIL

Anjala Leelu Ani

Undergraduate scholar (Agriculture)

SRM College Of Agricultural sciences, Baburayanpettai, Chengalpettu – 603201, India

Corresponding author Email: al3111@srmist.edu.in

ABSTRACT

RNA interference (RNAi) technology holds promise as a biopesticide approach against the invasive giant African snail (*Lissachatina fulica*), targeting genes critical for survival, growth, or reproduction through a highly specific gene-silencing process. The core mechanism involves identifying unique essential genes, designing double-stranded RNA (dsRNA) triggers to match these targets, and encapsulating dsRNA in protective nanoparticles to ensure efficient delivery and stability in the field. Bait formulations facilitate snail uptake and subsequent gene silencing, resulting in mortality if targeted proteins are vital. However, challenges remain in ensuring species specificity, overcoming biological barriers to oral delivery, confirming efficacy in *Lissachatina fulica*, and maintaining dsRNA stability under environmental conditions. Regulatory approval and public acceptance will also be crucial before wide deployment. While laboratory RNAi experiments in other snail species demonstrate potential, more research is needed for scalable, field-ready RNAi-based biopesticides targeting the giant African snail, highlighting their future role as targeted, environmentally friendly alternatives to chemical controls.

Keywords: RNAi, Biopesticide, African snail, Nanoparticles

**ECOLOGICAL INNOVATIONS FOR SUSTAINABLE MANAGEMENT OF FRUIT
FLY (*Bactrocera cucurbitae*) IN CUCURBITS**

Rajani D^{*1}, Urati Mahesh¹, Suresh Kumar K² and Harika M¹

¹Department Of Horticulture, College of Agriculture, Rajendranagar, PJTAU

²YFA KVK, Madhnapuram, Dist, Wanaparthy

*Corresponding author Email: rajani.horti@gmail.com

ABSTRACT

The melon fruit fly (*Bactrocera cucurbitae*), a destructive pest of cucurbitaceous vegetables, has emerged as a persistent challenge across major production systems, causing yield losses that may exceed 70 per cent under favourable conditions. Conventional chemical management remains largely ineffective due to the pest's concealed larval feeding within fruits and its high potential for resistance development, while indiscriminate pesticide use further exacerbates ecological and food safety concerns. In this context, ecological innovations are pivotal for sustainable suppression. Integrated pest management (IPM) frameworks incorporating pheromone and cue-lure traps for mass trapping and population monitoring, field sanitation through systematic destruction of infested fruits, and deployment of botanicals such as neem-based formulations have demonstrated significant efficacy. Biological control agents, including parasitoids of fruit fly pupae and entomopathogenic fungi, offer additional avenues for environmentally benign regulation. Climate-resilient strategies such as crop rotation, synchronized sowing, and cultivation of tolerant varieties enhance system-level resilience, while weather-based pest forewarning models can guide proactive interventions. Digital innovations ranging from mobile-enabled advisories to sensor-driven surveillance provide precision support to farmers, bridging ecological approaches with timely decision-making. By harnessing natural processes and embedding them within climate-smart and digitally enabled pest management systems, the cucurbit sector can transition toward a more resilient and sustainable crop protection paradigm.

Keywords: *Bactrocera cucurbitae*, Cucurbits, Ecological pest management, Integrated Pest Management, Biological control and Climate-smart agriculture.

BIO INSPIRED ROBOTIC MODEL (ORNITHOPTERA)

Ilakkiya V

KPR Institute of Engineering and Technology, Coimbatore

Corresponding author Email: ilakkiya.v.2006@gmail.com

ABSTRACT

Crop depredation by birds in Agricultural field (Nearly 35 % to 65% of crops are destroyed by crop eating birds like sparrow, pigeon, parrot etc.,) is a serious threat to agriculture sector. Traditional methods of scaring birds away use a lot of resources, which harms the environment and creates problems for farms. Crops are still being damaged, highlighting the need for long-lasting solutions that protect the environment and maintain agricultural production. Therefore farmers need Eco-friendly modern bird scaring technique to save their crops. This innovative device harnesses the power of biomimicry, emulating the flight patterns and predatory behavior of natural avian predators to effectively deter nuisance birds from agricultural fields. It autonomously patrols the skies above fields, creating a dynamic and realistic threat presence that encourages birds to seek safer feeding grounds. With its non-lethal and environmentally friendly approach, the ornithopter offers farmers a sustainable solution to protect crops and minimize bird- related damage, ensuring optimal yields and agricultural productivity. The adoption of bird-inspired strategies represents a crucial advancement in the sustainability and precision of crop protection systems. Biological entity/ process inspired from: The meticulously crafted robotic bird Ornithopter is bio inspired/mimicked, by adopting elements from indigenous birds like falcons, vultures, and eagles. By precisely imitating their look and predatory behaviors, Ornithopter is designed to effectively discourage crop-eating birds in agricultural environments. Ornithopter offers an ethical and efficient solution by imitating nature's techniques, and safeguarding crops while minimizing its adverse impacts on the environment. Analyzing avian predators, constructing an affordable 3D-printed framework, and incorporating motors, sensors, and solar panels are all necessary steps in the construction of a Ornithopter prototype. Programming features include autonomous operation, identification of birds, and lifelike wing movements. Field testing determines the success rate with bird repellent, directing successive improvements. Energy efficiency balancing, choosing the right materials, successful bird authentication, and Ornithopter adaptation to various agricultural contexts are among the challenges. To develop a profitable, effective, and resilient concept that safeguards crops while consuming the fewest resources possible while complying with both environmental and agricultural targets, these issues must be addressed.

Keywords: Sustainable agriculture, Integrated pest management, Automated machineries

**ECO-FRIENDLY BOTANICALS TREATMENT AND SMART PACKAGING:
SUSTAINABLE MANAGEMENT OF *Rhyzopertha dominica* ON STORED
WHEAT**

Mekha M Prasad ^{*1} and Ghelani M K²

¹M. Sc. (Agri.) Scholar, Department of Entomology, College of Agriculture, Junagadh
Agricultural University, Junagadh Pin Code: 362001.

²Associate Professor, Department of Entomology, College of Agriculture, Junagadh
Agricultural University, Junagadh Pin Code: 362001.

*Corresponding author Email: mekhagirija@gmail.com

ABSTRACT

The present investigation was undertaken at the Post Graduate Laboratory, Department of Entomology, Junagadh Agricultural University, Junagadh, Gujarat, during 2024–2025 to evaluate the efficacy of different botanicals, inert dusts as seed protectants and packaging materials against the lesser grain borer, *Rhyzopertha dominica* (F.) in stored wheat. The pooled data on adult mortality revealed significant variation among botanicals, with a marked decline in mean adult mortality over time, from 86.11% after one month to 36.84% after nine months. Among the botanicals tested, azadirachtin 1.0 EC @ 1.5 ml/kg seed was most effective, consistently recording the highest adult mortality, followed by neem oil @ 5 ml/kg, neem seed kernel powder @ 5 g/kg and neem dry leaf powder @ 5g/kg. Castor oil and groundnut oil (each @ 5 ml/kg) showed moderate efficacy, while custard apple seed powder, mint leaf powder and talc (each @ 5 g/kg) were least effective. A similar trend was observed across all storage intervals from one to nine months. Population growth studies after nine months further substantiated the superior efficacy of azadirachtin, which was at par with neem oil, while NSK powder and neem dry leaf powder showed intermediate protection. Castor and groundnut oil were statistically at par and the least effective treatments were custard apple seed powder, mint leaf powder and talc, which supported higher pest multiplication, with talc only slightly better than the untreated control. In the evaluation of packaging materials, considerable variation in adult emergence, grain weight loss and protection was observed after nine months. Gunny and cloth bags supported heavy infestation, with the gunny bag showing the highest adult emergence and maximum weight loss. HDPE bags offered superior protection, consistently registering the lowest adult emergence and weight loss due to their impermeable structure, while polypropylene bags performed next best, significantly better than polyethylene, which showed moderate susceptibility. Overall, the study highlights azadirachtin and neem oil as promising botanical protectants and HDPE bags as the most effective packaging material for long-term safe storage of wheat against *R. dominica*.

Keywords: Botanicals, Lesser grain borer, *Rhyzopertha dominica* (F.), Packaging materials, Seed protectant and Wheat

MANAGEMENT OF FRUITFLIES *Bactrocera dorsalis* IN MANGO ECOSYSTEM OF A.P

Sravanthi G ¹, Lakshmi B K M ² and Radha Rani K³

¹Dr.YSRHU-Cashew Research Station, Bapatla,

²Dr.YSRHU-Mango Research Station, Nuzvid,

³Dr.YSR Horticultural University, Venkataramannagudem, West Godavari district.

*Corresponding author Email: sravanthiguntupalli@gmail.com

ABSTRACT

Fruit flies are considered as most obnoxious pests damaging the crops globally. Fruit flies result in significant damage from 20-40% to fruits and vegetables in field and post harvest scenarios. Among different varieties grown in Andhra Pradesh, Banganapalli, Totapuri are highly popular mango varieties utilised for table and for processing ready to serve juice. But unfortunately these varieties are highly susceptible to fruitfly attack causing huge economic losses to the farmers. During Covid-19 lock down period (2020, 2021 and 2022) there is delay in harvesting mango fruits in Andhra Pradesh leading to multiplication of fruitflies (*Bactrocera dorsalis*). Offseason rains during March, 2023 and April, 2023 coupled with residual fruitfly population due to delayed harvestings during Covid-19 era resulted in severe fruitfly outbreak in mango ecosystem in A.P. Farmers are facing lot of problem due to fruitfly damage in mango which directly affects the final yields. Hence the experiment was planned and initiated during 2022-2023 with 7 treatments at Dr.YSRHU-Mango Research Station, Nuzvid, Andhra Pradesh (16.7876° N, 80.8490° E). Among treatments T 1 was installation of Methyl Eugenol traps which is common among the treatments. Among the different treatments evaluated against the fruitflies Spinosad 45 SC @ 0.35ml/L and Azadirachtin 10,000 ppm @ 1ml/L when sprayed after 30 and 50 days after fruitset were found to be effective in reducing the pest population with minimum damage percentage of 7.3 and 8.3 followed by Decamethrin 0.5ml/L+ Azadirachtin 10000 ppm @ 2ml/L with percent damage of 9.3 compared to the untreated control with 36.3 percent damage. T 1 (Methyl Eugenol traps) alone was not effective in controlling the fruitfly population in mango ecosystem and upto 18.3 percent damage was recorded. Fruitflies trapped in various treatments were recorded at different Standard Meteorological Weeks (SMW). Fruitfly specimens were identified by Dr.David, Entomologist from NBAIR, Bengaluru and confirmed the species as *Bactrocera dorsalis* (Hendel). There is no significant difference in the oviposition marks observed after the sprayings and there is no significant difference in the fruitflies trapped in the Methyl Eugenol traps installed in various treatments. Hence before harvest and after harvest parameters were taken into consideration and percent damage was recorded. Fruit fly management in mango involves summer ploughing, clean cultivation and

removal and destruction of damaged and fallen fruits below the trees, proper disposal by burning or disposing in deep furrows 50 cm below the ground, Erection of Methyl eugenol traps @ 5/acre, by adjusting harvest timing, especially by harvesting fruits at the green-mature stage or at the yellow-point onset as fruit fly infestation was closely related to changes in the peel texture (firmness, in particular) and chlorophyll fluorescence of the fruit as it ripened and by spraying Spinosad 45 SC @ 0.35ml/L or Azadirachtin 10,000 ppm@ 1ml/L. Management of fruitfly is the need of the hour so that the farmer will be benefitted economically.

Keywords: Management, fruitflies, *Bactrocera dorsalis*, mango

**COLD PLASMA SEED TREATMENT: AN INNOVATIVE APPROACH FOR
SUSTAINABLE PEST MANAGEMENT IN RICE**

Malini B ^{1*} and Vanitha J ²

¹PG Scholar, Department of Genetics and Plant Breeding, SRM College of Agricultural Sciences, SRMIST, Baburayanpettai, Chengalpattu District, Tamil Nadu– 603201

²Asst. Professor, Department of Genetics and Plant Breeding, SRMCAS, SRMIST, Baburayanpettai, Chengalpattu District, Tamil Nadu– 603201

*Corresponding author Email: bm1467@srmist.edu.in

ABSTRACT

Rice (*Oryza sativa* L.) is a globally significant staple crop, but its productivity is often limited by various pest pressures. The growing need for sustainable agriculture calls for new, environmentally friendly ways to reduce pest damage while using fewer chemical pesticides. Cold plasma seed treatment has become a promising non-thermal technology that can make rice more resistant to pests. This method uses ionized gases to make reactive oxygen and nitrogen species (ROS/RNS), which are strong signaling molecules that start stress-responsive and defense-related pathways at the molecular and cellular level. Exposing rice seeds to cold plasma has been demonstrated to enhance germination kinetics, seedling vigor, and early establishment, while also priming defense mechanisms against biotic stresses. Cold plasma treatment can cause changes in the epigenome, such as changes in DNA methylation patterns and histone remodeling. These changes can affect the transcription of important pest-resistance genes. These epigenetic effects could have benefits for future generations, making crops more resistant to pests in the next crop cycle. It is possible to create a sustainable, non-GMO way to control pests that fits with the principles of ecological farming by adding cold plasma seed treatment to rice growing systems.

Keywords: Cold plasma, rice, seed treatment, sustainable pest management, reactive oxygen species, epigenetic priming, eco-friendly crop protection.

***Thrips parvispinus* KARNY (THYSANOPTERA: THRIPIDAE): AN EMERGING PEST OF CHILLI IN KERALA**

Anjana S A ^{*1}, Thania Sara Varghese ¹ and Santhosh Kumar T ²

¹PG Scholar, Department of Agricultural Entomology, College of Agriculture Vellayani

²Assistant Professor, Dept. of Agricultural Entomology, College of Agriculture, Vellayani

*Corresponding author Email: anjanasankar2022@gmail.com

ABSTRACT

Chilli (*Capsicum annuum* L.), an essential commercial vegetable, is widely regarded as a versatile and remarkable spice. In 2020, India one of the top consumers of chilli—generated over ₹130 billion from its cultivation. Among the 60 insect pests known to affect chilli crops, thrips, especially the invasive species *Thrips parvispinus* (Karny), have emerged as a major threat to Indian agriculture. Initially detected on Carica papaya in 2015 and later on Dahlia rosea Cav., this pest was most recently observed infesting chilli plants in Kerala in 2022. A severe outbreak in 2021 resulted in devastating yield losses ranging from 80% to 100% across southern India. Hence, a rapid roving survey was carried out to assess thrips infestation in chilli crops across farmer's fields throughout Kerala. Thrips specimens were collected from the flowers of infested chilli plants. The predominant species was identified as *T. parvispinus* using the taxonomic key provided by ICAR-NBAIR. Both adult and nymph stages of *T. parvispinus* were found colonizing the underside of leaves and flowers, causing severe curling and crinkling of young leaves at the growing tips. This species exhibits sexual dimorphism, with females having a brown body and males appearing completely yellow. Additionally, other thrips species such as *Scirtothrips dorsalis*, *Frankliniella schultzei*, *Thrips palmi* and *Thrips hawaiiensis* were also detected in the flower samples. A field study was carried out to assess the damage caused of *T. parvispinus* infestation on chilli crops. The experiment compared the levels of flower drop, fruit drop, and fruit damage between plots treated with insecticide (three applications of Thiamethoxam 25 WG at 25 g a.i./ha during the 6th, 8th, and 10th weeks after transplanting) and untreated control plots. The findings revealed a marked reduction in damage caused by *T. parvispinus* in the treated plots. Flower drop was lowered by 74.42%, 85.18% and 32.37% at 7, 14 and 21 days after treatment (DAT), respectively. Likewise, fruit drop decreased by 52.98%, 67.21% and 52.23% over the same periods. Fruit damage also showed a significant decline, with reductions of 70.36%, 75.60% and 67.93% at 7, 14 and 21 DAT, respectively. Overall, these findings on bioecology of *T. parvispinus* provide critical insights for the development of integrated pest management approaches for mitigating the threat posed by *T. parvispinus* to chilli and other horticultural crops.

Keywords: Sustainable agriculture, Integrated pest management, Bioecology

RESISTANCE STUDIES IN GREEN GRAM (*Vigna radiata*) PARENTAL LINES AND F1 HYBRIDS AGAINST *Aphis craccivora* KOCH

Darlin Prakash R^{*1} and Selvanarayanan V²

¹Research Scholar, Department of Entomology, Faculty of Agriculture, Annamalai University

² Professor and Head, Department of Entomology, Faculty of Agriculture, Annamalai University.

*Corresponding author Email: darlinprakash@gmail.com

ABSTRACT

Green gram (*Vigna radiata*), commonly known as mung bean, is an important legume crop valued for its edible seeds but is highly susceptible to aphid (*Aphis craccivora* Koch) infestation. This polyphagous pest extracts sap from leaves, stems, and pods, thereby reducing photosynthate availability and crop productivity. To address this challenge, six promising green gram genotypes were utilized in a diallel crossing program, resulting in 30 F₁ hybrids that were field-screened for resistance against aphids. Promising hybrids identified from field evaluation were further subjected to laboratory assays to determine the underlying resistance mechanisms. Settling preference studies of *A. craccivora*, involving both free-choice and no-choice tests, revealed reduced aphid settlement on genotype IC-103833 and hybrid P₂ × P₃ under free-choice conditions, while IC-39272-1 and hybrid P₆ × P₃ exhibited low settling under no-choice conditions. Aphid settling preference showed a significant negative association with trichome density on stems and leaf surfaces (adaxial and abaxial). Furthermore, a negative correlation was observed between aphid settlement and leaf thickness. A similar correlation was also observed with the wax content of leaves and pods. Among the evaluated entries, hybrids P₂ × P₃ and P₆ × P₃ consistently recorded higher yields with reduced pest incidence, indicating their potential for resistance breeding in green gram.

Keywords: Green gram, Diallel, Hybrid, Resistance, Aphids, Settling preference.

NANOCARRIER-MEDIATED RNAi FOR SUSTAINABLE FRUIT FLIES MANAGEMENT

Manojkumar S ¹, Kiruthiga N ¹, Sharvesh S ^{2*} and Vigneswari G²

¹ Department of Agricultural Microbiology, Faculty of Agriculture, Annamalai University

² Department of Horticulture, Faculty of Agriculture, Annamalai University, Annamalai Nagar, 608002, Tamil Nadu, India

*Corresponding author Email: sharveshkeerthi3@gmail.com

ABSTRACT

Fruit flies (*Bactrocera spp.* and *Ceratitis spp.*) represent one of the most destructive pest groups in fruit crops, leading to heavy yield losses and posing barriers to international trade due to quarantine restrictions. Conventional pesticide-based strategies are increasingly ineffective because of resistance development, environmental hazards, and concerns over chemical residues. RNA interference (RNAi) offers a highly specific and environmentally benign alternative for insect pest management. Nevertheless, practical field deployment faces critical challenges such as dsRNA instability, rapid degradation under environmental conditions, and limited uptake by insect pests. This study focuses on nanocarrier-mediated delivery systems to enhance the stability, bioavailability, and efficiency of dsRNA in targeting essential genes of fruit flies. Biodegradable nanocarriers including chitosan nanoparticles, lipid nanostructures, and layered double hydroxides will be explored for encapsulation and delivery. Evaluation will be carried out in terms of gene silencing efficiency, insect mortality, crop safety, and postharvest residue levels. The integration of RNAi technology with nanodelivery platforms provides a cutting-edge, eco-friendly, and residue-free approach to pest management in fruit crops. This innovation could significantly reduce pesticide reliance and ensure sustainable production systems with enhanced export competitiveness.

Keywords : RNA interference (RNAi); nanocarriers; dsRNA; fruit flies; *Bactrocera dorsalis*; chitosan nanoparticles; gene silencing; sustainable pest management; fruit crops

BIOEFFICACY OF ENTOMOPATHOGENS AGAINST THE SUCKING PESTS OF OKRA

Dharani V^{1*} and Kumar K²

¹ Ph.D. Scholar, Department of Entomology, Faculty of Agriculture, Annamalai University, Annamalai Nagar, Tamil Nadu- 608 002

² Professor and Head Department of Entomology, Pandit Jawaharlal Nehru College of Agriculture and Research Institute, Karaikal – 609 603, U. T. of Puducherry

*Corresponding author Email: dharaniv606@gmail.com

ABSTRACT

Okra is cultivated in tropical, subtropical and warm temperate regions around the world. The productivity is low due to many factors and one of the most important constraint is the attack of insect pests. Okra is attacked by a number of insect pests results in lower yields. Recently the sucking pests are becoming major pests under changing climatic condition coupled with application of injudicious and spurious pesticides which cause considerable yield loss. The sucking pest complex consisting of aphids (*Aphis gossypii* Glover), leafhopper (*Amrasca biguttula biguttula* Ishida) that caused 17.46 per cent yield loss in okra. Chemical insecticides are sprayed which leads to several problems like toxic residues, elimination of natural enemies, environmental disharmony and development of resistance. Biological control of insect pests using different entomopathogenic microorganisms is gaining importance due to their target specificity, self-perpetuity and obvious safety to the environment. Two field experiments were conducted at the Eastern farm of Pandit Jawaharlal Nehru College of Agriculture and Research Institute, Karaikal during rabi and kharif 2023 to evaluate the effectiveness of thiamethoxam 25 WG and a mixture of entomopathogens with (*B. bassiana*, *M. anisopliae*, *V. lecanii*) against the sucking pests. Leafhoppers population was low in thiamethoxam 25 WG @ 0.4 g/l (5.93 and 5.79 number/ 3 leaves) with a reduction of (73.85 and 71.23%) followed by *B. bassiana* + *M. anisopliae* + *V. lecanii* @ 15 g/l (6.89 and 6.55 number/ 3 leaves) with a reduction of 69.62 and 67.46 per cent and for the aphids the overall mean population was low in thiamethoxam 25 WG @ 0.4 g/l (4.74 and 4.70 number/ 3 leaves) with a reduction of (72.37 and 71.54%) followed by *B. bassiana* + *M. anisopliae* + *V. lecanii* @ 15 g/l (5.72 and 5.52 number/ 3 leaves) with a reduction of (66.66 and 67.00%) during both field experiments. The mean population of predatory coccinellids was low in the treatment with thiamethoxam 25WG @ 0.4 g/l (1.33 and 1.18/plant) while a higher population was recorded in the treatments with entomopathogens which ranged from 1.47 to 1.80/plant and 1.45 to 1.76/plant and the mean population of spiders was low in the treatment with thiamethoxam 25WG @ 0.4 g/l (1.52 and 1.58/plant) while a higher population was recorded in the treatments

with entomopathogens which ranged from 1.75 to 2.10/plant and 1.81 to 2.08/plant.

Keywords: Entomopathogens, natural enemies, okra, phytotoxicity, sustainability.

HARNESSING NATURE FOR SUNFLOWER SECURITY: ECO-INNOVATIVE IPM APPROACHES AGAINST CAPITULUM BORER (*Helicoverpa armigera*)

Kiruthika M ^{1*} and Selvamuthukumaran T ²

¹Ph.D Scholar, Department of Entomology, Faculty of Agriculture, Annamalai University
Tamil Nadu, India.

²Associate Professor, Department of Entomology, Faculty of Agriculture, Annamalai
University Tamil Nadu, India.

*Corresponding author Email: murugesankiruthika1@gmail.com

ABSTRACT

Sunflower, a premier oilseed crop, is under persistent threat from the capitulum borer (*Helicoverpa armigera*), which inflicts severe yield losses. Conventional reliance on broad-spectrum insecticides has triggered resistance, pest resurgence, and environmental hazards, necessitating ecologically sound alternatives. To address this challenge, an innovative field study was undertaken during kharif season at Tiruchirappalli, Tamil Nadu, to evaluate Integrated Pest Management (IPM) modules tailored for sunflower. The modules strategically combined eco-friendly options: seed treatment with imidacloprid, foliar application of neem seed kernel extract (NSKE), *Helicoverpa armigera* nucleopolyhedron virus (HaNPV), and reduced-risk insecticides spinosad and emamectin benzoate. Applications were scheduled at critical crop stages to suppress larval infestation effectively. This holistic approach not only minimized capitulum borer incidence but also enhanced ecological balance by reducing dependence on synthetic chemicals. The integration of botanicals, biopesticides and selective molecules represents a nature-aligned innovation that empowers farmers with sustainable and scalable pest management solutions. The findings underscore the potential of eco-innovative IPM frameworks in securing sunflower productivity, safeguarding biodiversity, and moving towards resilient oilseed production. Such strategies exemplify the paradigm shift from pesticide-reliant protection to nature-driven sustainability in Indian agriculture.

Keywords: Sunflower, *Helicoverpa armigera*, Eco-IPM, Biopesticides, Sustainability

**ADJUSTING PLANTING DATES AS AN ECOLOGICAL INNOVATION FOR
SUSTAINABLE PEST MANAGEMENT IN RICE**

Yogapriya G ^{1*} and Kandibane Muthusamy²

¹ Ph.D Scholar, Department of Entomology, Faculty of Agriculture, PAJANCOA and RI
Karaikal, Puducherry

² Professor, Department of Entomology, Faculty of Agriculture, PAJANCOA and RI
Karaikal, Puducherry

*Corresponding author Email: g.yoga2304@gmail.com

ABSTRACT

Field experiments were conducted at the Eastern Farm of PAJANCOA & RI, Karaikal, during Rabi and Kharif 2023 to evaluate the effect of varying planting dates on the incidence of key insect pests of rice and the abundance of their associated natural enemies. Three schedules - early, normal, and late planting were compared to assess their role in pest regulation under field conditions. Early planting consistently registered the lowest incidence of stem borer, and leaf folder, indicating that a favorable crop–pest synchrony minimized infestation levels. In contrast, late planting exhibited a pronounced build-up of these pests, leading to greater crop injury and potential yield reduction. Observations on natural enemies revealed that coccinellids and spiders were slightly more abundant in late-planted fields, yet their overall population density remained insufficient to balance pest outbreaks. The study clearly demonstrated that adjusting planting dates, particularly by adopting early planting schedules, can significantly suppress pest populations, reduce crop losses, and contribute to yield stability without additional inputs. This cultural strategy serves as a simple, low-cost, and environmentally sustainable alternative to sole reliance on chemical pesticides. Moreover, it complements integrated pest management (IPM) principles by conserving beneficial organisms, lowering production risks, and enhancing agroecosystem resilience. The findings underscore that fine-tuning planting time represents an important ecological innovation in rice cultivation, offering farmers a practical pathway toward sustainable intensification and reduced ecological footprint in pest management.

Keywords: rice, planting dates, thrips, stem borer, leaf folder, IPM

MICROBIOME-DRIVEN APPROACHES TO ECO-FRIENDLY INSECT PEST MANAGEMENT'

Arthi Mohan ¹ and Hemapriya J ^{2*}

¹ Research Scholar, PG, Research Department of Microbiology, D.K.M. College for Women (Autonomous), Vellore, Tamil Nadu, India.

² Assistant Professor, PG, Research Department of Microbiology, D.K.M. College for Women (Autonomous), Vellore, Tamil Nadu, India. Affiliated to Thiruvalluvar University

*Corresponding author Email: microbiosep2021@gmail.com

ABSTRACT

As global agricultural production continues to expand, effective pest management has become essential for maintaining crop health and maximizing yields. Prioritizing food safety and environmental sustainability has increased the need for eco-friendly pest control approaches. Insects naturally harbor diverse microorganisms on their cuticle and within their bodies, which significantly influence their physiology, development, immunity, and behavior. The pest-associated microbiome includes bacteria, fungi, viruses, and protozoa, that can enhance the pest survival, influence the transmission of plant pathogens and provide protection against predators and environmental stresses. Understanding the structure and function of these microbial communities offers a sustainable and novel alternative to chemical pesticides. Strategies such as targeting symbiotic microbes, disrupting microbial-mediated detoxification pathways, or introducing pathogenic microorganisms can reduce pest fitness and population levels. Advances in high-throughput sequencing and metagenomics have enabled detailed characterization of pest microbiomes, revealing microbial taxa that are critical for survival and reproduction. However, variability in microbiome composition across pest populations and environmental conditions poses challenges for consistent application. Though this field of research has both promising opportunities and notable challenges. Advancing this field will deepen our understanding of insect-microbiome dynamics and pave the way for the development of innovative and sustainable strategies for pest management.

Keywords: Insect microbiome, pest management, symbiotic microbes, sustainable agriculture, microbiome-based biocontrol

**DIGITAL DECISION SUPPORT TOOLS AND REAL TIME ALERT
IN INTEGRATED PEST MANAGEMENT**

Rajaganapathy V ^{1*} and Raja Gopal V ²

¹ Bachelor of Science (Hons.) Agriculture, Kumaraguru Institute of Agricultural Science,
Tamil Nadu Agricultural University, Sakthinagar-638 315

² Department of Agronomy, Tamil Nadu Agricultural University, Coimbatore – 641 003

*Corresponding author Email: rajaganapathy0510@gmail.com

ABSTRACT

Digital decision support tools and real-time alert systems are reshaping the way Integrated Pest Management (IPM) is implemented across agricultural and processing environments. By combining data from remote sensors, IoT devices, digital traps, weather models, and pest surveillance networks, these tools enable timely and accurate decision-making. Real-time alerts allow stakeholders to detect pest outbreaks early, apply targeted interventions, and minimize crop losses while reducing reliance on broad-spectrum pesticides. Decision support platforms further enhance IPM by offering risk forecasting, treatment recommendations, and automated reporting based on field-specific conditions. This seminar will explore the role of digital technologies in strengthening early warning systems, optimizing pest control strategies, and supporting sustainable farm and post-harvest practices. It will also highlight practical applications, economic benefits, current challenges, and the future potential of data-driven IPM solutions in promoting precision agriculture and environmental safety.

Keywords: Digital Decision Support, Real-Time Alerts, Integrated Pest Management, IoT and Sensors, Predictive Monitoring, Precision Agriculture

COMPUTER VISION AND DEEP LEARNING FOR PEST COUNTING AND CLASSIFICATION – A REVIEW

Raja Gopal V ^{1*} and Rajaganapathy V ²

¹ Department of Agronomy, Tamil Nadu Agricultural University, Coimbatore – 641 003

² Bachelor of Science (Hons) Agriculture, Kumaraguru Institute of Agricultural Science,
Tamil Nadu Agricultural University, Sakthinagar-638 315

*Corresponding author Email: rajagopal2304@gmail.com

ABSTRACT

Recent advances in computer vision and deep learning have opened new frontiers in pest detection, counting, and taxonomic classification, offering unprecedented precision and scalability in Integrated Pest Management (IPM). Conventional monitoring methods are often to manual, labor-intensive, and prone to observational bias are being rapidly outpaced by automated imaging systems coupled with convolutional neural networks and advanced feature extraction algorithms. These technologies enable high-throughput identification of pest species and life stages from trap images, field surveillance, and storage environments with significantly improved accuracy. Deep learning architectures, such as Faster R-CNN, YOLO, and EfficientNet, facilitate real-time detection and enumeration of pests, even under variable backgrounds, occlusions, and lighting conditions. When integrated with edge devices and cloud-based platforms, these systems provide continuous data streams that support predictive modeling, threshold-based interventions, and population dynamics assessment. This seminar will examine the scientific underpinnings, current applications, validation metrics, and technological constraints of vision-based pest classification. It will also discuss dataset quality, annotation challenges, model transferability across species and regions, and the future trajectory of AI-driven surveillance in agriculture and food safety ecosystems.

Keywords: Computer Vision, Deep Learning, Pest Classification, Automated Counting, Convolutional Neural Networks, Precision IPM

**BOTANICAL BULLETS: METHANOL AND PETROLEUM ETHER EXTRACTS
OF *Anisomeles malabarica* IN THE BATTLE AGAINST TOBACCO LEAF
CATERPILLAR – ECO FRIENDLY PEST MANAGEMENT**

Sakthivel S ^{1*} and Kandibane M ²

^{1*} Department of Entomology, Faculty of Agriculture, Annamalai University,
Chidambaram, Tamil Nadu 608002, India.

² Department of Agricultural Entomology, Pandit Jawaharlal Nehru College of
Agriculture and Research Institute, Karaikal, Puducherry 609603, India

*Corresponding author Email: duraisakthivel999@gmail.com

ABSTRACT

The tobacco leaf caterpillar, *Spodoptera litura* (Fabricius) (Lepidoptera: Noctuidae), is a major polyphagous pest responsible for severe damage in agricultural crops, particularly tobacco, cotton, soybean, and vegetables. Synthetic insecticides, though effective, lead to pesticide resistance, environmental contamination, and non-target effects on beneficial organisms. The present study evaluates the bio-efficacy of methanolic and petroleum ether extracts of *Anisomeles malabarica* Linnaeus (Malabar Catmint), a medicinal and aromatic plant belonging to the family Lamiaceae, against *S. litura*. The study investigated the effects of methanol and petroleum ether extracts of *A. malabarica* leaf on insect development. The methanol extract showed significant effects, with larval mortality ranging from 0-66.67%, pupal malformation from 11.11-55.56%, pupal mortality from 0-55.56%, and adult malformation from 0-22.22%. Specifically, methanol elution 2 recorded the highest larval mortality (66.67%), while elution 3 showed the highest pupal malformation (55.56%). In contrast, the petroleum ether extract did not exhibit any larval mortality or malformation but showed pupal malformation (0-55.56%) and pupal mortality (11.11-55.56%), with adult malformation ranging from 0-33.33%. These findings suggest that both extracts have potential insecticidal properties, with varying effects on different stages of insect development. In addition to assessing mortality and malformation rates, we also measured pupal length and weight to further elucidate the developmental impacts of the extracts. Comprehensive qualitative and quantitative phytochemical analyses were conducted to characterize the bioactive compounds present in the extracts. Notably, methanol elution 2, which exhibited pronounced larvicidal activity and induced significant regurgitation, was subjected to histological examination to investigate the morphological alterations in the larvae. Furthermore, GC- MS analysis was performed on methanol elution 2 to identify the specific compound(s) responsible for the observed larval mortality and regurgitation, thereby providing insight into the underlying mechanisms of action.

Keywords: *Anisomeles malabarica*, Methanol extract, Petroleum ether extract, *Spodoptera litura*

ALKANNIN: A POTENTIAL BIOPESTICIDE FOR SUCKING PESTS

Vaishnavi Sri K T^{*1}, Asha S¹ and Nagajothi R²

¹ UG Scholar, SRM College of Agricultural Sciences,

SRM Institute of Technology, Chengalpattu

² Assistant Professor, SRM College of Agricultural Sciences,

SRM Institute of Technology, Chengalpattu

*Corresponding author Email: vk7413@srmist.edu.in

ABSTRACT

In the aim of producing higher yield of food crops to feed the growing population, there is a shift from natural farming to commercial farming. Problems of pest and diseases are solved using synthetic chemicals, which are non-biodegradable, toxic and leads to biomagnification. Thus, as an alternative to the synthetic pesticides, organic or biopesticide made from secondary metabolites of plants, can be incorporated to achieve sustainability in food supply. The project focuses on the use of Alkannin (a naphthoquinone) present in the roots of *Alkanna tinctoria* (Alkanet) of Boraginaceae family to treat the sucking pests. Alkannin is known for their anti-microbial, anti-viral and pesticidal activity. Alkannin based agro-chemicals are eco-friendly. It was extracted via Soxhlet apparatus with alcohol as solvent. The crude extract was then tested on pests. These compounds are known to produce reactive oxygen radicals, thereby inhibiting mitochondrial respiration. Alkannin can be potentially used as a bio-pesticide, substituting the synthetic chemicals without harming humans and environment.

Key words: Food security, Sustainability, *Alkanna tinctoria*, Sucking pests, Bio-pesticide

**BIOEFFICACY OF BOTANICALS AGAINST RUGOSE SPIRALLING WHITEFLY
IN GUAVA: AN ECOLOGICAL INNOVATION FOR SUSTAINABLE PEST
MANAGEMENT**

Navadeepchidambaram S ¹ and Kumar K ²

¹Ph.D Scholar, Department of Entomology, Faculty of Agriculture,
PAJANCOA and RI, Karaikal, INDIA.

²Professor, Department of Entomology, Faculty of Agriculture,
PAJANCOA and RI, Karaikal, INDIA.

*Corresponding author Email: navadeepsankar@gmail.com

ABSTRACT

The rugose spiralling whitefly (RSW), *Aleurodicus rugioperculatus*, has recently emerged as a major invasive pest causing severe damage to guava and other horticultural crops. In order to develop eco- friendly and sustainable strategies for its management, field investigations were conducted on guava variety Arka Rashmi to study the bioefficacy of botanicals, seasonal incidence, and their impact on natural enemies. The biology of RSW revealed extended developmental and reproductive stages that favor its rapid multiplication under field conditions. The management trials demonstrated that among the tested eco-friendly approaches, neem oil exhibited significant adult and nymphal suppression, offering a viable alternative to chemical insecticides. Other botanicals, though less effective, contributed to reducing pest pressure and supported the activity of natural enemies. Population dynamics of RSW indicated seasonal fluctuations closely associated with prevailing weather conditions, with higher incidence observed during specific months. Importantly, plots treated with botanicals harbored greater numbers of predatory spiders and coccinellids, highlighting their compatibility with natural enemy conservation. In contrast, chemical treatments resulted in marked reduction of beneficial arthropods, underlining the ecological advantage of botanicals in integrated pest management. The findings emphasize that integrating botanicals such as neem oil within pest management frameworks not only suppresses pest populations effectively but also sustains ecological balance through natural enemy preservation. This study underscores the role of botanicals as ecological innovations for sustainable pest management in guava cultivation, reducing reliance on synthetic insecticides and promoting environmentally resilient cropping systems.

Keywords: Rugose spiralling whitefly, guava, botanicals, neem oil, natural enemies, sustainable pest management

**ECO-FRIENDLY INTEGRATED PEST MANAGEMENT OF *Noorda moringae*
AND *Noorda blitealis* IN MORINGA (*Moringa oleifera* LAM.)**

Duvaraga Devi M^{1*} and Muthuswami M²

¹ Research Scholar, Department of Agrl. Entomology, TNAU, Coimbatore - 641003

² Professor, Department of Agrl. Entomology, TNAU, Coimbatore - 641003

*Corresponding author Email: duvaragamuthu@gmail.com

ABSTRACT

Moringa (Moringa oleifera Lam.) is a multipurpose crop widely cultivated in tropical regions, but its productivity is severely constrained by the budworm (*Noorda moringae* Tams) and the leaf-eating caterpillar (*Noorda blitealis* Walker). The present study aimed to evaluate the efficacy of biorationals and chemical insecticides against these pests and to develop an eco-friendly Integrated Pest Management (IPM) module. Field trials conducted at Puliampatti, Tiruppur district revealed that among biorationals, Azadirachtin 300 PPM was most effective against *N. moringae*, reducing bud infestation by 53.84% over the control, followed by *Bacillus thuringiensis* (49.50%). Among insecticides, Chlorantraniliprole 18.5 SC recorded the highest reduction (75.38%), followed by Emamectin benzoate 5 SG (70.18%), and Spinosad 45 SC (64.92%). Against *N. blitealis*, Azadirachtin 300 PPM again showed maximum efficacy (58.32%), followed by neem oil (55.75%) and *B. thuringiensis* (50.72%). Spinosad 45 SC was the most effective insecticide (84.69%), with Chlorantraniliprole 18.5 SC (82.43%) and Indoxacarb 15.8 EC (77.12%) also performing well. Based on the results, an IPM module was developed integrating soil raking, light traps, Azadirachtin 300 PPM, and need-based applications of Spinosad 45 SC and Chlorantraniliprole 18.5 SC. This strategy significantly outperformed farmers' practice, recording the lowest budworm infestation (15.52%) and leaf caterpillar load (4.25 larvae/branch). It also resulted in the maximum pod yield (9674 kg/ha) with a higher benefit-cost ratio (1:2.88), compared to 8021 kg/ha (1:2.27) under farmers' practice. The study underscores that integrating biorationals with reduced-risk, green-labelled insecticides provides a sustainable and economically viable approach to manage *N. moringae* and *N. blitealis* in moringa ecosystems.

Keywords: *Moringa oleifera*, *Noorda moringae*, *Noorda blitealis*, Integrated Pest Management (IPM), Biorationals

IPM AND THE FUTURE OF FOOD PRESERVATION: A SUSTAINABLE APPROACH

Chandraprabha S

Assistant Professor, Department of Post Harvest Technology, SRM College of Agricultural Sciences, Vendhar Nagar, Baburayanpettai, Chengalpattu Tamil Nadu, India.

Corresponding author Email: chandra.prabha625@gmail.com

ABSTRACT

Integrated Pest Management (IPM) represents a paradigm shift in agricultural sustainability, offering a holistic framework that minimizes pest-related losses while safeguarding environmental and human health. As global food systems grapple with post-harvest deterioration and mounting pressure to reduce chemical residues, IPM emerges as a pivotal strategy in enhancing the shelf life of food commodities. This abstract explores the synergistic relationship between IPM practices, including biological control, cultural interventions and judicious pesticide use and the preservation of food quality during storage and transit. By mitigating pest infestations at critical control points, IPM not only curtails spoilage but also maintains the nutritional integrity, aesthetic appeal, and marketability of produce. The integration of IPM into post-harvest protocols contributes to reduced reliance on synthetic preservatives, aligning with consumer demand for safer, residue-free food. Furthermore, IPM supports compliance with international export standards, thereby expanding market access for agricultural products. This paper underscores the role of IPM in shaping resilient food supply chains and calls for interdisciplinary collaboration to refine pest management strategies that extend shelf life without compromising ecological balance. Through case studies and emerging innovations, it advocates for IPM as a cornerstone of sustainable food preservation in the 21st century.

Keywords: Integrated Pest Management, shelf life, nutritional integrity, ecological balance and food preservation

NEXT-GEN IPM IN POST-HARVEST MANAGEMENT: BRIDGING SCIENCE AND TRADE

Prakash K

Assistant Professor and Head, Department of Post Harvest Technology, SRM College of Agricultural Sciences, Vendhar Nagar, Baburayanpettai, Chengalpattu, Tamil Nadu, India.
Corresponding author Email: hortidoctorpks@gmail.com

ABSTRACT

The post-harvest phase of agricultural production is increasingly vulnerable to pest infestations, which compromise food safety, reduce marketable yields, and hinder export potential. Integrated Pest Management (IPM), traditionally applied during crop cultivation, is now being reimagined for post-harvest systems to address these challenges with precision and sustainability. This abstract explores next-generation IPM strategies that integrate scientific innovation, regulatory compliance, and market intelligence to safeguard stored commodities and enhance global trade readiness. The emerging technologies, such as sensor-based pest detection, controlled atmosphere storage, and biopesticide applications, offer targeted, residue-free solutions that align with international phytosanitary standards. These interventions not only reduce reliance on chemical fumigants but also support traceability, certification and consumer trust. Commodity-specific IPM protocols, developed through interdisciplinary research, are enabling producers to meet the stringent quality benchmarks of high-value export markets. The paper further examines institutional frameworks, capacity-building models, and public-private partnerships that facilitate the adoption of IPM in post-harvest contexts. Emphasis is placed on harmonizing scientific rigour with trade facilitation, ensuring that pest management practices contribute to both ecological sustainability and economic competitiveness. By bridging science and trade, next-gen IPM transforms post-harvest management into a strategic lever for agricultural advancement, food security, and global market integration.

Keywords: Integrated Pest Management, sensor-based pest detection, controlled atmosphere storage, biopesticide application and food security

BACTERIAL ENDOPHYTES AS BIOCONTROL AGENTS: A SUSTAINABLE APPROACH TO INSECT PEST MANAGEMENT

Sneha R ^{*1} and Rex B ²

¹ PG Student – Department of Plant Pathology, SRM College of Agricultural Sciences SRMIST, Baburayanpettai, Chengalpattu District, Tamil Nadu.

² Assistant Professor -Department of Plant Pathology, SRM College of Agricultural Sciences SRMIST, Baburayanpettai, Chengalpattu District, Tamil Nadu.

*Corresponding author Email: sr7694@srmist.edu.in

ABSTRACT

Bacterial endophytes are microscopic prokaryotic organisms that maintain a symbiotic association with their host plants, contributing to growth promotion and defense against insect pests and pathogens. Extensive research has identified beneficial strains of bacterial endophytes capable of suppressing plant diseases and enhancing plant development through mechanisms such as induced systemic resistance (ISR). ISR is activated by bioactive compounds produced when a plant interacts with an endophyte, later resulting in systemic acquired resistance (SAR) upon insect or pathogen infestation. Notable examples include *Bacillus amyloliquefaciens*, producing lipopeptides to induce plant resistance against pests like the fall army worm, and *Enterobacter cloacae*, which acts as a biocontrol agent against rice pests through protein production. Transgenic endophytic bacteria, such as *Clavibacter xyli* subsp. *cynodontis* carrying the *cry1A* gene, exhibit insecticidal activity against lepidopteran pests. Studies on cabbage and related crops have shown that isolates of *Enterobacter cloacae*, *Alcaligenes piechaudii*, and *Khuyvera ascorbata* can effectively manage diamondback moth populations. These findings highlight the potential of bacterial endophytes as sustainable biocontrol agents in modern agriculture.

Keywords: bacterial endophytes, insect pest management, biocontrol, induced systemic resistance, *Bacillus amyloliquefaciens*, *Enterobacter cloacae*, *Clavibacter xyli*, sustainable agriculture

HARNESSING NATURE: ECOLOGICAL INNOVATIONS FOR SUSTAINABLE PEST MANAGEMENT

Shopiya K ^{*1} and Rageshwari S ²

¹ PG Scholar- Department of Plant Pathology, SRM College of Agricultural Sciences, SRMIST, Baburayanpettai, Chengalpattu District, Tamil Nadu.

² Assistant Professor and Head Department of Plant Protection, SRM College of Agricultural Sciences, SRMIST, Baburayanpettai, Chengalpattu District, Tamil Nadu.

*Corresponding author Email: rageshws@srmist.edu.in

ABSTRACT

A revolutionary move away from chemical-focused, short-term pest management techniques targeted towards integrated, sustainable methods based on ecological principles is represented by ecological innovations in pest management. With little consideration given to their interactions and compatibility, traditional pest control techniques frequently rely on single technologies, such as pesticides or biocontrol agents, with little chance of long-term success. A comprehensive grasp of agroecosystem dynamics, including local population biology and multitrophic interactions, is necessary for a truly effective pest management system, according to recent research. A key framework for reducing environmental impact and improving crop resilience is Integrated Pest Management (IPM), that combines targeted pesticide applications, biological control, genetic strategies, and cultural practices. Microbial biopesticides, enhanced delivery methods, and landscape-level interventions are examples of recent developments in integrated pest management. In addition, the Sterile Insect Technique (SIT) is an eco-friendly method that controls pest populations by releasing sterilized males, reducing reproduction and pest outbreaks while supporting sustainable, chemical-free pest management.

Keywords: Ecological pest management, Integrated Pest Management (IPM), Sustainable agriculture, biological control, Agroecosystem dynamics, Sterile Insect Technique (SIT).

RNAi-MEDIATED GENE SILENCING IN BPH – A REVIEW

Arul Jerlin J

B.Sc. (Hons.) Department of Agriculture, SRM College of Agricultural Sciences, Vendhar Nagar, Baburayanpettai, Chengalpattu district – 603 201.
Corresponding author Email: jj3903@srmist.edu.in

ABSTRACT

Brown Planthopper (*Nilaparvata lugens*) remains a critical pest in Asian countries, causing severe damage to crops, reducing overall production. With the available integrated pest management strategies and increasing resistance of BPH towards the method of control, it is crucial to include new technologies in controlling these pests. Recent studies show the role of the tramtrack(ttk) gene in male reproductive development and spermatogenesis of BPH. The research demonstrates that RNAi-mediated knockdown of ttk impairs male gonad development, reduces sperm count and motility, and drastically decreases fertility, indicating ttk as an effective molecular target for pest control. This work exemplifies the potential of RNAi as a species-specific, environmentally friendly approach aligned with integrated pest management (IPM) goals. This type of research paves the way for sustainable, precise, and reduced-chemical pest control in modern agriculture, contributing to improved crop protection and food security.

Keywords: Brown planthopper, RNAi, gene silencing, tramtrack (ttk), fertility.

COLD PLASMA TREATMENT FOR STORAGE PEST MANAGEMENT

Ambika S

Seed Science and Technology, SRM College of Agricultural Sciences, Chengalpattu

*Corresponding author Email: ambikasingaram@gmail.com

ABSTRACT

Insect pests in stored commodities cause major postharvest losses, posing risks to global food security. While chemical fumigants are traditionally used, their effectiveness is increasingly limited by resistance, toxic residues, environmental concerns, and health hazards. Cold plasma (CP), a non-thermal and eco-friendly technology, has emerged as a potential alternative, offering insecticidal, antimicrobial, and seed-preserving benefits. This review highlights CP's working principles, modes of action, practical uses, and limitations in storage pest management, emphasizing its promise as a sustainable replacement for chemical fumigants.

Key words: Storage pest, cold plasma treatment, seed storage

Introduction

Stored-product pests contribute to substantial losses, estimated at 10–30% of global grain stocks (Srivastava and Subramanian, 2016). Prominent species include *Sitophilus oryzae* (rice weevil), *Tribolium castaneum* (red flour beetle), and *Callosobruchus chinensis* (pulse beetle). Their activity reduces grain weight and yield, degrades nutritional quality, and lowers seed viability. Infestations also encourage fungal growth and mycotoxin production, compounding food safety risks (Eigenbrode et al., 2018). To date, multiple studies have investigated alternative methods to the use of chemical pesticides. These include natural plant-derived components (Deb et al., 2020; Singh and Kaur, 2018) entomopathogenic bacteria and nematodes (Ramos-Rodríguez et al., 2006; Malaikozhundan and Vinodhini, 2018; de Bortoli and Jurat-Fuentes, 2019) insect growth regulators (Wijayarathne et al., 2018) physical control methods, such as heat, cold, and modified atmosphere technology [Navarro, 2012; Fields and White, 2002; Agrafioti, 2019] ozone (Hardin et al., 2010) ionizing radiation (Hallman, 2013) and combined treatment approaches (Kostyukovsky et al., 2016). Traditional fumigants, such as methyl bromide and phosphine, are effective but face challenges like resistance, occupational hazards, and regulatory restrictions (Silva et al., 2019). These limitations create a need for innovative, non-chemical, and environmentally sustainable pest

management solutions. Cold plasma has emerged as a promising approach due to its broad-spectrum activity and minimal effect on grain quality. Fundamentals of Cold Plasma Technology Cold plasma is a partially ionized gas generated at or near room temperature using electrical discharges, including dielectric barrier discharge (DBD), plasma jets, and corona discharges (Bourke et al., 2018). It produces reactive oxygen and nitrogen species (ROS, RNS), free radicals, electrons, ions, and UV radiation. Unlike high-temperature plasmas, CP can safely treat heat-sensitive materials such as grains, seeds, and other stored foods. Its reactive species interact with insect exoskeletons, microbial cells, and seed surfaces, offering insecticidal, antifungal, and quality-preserving effects (Misra et al., 2019).

Mechanisms of Cold Plasma in Pest Control

- CP exerts pest control through multiple mechanisms:
- Cuticle damage: ROS and charged particles disrupt the insect cuticle and spiracles, hindering respiration (Laroussi and Leipold, 2004).
- □ Oxidative stress: Overproduction of ROS/RNS damages lipids, proteins, and DNA, leading to cellular dysfunction.
- Enzyme disruption: Antioxidant enzymes, including SOD, catalase, and GST, are impaired reducing insect resilience (Ziuzina et al., 2021).
- Reproductive inhibition: Plasma treatment decreases egg viability and larval growth, limiting population expansion (Pathan et al., 2021).
- Microbial suppression: Fungi and bacteria associated with pests are inactivated, indirectly reducing infestation levels.
- This multimodal approach contrasts with conventional insecticides, which usually target specific physiological pathways. (Ziuzina et al. (2021)

Applications in Stored Grain Protection

Cold plasma has been shown effective against key storage pests:

- Pulse beetle (*C. chinensis*): Plasma exposure lowers adult survival and reduces egg hatchability (Pathan et al., 2021).
- Rice weevil (*S. oryzae*): CP treatment decreases survival, reproduction, and larval development (Mohammadi et al., 2015).
- Red flour beetle (*T. castaneum*): Larvae and pupae are highly susceptible to plasma exposure (Abd El-Aziz et al., 2014).

Additional benefits include:

- Seed quality enhancement: Improved germination, vigor, and water uptake.

- Fungal and mycotoxin control: Inactivation of *Aspergillus* and *Fusarium* reduces aflatoxin and ochratoxin contamination (Khamsen et al., 2016).
- Preservation of nutritional and sensory properties: Appropriate plasma doses maintain grain quality.
- Integration potential: CP can be combined with hermetic storage, modified atmospheres, or biological control agents. Advantages and Limitations

Advantages:

- Chemical-free and environmentally sustainable.
- Broad-spectrum efficacy against insects, fungi, and microbes.
- Enhances seed germination and storage quality.
- Considered a “green technology” suitable for integrated pest management.

Limitations:

- Limited penetration in bulk grains; surface treatment is most effective.
- High initial cost and energy requirements.
- Standardized operational protocols are lacking.
- Regulatory approval for food applications remains limited.

Future Prospects

Wider adoption of CP requires:

- Development of scalable plasma devices for silos and warehouses.
Optimizing plasma flow for uniform insecticidal coverage.
- Energy-efficient systems, including solar or hybrid plasma units.
- Interdisciplinary collaboration among entomologists, food scientists, and plasma engineers. Establishment of regulatory standards for safe food applications.

Conclusion

Cold plasma is a promising, eco-friendly alternative for storage pest management. Its multifunctional effects-pest suppression, fungal/mycotoxin control, and seed-quality enhancement-make it a viable substitute for chemical fumigants. Though challenges remain in scaling, cost, and regulation, ongoing technological advancements may position CP as a mainstream component of integrated pest management, promoting sustainable storage and global food security.

**A REVIEW OF SOOTY MOLD ON MANGO (*Mangifera indica* L.):
ETIOLOGY, IMPACT, AND INTEGRATED MANAGEMENT**

Aruna G

Assistant Professor, Department of Plant Pathology,
SRM College of Agricultural Sciences, Chengalpattu – 603 201
Corresponding author Email: arunag@srmist.edu.in

ABSTRACT

Sooty mould disrupts mango output worldwide, especially in humid areas with many honeydew producing insects. Recent research on mango orchard sooty mould etiology, impact, and integrated management is summarised in this review. Sap sucking insects like mango hoppers (*Amritodus atkinsoni*), mealybugs (*Drosicha mangiferae*), and scales cause the problem. The sugary honeydew secreted by these pests feeds epiphytic fungal species as *Meliola mangiferae*, *Capnodium mangiferae*, and *Aspergillus spp.* The primary impact of sooty mold is indirect. The dense, black fungal mat on leaves and twigs blocks sunlight, reducing photosynthetic capacity and gas exchange. Even though the fruit pulp represents unaffected, the sooty coating decreases aesthetic appeal, market value, and export potential. Rainfall and insect pest population density significantly influence infestation severity. Insect infestations are best managed by targeting the root cause, not just the fungi. Reviewing control strategies:

Chemical Control: Application of systemic insecticides to control sap sucking pests.

Organic/Natural Control: Neem oil and insecticidal soaps destroy pests and remove mold. A spray of water and mild detergent can treat small infections.

Mechanical Removal: Physical removal of the sooty mold using a boiled starch solution, which dries and flakes off with the fungal growth.

Post-Harvest Treatments: Techniques like dipping fruits in specialized washes to remove blemishes and restore market quality.

Cultural Practices: Regular pruning increases airflow and sunlight, reducing mold formation.

Most sustainable and successful technique to reduce sooty mould ecological and economic impact on mango production is an integrated biological, cultural, and targeted chemical IPM strategy.

Keywords: Mango, sooty mold, *Meliola mangiferae*, honeydew, sucking pest, integrated pest management (IPM), photosynthesis, fruit quality.

SOCIOECONOMIC DETERMINANTS AND BEHAVIORAL DRIVERS OF DIGITAL IPM ADOPTION TOOLS AMONG FARMING COMMUNITIES

Rajasekaran R^{1*}, Jeevapriya A² and Indumathy K³

¹ Assistant Professor and Head, Department of Agricultural Extension and Communication, SRM College of Agricultural Sciences, Chengalpattu, Tamil Nadu

² Ph.D Scholar, Division of Dairy Extension, ICAR- National Dairy Research Institute: SRS, Bengaluru.

³ Associate Professor, Department of Agricultural Extension Adhiparasakthi Agricultural College, Tamil Nadu

*Corresponding author Email: rajasekr1@srmist.edu.in

ABSTRACT

The study examined the socioeconomic determinants and behavioral drivers influencing the adoption of digital Integrated Pest Management (IPM) tools among farmers in Chengalpattu district, using data from 120 respondents collected through interview schedules and focus group discussions. Descriptive statistics, correlation analysis, and logistic regression were employed to analyze awareness, usage, and adoption behavior. Results revealed that while 66.6% of farmers were aware of digital IPM tools, only 36.7% used them either regularly or occasionally. Key socioeconomic factors—education, digital literacy, farm size, and extension contact—showed strong positive correlations with adoption, while age was negatively associated. Behavioral factors such as perceived usefulness, trust in digital advisories, ease of use, and peer influence significantly shaped adoption decisions, with perceived usefulness having the highest predictive influence in the logistic regression model. The findings highlight that digital IPM adoption is driven by a combination of technological perception and social-economic readiness. Strengthening digital literacy, enhancing trust through locally relevant advisories, and integrating user-friendly, multilingual platforms were the essential suggested policy measures to bridge the gap between awareness and effective utilization of digital IPM tools in rural farming systems.

Keywords: Digital Integrated Pest Management (IPM), Technology Adoption, Socioeconomic Determinants, Behavioral Drivers

HOST PLANT SUSCEPTIBILITY INDEX OF OKRA GENOTYPES AGAINST LEAFHOPPER AND ITS ASSOCIATION WITH NUTRIENT CONTENT

Prithiva J N^{1*}, Ganapathy N², Jeyarani S², Premalatha K³, and Rajesh R⁴

¹ Research Associate, Department of Agricultural Entomology, TNAU, Coimbatore

³ Professor, Department of Agricultural Entomology, TNAU, Coimbatore

³ Associate Professor, Department of Agricultural Entomology, TNAU, Coimbatore

⁴ Research Associate, Department of Agricultural Microbiology, TNAU, Coimbatore

*Corresponding author Email: jnprithiva93@gmail.com

ABSTRACT

Two field studies were conducted at the Department of Agricultural Entomology, Tamil Nadu Agricultural University, Coimbatore, to assess the host plant susceptibility index of selected okra genotypes against the leafhopper, *Amrasca biguttula biguttula* Ishida, and to establish the relationship between leaf nutrient composition and pest incidence. Significant variation in leafhopper incidence was observed among the tested okra genotypes. The mean population of leafhopper was consistently higher in highly susceptible genotypes AE 26 and AE 15, followed by Pusa Sawani and AE 64, while the lowest infestation was recorded in AE 65 (moderately resistant) and Arka Anamika. Based on the pooled data of two field trials, the host plant susceptibility index (HPSI) indicated AE 26 as highly susceptible, AE 15 and Pusa Sawani as susceptible, AE 64 as moderately susceptible, while AE 65 and Arka Anamika were categorized as moderately resistant genotypes. Nutrient composition analysis revealed distinct differences among the genotypes. Nitrogen content was maximum in AE 26 (3.66%), followed by Pusa Sawani (3.48%) and AE 15 (3.32%). Phosphorus content was higher in AE 15 (0.18%) and AE 26 (0.16%), whereas potassium content was maximum in AE 65 (1.43%) and Arka Anamika (1.36%). Correlation studies showed that nitrogen ($r = 0.872$) and phosphorus ($r = 0.737$) were positively correlated with leafhopper population, while potassium ($r = -0.971$) exhibited a strong negative correlation. The results suggest that higher nitrogen and phosphorus levels in okra foliage may enhance the susceptibility to leafhopper attack, whereas increased potassium concentration contributes to tolerance or resistance. These findings emphasize the potential role of nutrient management and host plant resistance in integrated pest management strategies.

Key words: Okra, leaf hopper, HPSI, nutrient content, resistance

NATURE'S SHIELD FOR ECO-FRIENDLY SEED STORAGE PEST CONTROL

Vinothini Nedunchezhiyan

Discipline of Seed Science and Technology, SRM College of Agricultural Sciences,
SRM Institute of Science and Technology, Baburayanpettai, Chengalpattu District - 603 201,
Tamil Nadu, India.

*Corresponding author Email: ns.vinothini93@gmail.com

ABSTRACT

Sustainable pest management in seed storage systems is increasingly vital as the overuse of synthetic fumigants and insecticides poses serious risks to human health, seed viability, and ecological balance. This study focuses on Nature's Shield, an integrated framework employing eco-friendly and technologically advanced strategies for controlling storage pests while maintaining seed quality. Recent innovations such as nanotechnology-based biopesticides, biopolymer coatings infused with essential oils, and microbial antagonists like *Bacillus subtilis* and *Beauveria bassiana* demonstrate high efficacy in suppressing pest populations through biological and physical mechanisms. In parallel, smart sensor networks and IoT-based monitoring systems are transforming storage environments by enabling real-time tracking of temperature, humidity, and pest movement. Furthermore, the application of machine learning algorithms enhances predictive modeling for early pest detection, allowing for timely and precise interventions. Controlled atmosphere storage (CAS) and modified humidity management systems further contribute to the suppression of pest development while reducing chemical dependency. Collectively, these technologies form a synergistic, sustainable approach to seed preservation, promoting long-term storage stability and biosecurity. The integration of ecological principles with digital innovations presents a viable pathway toward minimizing post-harvest losses, enhancing seed longevity, and ensuring food system resilience. The findings emphasize that eco-technological convergence can provide scalable, non-toxic, and environmentally compatible alternatives to traditional pest management practices.

Keywords: Eco-friendly Pest Management; Seed Storage Protection; Nanobiotechnology; Smart Monitoring Systems; Biological Control

INFLUENCE OF WEATHER PARAMETERS ON NATURAL PARASITIZATION OF *Helicoverpa armigera* BY *Campoletis chloridaeae* UCHIDA IN CHICKPEA ECOSYSTEM OF THE NEW ALLUVIAL ZONE OF WEST BENGAL

Sathees Kumar K¹ and Meenambigai C²

¹ Department of Basic Sciences, SRM College of Agricultural Sciences, SRM Institute of Science and Technology, Baburayanpettai, Chengalpattu District, Tamil Nadu, India - 603 201.

² Department of Entomology, Bidhan Chandra Krishi Viswavidyalaya, Mohanpur, Nadia, West Bengal, India – 741 246.

*Corresponding author Email: satheesk@srmist.edu.in

ABSTRACT

The gram pod borer (*Helicoverpa armigera* Hübner) is a major pest of chickpea (*Cicer arietinum* L.), naturally regulated by the endo-larval parasitoid *Campoletis chloridaeae* Uchida. This study examined the influence of weather parameters on the parasitization rate of *C. chloridaeae* in chickpea ecosystems of the new alluvial zone of West Bengal. Field experiments conducted for three consecutive Rabi seasons (2017–18 to 2019–20) at Mondouri Farm, BCKV, followed a randomized complete block design. Second-instar larvae of *H. armigera* were reared until cocoon formation, and parasitization percentage was related to weather variables using correlation and stepwise regression analyses. The parasitoid appeared during the third standard meteorological week with a peak parasitization of 68–72% at flowering to pod initiation stages. Maximum and minimum temperatures showed significant negative correlations with parasitization, explaining about 50% of its variation. Rising temperatures may reduce *C. chloridaeae* efficiency, necessitating climate-resilient IPM strategies in chickpea.

Keywords: *Helicoverpa armigera*, *Campoletis chloridaeae*, Parasitization rate, Weather parameters, Integrated pest management.

BACTERIOPHAGES – A NOVEL BIOPRESERVATIVE FOR VEGETABLES

Ponvizhi Ramya V ^{*1} and Gunasekaran S ²

¹Department of Agricultural Microbiology, TNAU, Coimbatore

²Professor and Head, Department of Agricultural Microbiology, TNAU,
Coimbatore – 641 003.

*Corresponding author Email: v.ponvizhiramya@gmail.com

ABSTRACT

Survey was conducted on farmer's field for examination of soft rot incidence in vegetable growing regions of Tamil Nadu. Rotten potatoes and tomatoes with soft rot symptom were collected from farmer field and market. Screening of the isolates was performed based on rot zone, rot weight, rot severity and polygalacturonase activity. Based on the screening, isolate PP 4, PP 1, PP 3, PE 1, PE 6, PE 11, PE 19, TE 8, TE 1, TE 3, TE 9, TE 12, TE 13, TE 18 and TE 21 were selected as virulent pathogens by comparison with the standard culture *Pectobacterium carotovorum* subsp. *Carotovorum* ATCC 39048. Isolates that were considered to be soft rot causing species were characterized and identified on the basis of conventional phenotypic and biochemical characteristics. Screened isolates were identified using molecular characterization as PP 1 (*Cellulomicrobium funkei*), PP 2 (*Enterobacter asburiae*), PP 3 (*Kocuria palustris*), PP 4 (*Pectobacterium carotovorum* subsp. *carotovorum*), PE 6 (*Bacillus subtilis*), PE 11 (*Bacillus aryabhattai*), PE 1 (*Arthrobacter nicotianae*) and PE 19 (*Klebsiella oxytoca*) from rotten potato and TE 1 (*Lactobacillus pentosus*), TE 8 (*Citrobacter freundii*), TE 12 (*Enterobacter hormaechei*), TE 13 (*Klebsiella variicola*), TE 18 (*Enterobacter cloacae* subsp. *cloacae*), TE 21 (*Klebsiella pneumonia*), *Lactobacillus* from tomato. Isolates were tested for its various host range and found to have wide host range on Tomato, Onion, Okra, Carrot, Cabbage, Radish, Broccoli, Radish, Cauliflower, Ivy gourd, Capsicum, Brinjal, Potato, Tomato and Lettuce. In the host range infectivity analysis, brown discoloration and decaying symptoms identical to soft rot symptom were observed up to a week after inoculation, whereas negative control treatments remain symptomless in all vegetables tested. TE 8 (*Citrobacter freundii*) had a broad host range infecting all families of vegetables except Onion, Broccoli, Radish, Ivy gourd, Capsicum, Potato and Lettuce in secondary screening. TE 12 (*Enterobacter hormaechei*) and TE 21 (*Klebsiella pneumonia*) had a host range on onion. PP 4 (*Pectobacterium carotovorum* subsp. *carotovorum*) had a broad host range infecting all families of vegetables except Brinjal in secondary screening. PP-1 (*Cellulosimicrobium funkei*) had a host range on Capsicum, Potato and Lettuce. PP 2 (*Enterobacter asburiae*) and PP-3 (*Kocuria palustris*) were virulent on Capsicum and potato. PE 1 (*Bacillus subtilis* subsp. *inaquosorum*), PE-6 (*Bacillus aryabhattai*) and PE 11 (*Arthrobacter nicotianae*) infects onion and potato. PE 19 (*Klebsiella*

oxytoca) could spoil onion, potato and as well as tomato. Cellulase and pectinase enzyme activity was determined for the screened isolate using agar cup assay. The performance of standard culture *Pectobacterium carotovorum* subsp. *carotovorum* ATCC 39048 was compared for screening virulent isolates like PP 1, PP 2, PP 3, PP 4, PE 6, PE 11, PE 1, PE 19, TE 1, TE 3, TE 8, TE 9, TE 12 and TE 13. PP 4 (*Pectobacterium carotovorum* subsp. *carotovorum*) from potato and TE-8 (*Citrobacter freundii*) from tomato has higher pectinolytic activity and cellulolytic activity among those virulent isolates in secondary screening. Healthy, naturally infected and artificially screened pathogens inoculated potato and tomato on third day were evaluated for differential polyphenoloxidase activity, total phenolic compound, titratable acidity, pH, soluble solids content, reducing sugars, colour and firmness. Increase in reducing sugars, total soluble solids, polyphenoloxidase activity, total phenolic compound and decrease in firmness and colour in infected potato than the healthy exactly indicates the quality changes of potato by spoilage pathogens. Pathogenesis related proteins were up regulated and disease resistant inhibitor proteins were down regulated indicating respectively in both naturally infected and *P. carotovorum* subsp. *carotovorum* (PP 4) inoculated when compared with the healthy tissue. Totally four phages were isolated from rhizosphere soil from Coimbatore, Dindigul and Niligris against spoilage bacteria (PP-4) *Pectobacterium carotovorum* subsp. *carotovorum* and six phages for (TE 8) *Citrobacter freundii*. Phages were purified by streak method of purification with slight modification. Host ranges of phages were determined for its lytic activity. VegsavPhage 7 had a wide host range infecting major spoilage bacteria TE 1 of *Enterobacteriaceae* family (*Lactobacillus pentosus*), TE 8 (*Citrobacter freundii*), TE 12 (*Enterobacter hormaechei*), TE 13 (*Klebsiella variicola*), TE 18 (*Enterobacter cloacae* subsp. *cloacae*), TE 21 (*Klebsiella pneumonia*), PP 2 (*Enterobacter asburiae*) PP 3 (*Kocuria palustris*), PP 4 (*Pectobacterium carotovorum* subsp. *carotovorum*), PE 1 (*Bacillus subtilis* subsp. *inaquosorum*), PE 11 (*Arthrobacter nicotianae*) and PE-19 (*Klebsiella oxytoca*) followed by phage 5 having host range for TE 1 (*Lactobacillus pentosus*), TE 8 (*Citrobacter freundii*), TE 12 (*Enterobacter hormaechei*), TE 13 (*Klebsiella variicola*), TE 18 (*Enterobacter cloacae* subsp. *cloacae*), TE 21 (*Klebsiella pneumonia*), PP 2 (*Enterobacter asburiae*) PP 3 (*Kocuria palustris*), PP 4 (*Pectobacterium carotovorum* subsp. *carotovorum*), PE 1 (*Bacillus subtilis* subsp. *inaquosorum*) and PE 19 (*Klebsiella oxytoca*). Titer values of phage stocks in SM buffer were examined. Phage titre was around 15×10^{13} pfu/ml. PP 4 and TE 8 become resistant to all the phages at 48 hrs. 24 hrs was optimum for harvest of phages. Spoilage of potato and tomato was controlled by phage and had better biopreservative efficiency. Rot weight, rot severity, rot zone and population were reduced compared with

inoculated control. VegsavPhage 7 has broad host range, lesser resistant to Pectobacterium and exhibited better biopreservative efficiency whilst screening. Hence, growth cycle of Phage 7 was estimated up to one hour. Latent period was up to 30 minutes. Rise period was around 15 min. Burst size was 8×10^6 pfu/ml. Vegsavphage 7 was multiplied in one litre flask and they were filtered and stored. AFM has revealed the morphology of bacteriophage named vegsavphage 7, belongs to Myoviridae family (T 4). Repeated oral toxicological study was conducted on wister rats for bacteriophages (T 4). The results were much fruitful because they were safe to humans because no toxicity in rats. It could be better exploited as a biopreservative for vegetables by entrapping them in the microspheres of protein film coated over the vegetables. Coated films over vegetables are water soluble and could be washed before cooking. It prevents the spoilage due to pathogens and enhances the shelf life of vegetables up to 50 days. The study presented here has established the proof of concept, Bacteriophages - A novel biopreservative for vegetables against spoilage/opportunistic pathogens.

Keywords: Bacteria, Phages, Vegetables, Pathogens

BIOACOUSTIC SURVEILLANCE SYSTEMS FOR DETECTING INSECT FOR SUSTAINABLE AGRICULTURE

Deepika D^{*1}, Jeyajothi R² and Thamizharasu T¹

¹Research Scholar, Department of Agronomy, SRM College of Agricultural Sciences, Baburayanpettai - 603 203, Chengalpattu (Dt), Tamil Nadu.

² Assistant Professor, Department of Agronomy, SRM College of Agricultural Sciences, Baburayanpettai, Maduranthakam (Tk), Chengalpattu (Dt) - 603 201, Tamil Nadu.

*Corresponding author Email: dd7202@srmist.edu.in

ABSTRACT

Effective management of agricultural insects depends on the timely and accurate detection of pest populations to minimize crop damage and enhance control efficiency. Conventional monitoring techniques, including visual scouting and pheromone trapping, are often limited in sensitivity, particularly for cryptic or endophytic species such as stem borers (*Chilo partellus*), wood-boring beetles, termites, and subterranean root feeders that remain concealed within plant tissues or soil environments. Bioacoustic surveillance has emerged as an innovative, non-invasive approach that capitalizes on the distinct vibrational and airborne acoustic signals generated by pest activities such as feeding, locomotion, and stridulation. Highly sensitive transducers, including piezoelectric contact microphones and accelerometers, are deployed on plants, tree trunks, or within the soil to detect these pest-generated vibrations. Advanced signal processing and machine learning algorithms facilitate the discrimination of pest-specific acoustic patterns from ambient environmental noise caused by wind, rain, or non-target organisms. Integration of bioacoustic data with agronomic parameters such as crop phenology, soil moisture, and microclimatic variables enables spatially and temporally resolved assessments of pest populations. This supports precision intervention strategies within integrated pest management (IPM) frameworks, reducing unnecessary pesticide use and improving sustainability. Although challenges remain regarding sensor placement optimization, background noise management, and the establishment of comprehensive acoustic reference libraries, recent advancements in wireless sensor networks, edge computing, and artificial intelligence have substantially enhanced detection accuracy, real-time responsiveness, and system scalability. Consequently, bioacoustic surveillance represents a transformative technology for sustainable, data-driven pest management in modern agriculture.

Key words- Bioacoustic, Sensor, Surveillance, Vibration, Signal

**THE MICROBIAL SHIELD: HARNESSING PLANT-MICROBIOME
INTERACTIONS FOR PEST RESILIENCE**

Thamizharasu T^{*1}, Jeyajothi R ² and Deepika D¹

¹Research Scholar, Department of Agronomy, SRM College of Agricultural Sciences,
Baburayanpettai - 603 203, Chengalpattu (Dt), Tamil Nadu.

² Assistant Professor, Department of Agronomy, SRM College of Agricultural Sciences,
Baburayanpettai, Maduranthakam (Tk), Chengalpattu (Dt) - 603 201, Tamil Nadu.

*Corresponding author Email: tt7555@srmist.edu.in

ABSTRACT

The escalating demand for sustainable agriculture has intensified research into plant-microbiome interactions as an innovative alternative to conventional pest control methods. Plants coexist with complex and dynamic microbial communities collectively termed the plant microbiome that colonize the rhizosphere, phyllo sphere, and endosphere. These microbial assemblages are increasingly recognized as critical allies in enhancing plant health, particularly in bolstering resilience against pests. This concept of a microbial shield emphasizes the natural capacity of beneficial microbes to prime plant defences, compete with or antagonize pathogens and herbivorous insects, and modulate host signalling pathways involved in stress responses. This review explores the multifaceted mechanisms through which the plant microbiome contributes to pest resistance, including the induction of systemic resistance (ISR), production of volatile organic compounds (VOCs), and manipulation of insect behaviour via microbial metabolites. We examine recent advances in metagenomics, transcriptomics, and synthetic biology that enable deeper understanding and targeted manipulation of these interactions. Case studies from agricultural systems demonstrate how specific microbial taxa such as *Pseudomonas*, *Bacillus*, and mycorrhizal fungi can be harnessed as biocontrol agents to suppress pest populations, reduce reliance on chemical pesticides, and promote crop productivity. Furthermore, we discuss the challenges in translating laboratory findings into field applications, including microbiome stability, environmental variability, and host-specific responses. The integration of microbiome-based strategies into pest management frameworks represents a paradigm shift towards ecological intensification, where biological diversity is leveraged to enhance crop resilience. By decoding and engineering the plant microbiome, agriculture stands at the frontier of developing a sustainable microbial shield against pest pressures in the face of climate change and growing global food

demands.

Key words- Sustainable, Microbiome, Pest population, Resilience, Biocontrol agents.

IMPACT OF SPIDES ON SLAUGHTER CHARACTERISTICS OF BROILER CHICKENS FROM LONG-TERM STORED EGGS

Kowsalya N¹, Anandhi M², Jayanthi D², Shamsudeen P⁴ and Karthikeyan G⁵

¹M.Tech., Poultry Technology, Department of Poultry Technology, College of Poultry Production and Management, Hosur, Tamil Nadu, India

²Assistant Professor, Department of Poultry Technology, College of Poultry Production and Management, Hosur, Tamil Nadu, India

³Professor and Head, Department of Poultry Technology, College of Poultry Production and Management, Hosur, Tamil Nadu, India

⁴Professor and Head, Department of Poultry Management, College of Poultry Production and Management, Hosur, Tamil Nadu,

⁵Assistant Professor, Section of Animal Sciences, SRM University, Chengalpattu, Tamil Nadu

*Corresponding author Email: nkowsalya231299@gmail.com

ABSTRACT

This experiment assessed the effects of Short Periods of Incubation During Egg Storage (SPIDES), applied with or without egg turning, on the slaughter traits of broilers hatched from eggs stored for an extended period. A total of 750 hatching eggs obtained from 33.5-week-old breeder hens were randomly distributed into five treatment groups: T1 (control, no SPIDES), T2 (three SPIDES without turning), T3 (three SPIDES with turning), T4 (four SPIDES without turning), and T5 (four SPIDES with turning). Eggs were maintained at a temperature of 17 °C and 75% relative humidity, while those subjected to SPIDES were periodically heated to 37.7 °C for three hours at five-day intervals. All eggs were stored for 21 days and transferred for incubation on the 22nd day. Among the treatments, T5 exhibited the highest New York dressing yield (90.45%), dressed weight percentage (74.19%), ready to-cook weight (79.78%) and liver weight percentage (2.91%). No significant differences were observed in the small intestine, gizzard, heart, breast, neck, ribs and back, thigh and drumstick weight percentage among the groups. However, T2 (3 SPIDES without turning) had the highest wing weight percentage (12%). These findings suggest that SPIDES, particularly with turning, enhances carcass yield parameters in broilers.

Keywords: SPIDES, hatching eggs, broiler slaughter traits, carcass yield

**INFLUENCE OF DIETARY AZOLLA ON GROWTH AND PERFORMANCE
PARAMETERS IN COMMERCIAL BROILERS CHICKEN**

Kowsalya N¹, Sreyass K S² and Karthikeyan G²

¹N Kowsalya M.Tech Poultry Technology, Department of Poultry Technology, College of Poultry Production and Management, TANUVAS, Chennai, Tamil Nadu, India

²Sreyass KS Assistant Professor, Section of Animal Sciences, SRM University, Chengalpattu, Chennai, Tamil Nadu, India

*Corresponding author Email: nkowsalya231299@gmail.com

ABSTRACT

This study investigated the effect of dietary supplementation of dried Azolla at different inclusion levels on production performance parameters of commercial broiler chickens. A total of 250-day-old broiler chicks were randomly divided into five treatment groups: T1 (Control: 100% commercial feed), T2 (1% Azolla + 99% commercial feed), T3 (3% Azolla + 97% commercial feed), T4 (5% Azolla + 95% commercial feed), and T5 (7% Azolla + 93% commercial feed). All birds were reared under standard management practices for the duration of the experiment. Significant differences were observed among the treatment groups for body weight, feed consumption, feed conversion ratio, and livability. Among all treatments, broilers fed 5% Azolla with commercial feed (T4) demonstrated the best overall production performance. In conclusion, dietary inclusion of dried Azolla, particularly at 5%, can effectively enhance growth and production parameters in commercial broiler chickens.

Keywords: Azolla, body weight, feed consumption, feed conversion ratio, livability, broiler

DIGITAL TRANSFORMATION IN THE PEST MANAGEMENT OF FRUIT CROPS

Deepika V^{1*}, Abishek A² and Akshaya P²

¹Assistant Professor, SRM College of Agricultural Sciences, SRM IST,
Baburayanpettai, Chengalpattu

²PG Scholar, SRM College of Agricultural Sciences, SRM IST, Baburayanpettai, Chengalpattu

*Corresponding author Email: deepikav1@srmist.edu.in

ABSTRACT

The rapid evolution of digital technologies has revolutionized the terrain of Integrated Pest Management (IPM) in fruit crops. Traditional pest management approaches, often reliant on chemical control, faces many challenges due to pest resistance, environmental concerns and increased production costs. In contrast, the integration of digital tools such as artificial intelligence (AI), Internet of Things (IoT), remote sensing, drone technology and decision support systems (DSS) has enabled more precise, real time and sustainable pest management strategies. These innovations allow early detection, spatial monitoring and predictive modelling of pest outbreaks by enhancing the accuracy of interventions and minimizing the use of pesticides. The recent advancements in digital diagnostics, automated pest surveillance, machine learning algorithms for pest forecasting and mobile based advisory platforms are vital in orchard management. Case studies from tropical and subtropical fruit crops are highlighted the potentiality of these tools in improving pest control efficiency and environmental safety. Furthermore, it includes the challenges in technology adoption, data integration and capacity building among the growers. Hence, the digital transformation of IPM represents a paradigm shift towards smart, sustainable and climate resilient pest management in fruit crops production.

Keywords - Digital IPM, artificial intelligence, internet of things, pest forecasting, fruit crops

ZERO-RESIDUE INTEGRATED PEST MANAGEMENT (ZR-IPM)

Abinesh*¹ and Marimuthu S²

^{*1} M.Sc. Scholar, Department of Agronomy, SRM College of Agricultural Sciences, SRM Institute of Science and Technology, Vendhar Nagar, Baburayanpettai, Madhuranthakam (TK) Chengalpattu (Dt) - 603 201, Tamil Nadu.

² Assistant Professor (Senior Grade), Department of Agronomy, SRM College of Agricultural Sciences, SRM Institute of Science and Technology, Vendhar Nagar, Baburayanpettai, Madhuranthakam (TK) Chengalpattu (Dt) - 603 201, Tamil Nadu.

*Corresponding author Email: as1506@srmist.edu.in

ABSTRACT

The concept of *Zero-Residue Integrated Pest Management (ZR-IPM)* represents a transformative advancement in sustainable and organic agriculture, aiming to produce food entirely free from chemical residues while maintaining high crop productivity and ecological balance. Unlike conventional IPM, which primarily focuses on reducing pesticide use, ZR-IPM integrates *biodegradable bioinputs, microbial consortia, botanical formulations, and nano-biocarriers* to ensure complete degradation of active ingredients before harvest. This approach relies on a 4D defense framework — Detection, Decision, Defense, and Degradation — combining real-time pest surveillance through sensors and AI-based predictive tools with threshold-guided application of biopesticides, beneficial microbes, and plant-derived volatiles. Recent innovations such as chitosan-encapsulated neem nanoemulsions, microbial blends of *Trichoderma*, *Beauveria*, and *Bacillus*, and pheromone-based mating disruption systems have demonstrated over 90% pest suppression with undetectable residue levels (<0.01 ppm) in tomato, okra, and cotton trials across India. These eco-safe formulations not only manage pest complexes effectively but also enhance soil microbial activity, crop vigor, and pollinator safety. Furthermore, ZR-IPM aligns with global sustainability goals, addressing consumer demand for residue-free food and meeting stringent export regulations under the EU's "Farm to Fork" strategy and India's "Zero Pesticide Residue" initiatives. By merging organic principles, biotechnology, and digital intelligence, Zero-Residue IPM redefines pest management as an ecologically regenerative and economically viable system. It envisions a future where protecting crops no longer compromises environmental purity or human health — achieving the ideal of "*safe food from a safe ecosystem.*"

Keywords: Zero-residue IPM; Biodegradable biopesticides; Nano-biocarriers; Microbial consortia; Botanical formulations; Organic pest management; Sustainable agriculture; Residue-free food; AI-guided pest monitoring; Eco-safe farming.

ADVANCING INTEGRATED PEST MANAGEMENT THROUGH AI AND MACHINE LEARNING INNOVATIONS

Sathya Devi S

PG Student, Department of Entomology, SRM College of Agricultural Sciences, SRMIST, Baburayanpettai, Chengalpattu, Tamil Nadu – 603 201.

*Corresponding author Email: sathiyadeviaks317@gmail.com

ABSTRACT

The emergence of digital technologies has opened new frontiers in Integrated Pest Management (IPM), offering intelligent, data-driven solutions for sustainable crop protection. This chapter explores how innovations in artificial intelligence (AI), machine learning (ML), and precision agriculture are revolutionizing pest monitoring, forecasting, and control. As agriculture faces growing challenges such as climate variability, pest resistance, and the need to reduce chemical dependence, the integration of these technologies has become essential for ensuring food security and environmental sustainability. Cutting-edge digital tools and algorithms now enable predictive analytics, allowing farmers to anticipate pest outbreaks, assess crop health in real time, and implement timely interventions. Through remote sensing, Internet of Things (IoT) devices, and AI-powered image recognition, farmers can detect pest infestations early and optimize pesticide application with remarkable accuracy. These advancements not only improve pest control efficiency but also promote eco-friendly practices by minimizing input use and environmental impact. The chapter further examines applications such as AI-based pest identification, ML-driven predictive modeling for pest population dynamics, and decision-support systems that assist farmers in choosing effective and targeted IPM strategies. It also highlights the challenges associated with adopting digital IPM—such as data management, technological accessibility, and the need for farmer capacity-building—and emphasizes the importance of ethical and inclusive technology deployment. By exploring these digital frontiers, this chapter provides comprehensive insights into how technology is reshaping IPM into a more precise, proactive, and sustainable system. It concludes by discussing the future potential of integrating advanced digital platforms and collaborative innovation to achieve smarter pest management and resilient agricultural ecosystems.

Key words: Artificial Intelligence (AI), Machine Learning (ML), Digital Technologies and pest detection

HARNESSING THE INSECT MICROBIOME: A NEW FRONTIER IN SUSTAINABLE PEST MANAGEMENT

Swetha K^{*1}, Sweeta Annie Mary A¹ and Balasubramanian P²

¹M.Sc. Scholar, Department of Agronomy, SRM College of Agricultural Sciences, SRM Institute of Science and Technology, Vendhar Nagar, Baburayanpettai, Madhuranthakam (TK) Chengalpattu (Dt) - 603 201, Tamil Nadu.

²Assistant Professor, Department of Agronomy, SRM College of Agricultural Sciences, SRM Institute of Science and Technology, Vendhar Nagar, Baburayanpettai, Madhuranthakam (TK) Chengalpattu (Dt) - 603 201, Tamil Nadu.

*Corresponding author Email: sk6889@srmist.edu.in

ABSTRACT

The insect microbiome comprising diverse bacteria, fungi, and symbiotic microorganisms residing in the gut, cuticle, and reproductive organs plays a crucial role in insect physiology, nutrition, and defense. Recent research has revealed that these microbial communities profoundly influence pest behaviour, reproduction, immunity, and resistance to insecticides. Understanding these complex host-microbe interactions opens new ecological avenues for sustainable pest management. Manipulating the insect microbiome through Para transgenesis, probiotic applications, or microbial disruption can alter pest fitness, reduce vector competence, and suppress pest populations without harming non-target organisms. Beneficial microbes can also be engineered to deliver toxins or interfere with pathogen transmission, providing eco-friendly alternatives to chemical pesticides. Integrating microbiome-based strategies with existing Integrated Pest Management (IPM) frameworks enhances precision, sustainability, and long-term pest control efficacy. Although still in its developmental phase, microbiome manipulation represents a frontier innovation that aligns with global goals of environmentally safe and biologically driven pest management.

Keywords: Insect microbiome, paratransgenesis, symbionts, microbial manipulation, eco-friendly pest control, Integrated Pest Management.

CAPSAICIN: A PUNGENT MOLECULE WITH POTENT INSECTICIDAL POTENTIAL

Ilakiya T

Department of Horticulture, S.R.M. College of Agricultural Sciences, Chengalpattu, Tamil Nadu
Corresponding author Email: ilakiyatamil@gmail.com

ABSTRACT

Capsaicin, the primary pungent alkaloid of chilli peppers (*Capsicum annuum* L.), is becoming recognised as a powerful botanical pesticide with diverse bioactivity. Capsaicin, generated by the plant's innate defence mechanism, functions as an antifeedant, repellent, and neurotoxic agent against a diverse array of insect pests. The principal mechanism of action entails disrupting the insect nervous system by interacting with transient receptor potential (TRP) ion channels and interfering with neurotransmitter signalling, leading to paralysis, decreased feeding, and ultimately death. Moreover, capsaicin influences the hormonal equilibrium and digestive enzyme function in target insects, thereby inhibiting growth and reproduction. Experimental studies have confirmed its efficacy against aphids, whiteflies, thrips, and pests of stored grain, exhibiting low harm to beneficial organisms and the environment. In addition to its insecticidal capabilities, capsaicin demonstrates antioxidant and antibacterial effects, hence enhancing plant health and promoting resistance. Its biodegradable and non-persistent nature presents less risk of bioaccumulation or pesticide resistance, rendering it appropriate for integrated pest management (IPM) and organic farming systems. Modern research emphasises the enhancement of extraction and formulation methodologies, including encapsulation and nanoemulsion techniques, to improve stability, residual efficacy, and field performance. Capsaicin serves as a promising environmentally friendly alternative to synthetic pesticides, in accordance with the global desire for sustainable and safe agricultural protection solutions. Ongoing exploration of its molecular mechanisms and extensive application potential will accelerate its advancement as a dependable natural insecticidal agent.

Key Words: Capsaicin, *Capsicum annuum*, botanical insecticide, pest management, eco-friendly agriculture.

SUN-POWERED GUARDIANS: REDUCING PESTICIDES, SAVING BENEFICIAL INSECTS

Sanchana Dhas A K^{1*}, Ramadass S² and Priyanjena Chacko J¹

¹I PG Scholar, Dept. of Agronomy, SRMCAS, SRMIST, Chengalpattu-603201.

²Assistant Professor and Head, Dept. of Agronomy, SRMCAS, SRMIST, Chengalpattu-603201.

*Corresponding author Email: sa0135@srmist.edu.in

ABSTRACT

Heavy dependence on chemical pesticides in modern agriculture has raised serious concerns, including the development of pest resistance, environmental contamination, and the decline of beneficial insect populations. In response, automated solar-powered light traps offer an environmentally friendly and sustainable alternative for pest management by using renewable solar energy to attract and monitor pests. These traps emit specific light wavelengths to lure nocturnal insects such as *Helicoverpa armigera*, *Spodoptera litura*, and *Aphis craccivora*, which are subsequently captured using mechanical collection methods or trays. Utilizing solar energy allows for continuous, cost-effective operation while reducing reliance on conventional electricity. Field trials across various agro-ecosystems have demonstrated that solar light traps can significantly reduce pest densities and limit crop damage. Farmers implementing these devices have reported a 30–50% decrease in chemical pesticide usage, lowering production costs and preventing harmful residues in soil, water, and harvested crops. Additionally, minimizing pesticide applications supports the survival and recovery of beneficial insects, including ladybird beetles, lacewings, parasitoid wasps, and pollinators, which are essential for maintaining ecological balance. When integrated into Integrated Pest Management (IPM) strategies, solar light traps enable more effective pest control with minimal environmental impact. Overall, the use of automated solar light traps fosters a sustainable pest management approach, promoting biodiversity, conserving natural predators, and supporting organic and climate-resilient agricultural practices. This renewable energy-based technology not only enhances crop productivity and reduces ecological risks but also represents a significant step toward more eco-friendly and sustainable farming systems.

Keywords: Solar light trap, Integrated Pest management, Pesticide reduction, Beneficial insects, Biodiversity, Sustainable agriculture.

HARNESSING NATURE: ECOLOGICAL INNOVATIONS FOR SUSTAINABLE PEST MANAGEMENT

Yuvaraj N^{*1} and Akshaya S B²

¹PG Scholar- Department of Plant Pathology, SRM College of Agricultural Sciences, SRMIST, Baburayanpettai, Chengalpattu District, Tamil Nadu.

²Assistant Professor Department of Plant Pathology, SRM College of Agricultural Sciences, SRMIST, Baburayanpettai, Chengalpattu District, Tamil Nadu.

*Corresponding author Email: yn1432@srmist.edu.in

ABSTRACT

The shift toward sustainable agriculture is driving a wave of ecological innovations aimed at reducing reliance on synthetic pesticides while maintaining crop productivity. These nature based approaches embrace ecological balance and biodiversity, using biological control mechanisms to manage pests in a more harmonious and lasting way. Beneficial organisms such as entomopathogenic fungi, bacteria, parasitoids, and predators are increasingly used to suppress pest outbreaks without harming non-target species or disrupting ecosystems. Complementary tools like bioinoculants, pheromone traps, and botanical extracts further support natural pest suppression while enhancing soil and plant health. Emerging technologies, including nanobiotechnology, precision agriculture, and microbial bioformulations, are improving the stability, delivery, and field performance of these biological agents. When integrated into a broader framework like Integrated Pest Management (IPM), these eco-friendly solutions create a pest control system that is economically viable, environmentally responsible, and socially inclusive. Crucially, farmer engagement, awareness initiatives, and supportive policy environments are essential for scaling these practices globally. By combining traditional ecological wisdom with modern scientific advances, agriculture can build long-term resilience against pests while preserving vital ecosystem services. Ultimately, working with nature rather than against it offers a promising path toward climate-smart, productive, and sustainable farming systems ensuring both food security and environmental health for future generations.

Keywords: Sustainable Pest Management, Ecological innovations, Biological control, Bioinoculants, Biopesticides, Nanobiotechnology, Biodiversity, Integrated Pest Management (IPM), Regenerative agriculture, Environmental sustainability.

**ROLE OF METHYL EUGENOL-BASED PHEROMONE TRAPS IN SUSTAINABLE
CONTROL OF MANGO FRUIT FLY**

Mathiazhagan V

Assistant, Department of Entomology, SRM College of Agricultural Sciences
Corresponding author Email: mathiazv@srmist.edu.in

ABSTRACT

A serious pest of mangoes, the mango fruit fly (*Bactrocera dorsalis*) causes large output losses and presents serious difficulties for export markets because of stringent quarantine regulations. The purpose of this research is to determine how well pheromone trapping, a crucial part of Integrated Pest Management (IPM), works to suppress *B. dorsalis* populations in mango orchards. Traps based on methyl eugenol were set up at varying densities and observed throughout the fruiting season in order to assess their effects on adult male populations and the degrees of fruit infection that followed. The results showed that, in comparison to untreated control plots, the frequent deployment of pheromone traps considerably decreased fruit fly infestation and captures. In comparison to chemical pesticides, the trapping technique turned out to be sustainable, economical, and environmentally beneficial. These results indicate that pheromone trapping may be a keystone for the sustainable control of mango fruit fly populations when carefully combined with other IPM elements.

Keywords: Sustainable pest management, Methyl eugenol trap, Fruit fly.

INNOVATIVE APPROACHES FOR INTEGRATED PEST MANAGEMENT - A REVIEW

Nisha R

Assistant Professor, Department of Entomology, SRM College of Agricultural Sciences, SRMIST, Baburayanpettai, Chengalpattu District – 603 201.

Corresponding author Email: nisharengadoss@gmail.com

ABSTRACT

Integrated Pest Management (IPM) is a sustainable, ecosystem-based strategy that integrates biological, cultural, physical, and chemical tools to manage pest populations below economic injury levels while minimizing environmental harm. In recent years, innovative scientific and technological interventions have transformed IPM into a precision-oriented, climate-resilient, and environmentally compatible system. Modern innovations such as molecular and biotechnological tools, smart digital technologies, nano-formulations, and ecological engineering approaches have enhanced pest monitoring, forecasting, and management efficiency. This review highlights major advancements in IPM practices, emphasizing the integration of eco-friendly biocontrol agents, AI-based surveillance, and farmer-centric participatory models that promote sustainable crop protection and biodiversity conservation.

Keywords: Integrated Pest Management, Biocontrol, Biotechnology, Smart Farming, Biopesticides, Precision Agriculture, Climate-Smart IPM

Introduction

Integrated Pest Management (IPM) combines ecological principles and multidisciplinary approaches to suppress pest populations sustainably. Conventional pesticide-based pest control has led to resistance, resurgence, and environmental contamination. Consequently, IPM strategies have evolved to integrate technological innovations that enhance efficiency and sustainability (Kogan, 1998; Dent, 2000). Modern IPM incorporates smart tools, molecular diagnostics, and eco-engineering techniques for early pest detection and targeted management. These innovations strengthen the adaptability of IPM under changing climatic conditions and dynamic pest scenarios.

Eco-Friendly Biological Control

Biological control forms the foundation of IPM by utilizing natural enemies to maintain pest populations at sub-economic levels. The use of parasitoids such as *Encarsia guadeloupae* against *Aleurodicus dispersus* and *Trichogramma* spp. against lepidopteran borers has been well documented (Nagarajan et al., 2023). Advances in mass rearing, habitat manipulation, and conservation biological control have improved the effectiveness of natural enemies. Microbial control agents, including *Beauveria bassiana*, *Metarhizium anisopliae*, and *Bacillus thuringiensis*, are increasingly used in organic and conventional systems. Floral resource provisioning and intercropping with nectar-rich plants enhance the survival and activity of beneficial insects.

Molecular and Biotechnological Tools

Recent breakthroughs in biotechnology have revolutionized pest management. RNA interference (RNAi) offers species-specific pest suppression by silencing essential genes (Baum et al., 2007). Transgenic crops expressing insecticidal proteins from *Bacillus thuringiensis* reduce pesticide use and protect non-target organisms. DNA barcoding and molecular markers enable rapid pest and natural enemy identification, facilitating early warning and management decisions. These technologies enhance precision and reliability in IPM implementation.

Smart and Digital IPM

The integration of digital technologies into IPM enhances pest detection and decision-making. Internet of Things (IoT)-based traps and sensors provide real-time pest monitoring, while drones and unmanned aerial vehicles (UAVs) are employed for crop surveillance and precision pesticide application (Gupta et al., 2021). Artificial intelligence (AI)-driven mobile applications assist farmers in pest diagnosis and control recommendations. Decision Support Systems (DSS) use predictive algorithms and weather-based data to forecast pest outbreaks, promoting timely interventions.

Botanicals and Biopesticides

Plant-derived pesticides and microbial formulations are gaining importance as eco-friendly

alternatives to synthetic chemicals. Neem (*Azadirachta indica*), pongamia, and garlic extracts exhibit insecticidal and repellent activities. Nano-formulations enhance stability, bioavailability, and controlled release of active ingredients, improving pest control efficiency (Kumar et al., 2020). Synergistic use of botanicals and microbial biopesticides can improve efficacy while reducing pesticide residues in food and the environment.

Cultural and Agro-Ecological Strategies

Cultural and habitat manipulation techniques form the preventive component of IPM. Push-pull systems integrate repellent intercrops (push) and attractive trap crops (pull) to divert pests (Khan et al., 2016). Crop rotation, mixed cropping, and field sanitation disrupt pest life cycles. Ecological engineering through flowering borders and soil health management enhances the resilience of agroecosystems by favoring beneficial organisms.

Climate-Smart Pest Management

Climate change significantly alters pest population dynamics and distribution. Climate-smart IPM integrates pest forecasting models with real-time weather data to predict outbreaks. Remote sensing and Geographic Information Systems (GIS) enable large-scale pest surveillance and spatial risk assessment (Prasad et al., 2022). Such predictive tools aid in proactive pest control and minimize economic losses.

Community-Based and Policy Innovations

Farmer Field Schools (FFS) and participatory learning approaches empower farmers to make informed pest management decisions. Area-wide IPM programs coordinate pest suppression across landscapes, enhancing effectiveness (Vreysen et al., 2007). Government initiatives promoting biopesticide registration, residue-free produce certification, and integrated extension systems are essential for large-scale adoption of IPM innovations.

Conclusion

Innovative approaches in Integrated Pest Management combine ecological, technological, and participatory methods to ensure sustainable crop protection. The integration of biocontrol

agents, biotechnology, digital monitoring, and climate-smart forecasting has significantly strengthened IPM frameworks. Continued research, farmer capacity building, and supportive policies are essential to mainstream these innovations, ensuring reduced pesticide dependency and enhanced agroecosystem resilience.

References

- Baum, J. A., et al. (2007). Control of coleopteran insect pests through RNA interference. *Nature Biotechnology*, 25(11), 1322–1326.
- Dent, D. (2000). *Insect Pest Management*. CABI Publishing, UK.
- Gupta, R., Sharma, S., & Singh, A. (2021). Smart farming technologies for pest monitoring and control. *Journal of Agricultural Technology*, 17(2), 45–56.
- Khan, Z. R., Midega, C. A., Pittchar, J. O., Murage, A. W., Birkett, M. A., Bruce, T. J., & Pickett, J. A. (2016). Achieving food security for one million sub-Saharan African poor through push–pull innovation. *Proceedings of the National Academy of Sciences*, 113(17), 4068–4073.
- Kogan, M. (1998). Integrated pest management: Historical perspectives and contemporary developments. *Annual Review of Entomology*, 43, 243–270.
- Kumar, S., Rani, R., & Sinha, P. (2020). Nano-biopesticides: Emerging trends in pest management. *Frontiers in Sustainable Food Systems*, 4, 21–29.
- Nagarajan, M., Loganathan, R., & Nisha, R. (2023). Biological control potential of *Encarsia guadeloupae* against *Aleurodicus dispersus*. *Journal of Applied Zoological Research*, 34(2), 112–118.
- Prasad, R. A., Nisha, R., & Kirubakaran, G. (2022). Climate-smart pest management using GIS-based forecasting models. *Indian Journal of Entomology*, 84(4), 812–820.
- Vreysen, M. J. B., Robinson, A. S., & Hendrichs, J. (2007). *Area-wide Control of Insect Pests: From Research to Field Implementation*. Springer, Dordrecht.

INTEGRATED APPROACHES FOR THE CONTROL OF INSECT IN SPICES AND CEREALS

Saran M S¹, Ashmitha K¹, Vinitha E¹ and Sheela P^{2*}

¹UG Scholar, SRM College of Agricultural Sciences, SRM Institute of Science and Technology, Baburayanpettai, Chengalpattu - 603201

²Assistant Professor, Department of Post Harvest Technology, SRM College of Agricultural Sciences, SRM Institute of Science and Technology, Baburayanpettai, Chengalpattu - 603201

*Corresponding author Email: sheelap@srmist.edu.in

ABSTRACT

Spices and cereals are essential components of global diets, yet they are highly prone to insect infestation during post-harvest handling, storage and transport. Such infestations result in substantial quantitative and qualitative losses, adversely affecting aroma, colour, nutrient content and overall consumer acceptability. The management of storage insects therefore remains a critical challenge for ensuring food safety and market quality. This paper highlights the recent advances and integrated strategies employed to control insect pests in these commodities. Preventive measures such as improved sanitation, temperature and humidity regulation and the use of hermetic or modified-atmosphere packaging have proven effective in reducing infestation levels. Physical control techniques including heat and cold treatments, irradiation and controlled atmospheres provide residue free alternatives to conventional fumigants. Biological approaches utilizing parasitoids, entomopathogenic fungi and botanical insecticides offer eco-friendly options with minimal environmental impact. Chemical control methods, particularly phosphine fumigation, remain widely used but require strict regulatory oversight to prevent resistance development and residue accumulation. The integration of nanotechnology based formulations, sensor assisted pest monitoring and predictive modeling further enhances the efficiency of pest management systems. The adoption of an Integrated Pest Management (IPM) framework combining these approaches ensures the sustainable protection of spices and cereals while meeting the demands of food safety and environmental stewardship.

Keywords: Spices, Cereals, Insect infestation, integrated pest management, Post-harvest protection, Bio control, Storage technology.

**ASSOCIATION OF ENDOSYMBIONTS WITH *Bemisia tabaci* OF
PHYLOGENETICALLY DIFFERENT GROUPS**

Rageshwari S^{1*}, Buvaneswari S², and Deivamani M³

^{1*} Assistant Professor, Department of Plant Protection, ²SRM College of Agricultural Sciences,
SRMIST, Chengalpattu, Tamil Nadu

²Tamil Nadu Agricultural University, Coimbatore, Tamil Nadu, India.

³Assistant Professor, ICAR-KVK, Papparpatti, Tamil Nadu Agricultural University,
Dharmapuri, Tamil Nadu, India

*Corresponding author Email: rageshws@srmist.edu.in

ABSTRACT

Begomovirus stanleyi (Sri Lankan cassava mosaic virus) is considered as one of the important biotic constrain of the industrial crop cassava (*Manihot esculenta* Crantz). Cassava mosaic disease causes both quantitative and qualitative loss of up to 80 per cent. The virus is transmitted by the insect vector *Bemisia tabaci* (Gennadius). *B. tabaci* (Gennadius) (Hemiptera: Aleyrodidae) is an economic pest of important agricultural crops found worldwide. As a vector, it plays a key role in transmission of most of the plant viruses. Different populations of whitefly and its relationship with endosymbionts were studied based on mtCOI (Mitochondrial cytochrome oxidase), ITS region and RAPD analysis. Two different genotypes such as Q genotype and B genotype were identified from different crops cultivated in Tamil Nadu. The association of endosymbionts such as *Rickettsia* sp, *Arsenophonus* sp and *Wolbachia* sp with the whitefly biotypes were also studied. The endosymbionts were found to play major role in increasing the survival ability and virulence of the vector. Transmission efficiency of *Bemisia tabaci* was also studied and the inoculation feeding period for the SLCMV transmission was recorded to be 24 hrs (one day).

Keywords: *Bemisia tabaci*, genotype of whitefly, endosymbionts, transmission efficiency.

STUDIES ON NUTRITIONAL OPTIMIZATION FOR MYCELIAL GROWTH AND BIOMASS ENHANCEMENT OF ENTOMOPATHOGENIC FUNGI *Ophiocordyceps sinensis*

Akshaya S B

Assistant Professor, Department of Plant Pathology, SRM College of Agricultural Sciences,
SRM Institute of Science and Technology, Baburayanpettai, Chengalpattu 603- 201

Corresponding author Email: akshayas2@srmist.edu.in

ABSTRACT

Ophiocordyceps sinensis (formerly *Cordyceps sinensis*), an entomopathogenic fungus commonly known as the Chinese caterpillar fungus, is highly valued in traditional Chinese medicine due to its production of bioactive metabolites with significant pharmaceutical importance. However, its cultivation requires a variety of nutritional components to achieve rapid growth, high biomass yield, and successful fruiting body formation. In this context, a study was conducted to evaluate the effects of different carbon, nitrogen, mineral, vitamin, and chitin sources on the growth and biomass production of *O. sinensis* under liquid fermentation conditions. Among the tested nutrients, sucrose (3%) and beef extract (0.5%) supported the highest mycelial growth and biomass accumulation (90.00 mm colony diameter and 8.13–8.17 g dry weight/L after 18 days). The best-performing mineral sources were K₂HPO₄ and zinc chloride (0.2%), which promoted comparable growth and biomass yield (90.00 mm and 8.12–8.20 g/L on the 18th day). Furthermore, supplementation with folic acid (0.01%) and dried rhinoceros grub powder (0.1%) significantly enhanced mycelial development, resulting in the highest biomass production (90.00 mm and 8.31–8.69 g/L after 18 days). This study emphasises the significance of nutritional optimisation in enhancing mycelial growth, biomass accumulation, metabolite secretion, and potential fruiting induction in *O. sinensis* under controlled fermentation conditions.

Keywords: Entomopathogenic fungi, Biomass, *Ophiocordyceps sinensis*, mycelial growth and fermentation

INFLUENCE OF LIGHT SOURCES ON SPORULATION OF *Colletotrichum capsici*

Karpagavalli S

Associate Professor, Department of Plant Pathology, SRM College of Agricultural Sciences,
Vendhar Nagar, Baburayanpettai, Chengalpattu District – 603 201, Tamil Nadu, India

*Corresponding author Email: karpagas1@srmist.edu.in

ABSTRACT

Turmeric is a most cultivated spice crop and is cherished for its aroma, taste and oleoresin content and used as herbal medicine and colouring powder for textile industry. India is the world largest producer of turmeric, followed by China, Indonesia, Bangladesh and Thailand. It occupies about six per cent of the total area under spices and condiments in India. The main turmeric producing states in India are Andhra Pradesh, Tamil Nadu, Orissa, Karnataka, West Bengal, Gujarat and Kerala. As a major crop, which is infected by several diseases *viz.*, rhizome rot, leaf spot and leaf blotch. Leaf spot incited by *Colletotrichum capsici* [Sydow (Butler & Bisby)] in turmeric is highly destructive causing yield loss up to 62 per cent with reduction in quality. Turmeric leaves showed typical leaf spot symptoms were collected and pathogen was isolated from the lesions. The effect of light on growth and sporulation of *C. capsici* was studied on potato dextrose agar by exposing the pure cultures in Petri plate to 12 h. dark + 12 h. light, 6 h. dark + 18 h. light, 3 h. dark + 21 h. light, 18 h. dark + 6 h. light dark and 21 h. dark + 3 h. light. The plates were incubated at 28±1°C for seven days at continuous dark by covering with aluminium foil. The control plates were kept under natural light and dark conditions without covering aluminium foil. Observations on colony diameter and sporulation were recorded. *C. capsici* showed the excellent conidia production under 3 h. dark + 21 h. light and continuous dark for seven days and the good sporulation was recorded in 6 h. dark + 18 h. light and continuous dark for seven days. Poor conidia production was observed in 21 h. dark + 3 h. light and continuous dark for seven days.

Key words: Turmeric Leaf spot, *Colletotrichum capsici*, Sporulation.

ECO-FRIENDLY SUPPRESSION OF GUAVA WHITEFLY THROUGH BIOLOGICAL CONTROL AGENTS WITHIN AN IPM STRATEGY

Vinothkumar G

Field Assistant, Department of Entomology, SRM College of Agricultural Sciences
Corresponding author Email: vinothkg1@srmist.edu.in

ABSTRACT

Guava (*Psidium guajava* L.) production is being affected by whitefly (*Dialeurodes citri*) infestations, a serious pest that causes severe damage by sucking sap and encouraging the growth of sooty mildew. This study assesses the efficacy of biological management employing natural enemies predators and parasitoids as a key component of Integrated Pest Management (IPM) for long-term whitefly suppression. Field trials were conducted in guava orchards to examine the impact of releasing major natural enemies, such as the green lacewing (*Chrysoperla carnea*), a predator, and (*Encarsia guadeloupae*), a parasite. Regular monitoring revealed significant reductions in whitefly populations in treated plots compared to untreated controls. To maximize the impact of predator and parasitoid releases, they were timed to coincide with peak whitefly populations. The findings showed that whitefly populations below economic threshold levels can be successfully managed by biological management alone, without the use of chemical interventions. Additionally, there was a decrease in the comeback of pests and the preservation of native beneficial insects. This study backs the use of parasitoids and predators as a practical, environmentally responsible method of managing whiteflies in guava, which is consistent with IPM and sustainable agriculture.

Keywords: Guava whitefly, Biological control, Integrated Pest Management (IPM), Natural enemies, Eco-friendly pest control

FIELD EVALUATION OF *Trichogramma chilonis* AND *Chrysoperla carnea* FOR THE BIOLOGICAL CONTROL OF *Helicoverpa armigera* IN TOMATO

Ramya P

Lab Assistant, Department of Entomology, SRM College of Agricultural Sciences
Corresponding author Email: ramyap3@srmist.edu.in

ABSTRACT

Tomato (*Solanum lycopersicum* L.) is a widely cultivated crop prone to infestation by the fruit borer, *Helicoverpa armigera*, a major pest that causes substantial yield and economic losses. This study evaluates the effectiveness of two biological control agents—*Trichogramma chilonis* (egg parasitoid) and *Chrysoperla carnea* (predator of early larval stages)—as key components of an Integrated Pest Management (IPM) strategy for managing *H. armigera* in tomato cultivation. Field experiments involved periodic releases of *T. chilonis* to parasitize eggs and reduce larval emergence, complemented by releases of *C. carnea* to prey on surviving early instar larvae. Monitoring results indicated a significant decrease in pest population and fruit damage in biologically treated plots compared to untreated controls. The combined use of these natural enemies maintained pest levels below the economic threshold and reduced the reliance on chemical insecticides. Furthermore, the conservation of other beneficial arthropods and improved fruit quality were observed in IPM-managed fields. These findings support the integration of both *T. chilonis* and *C. carnea* into a sustainable, eco-friendly IPM program for effective control of *H. armigera* in tomato ecosystems.

Keywords: Tomato fruit borer, Biological control, Integrated Pest Management (IPM), Egg parasitoid, Predator

**ENHANCING ANTIXENOSIS-MEDIATED RESISTANCE TO YELLOW STEM BORER
(*Scirpophaga incertulas* WALKER) THROUGH INTERCROPPING AND BORDER
CROP STRATEGIES IN IRRIGATED RICE ECOSYSTEMS**

Vairam Namachivayam¹ and Murugan Nagarajan^{*2}

¹Assistant Professor, SRM Valliammai Engineering College, Kattankulathur,
Chengalpattu, Tamil Nadu.

^{*2}Assistant Professor-Senior Grade, SRM College of Agricultural Sciences, Baburayenpettai,
Chengalpattu, Tamil Nadu.

*Corresponding author Email: murugan.agri@gmail.com

ABSTRACT

The yellow stem borer (*Scirpophaga incertulas* Walker) remains one of the most destructive pests of irrigated rice (*Oryza sativa* L.) across South and Southeast Asia, causing considerable yield losses. Continuous reliance on chemical insecticides has led to pest resurgence, resistance, and environmental degradation. To address these challenges, the present research focused on exploring host plant resistance, particularly antixenosis (non-preference), as an eco-friendly mechanism for pest suppression. The study was carried out during 2024–2025 at SRM Valliammai Engineering College, Kattankulathur, to evaluate the influence of intercropping and border crops on enhancing antixenosis-mediated protection against the yellow stem borer in irrigated rice systems. A Randomized Block Design (RBD) was adopted with the rice variety Co51 as the main crop, intercropped with sesame (*Sesamum indicum*), cowpea (*Vigna unguiculata*) and marigold (*Tagetes erecta*) and bordered with repellent crops such as maize (*Zea mays*) and sorghum (*Sorghum bicolor*). Observations on pest parameters, including the number of egg masses per hill, dead hearts, whiteheads, larval incidence, and grain yield, were recorded at key crop growth stages. Behavioral observations of oviposition preference and adult moth activity were also conducted to assess non-preference dynamics. The results revealed that rice plots intercropped with marigold + bordered with maize exhibited the lowest oviposition rate and larval incidence, recording a 42.8% reduction in dead hearts and 36.5% higher yield compared to the monocrop control. The sesame intercrop with sorghum border also demonstrated a strong deterrent effect, confirming that visual and olfactory interference from intercrops and border plants can disrupt the pest's host-location behaviour. Statistical analysis (ANOVA) indicated significant treatment differences at $p \leq 0.05$ for all recorded parameters. Overall, the experiment demonstrated that specific combinations of intercrops and border crops can effectively enhance antixenosis

expression in rice, thereby reducing pest pressure and pesticide dependence. The study provides a promising framework for developing ecological engineering-based Integrated Pest Management (IPM) models tailored for Tamil Nadu's irrigated rice ecosystems. The outcomes align with India's sustainable agriculture goals by promoting biodiversity-friendly, low-chemical, and climate-resilient rice cultivation systems.

Keywords: Rice, yellow stem borer, Antixenosis, inter-crop and Pest Management

Introduction

Rice (*Oryza sativa* L.) sustains more than half the global population. In South and Southeast Asia, *Scirpophaga incertulas* Walker (yellow stem borer, YSB) remains a persistent pest, causing yield losses up to 30–40% (Balaji et al., 2025). Conventional control through insecticides such as chlorantraniliprole and fipronil provides temporary relief but contributes to resistance, pest resurgence, and biodiversity loss (Patel et al., 2025).

Host plant resistance, particularly antixenosis, is a promising avenue within Integrated Pest Management (IPM). Antixenosis involves deterring pest colonization or oviposition through visual or chemical cues, thereby preventing initial infestation. Border crops and intercrops can further enhance antixenosis by disrupting pest orientation and host-seeking behavior (Nayak & Das, 2021). Despite progress in varietal resistance screening, there is limited research combining intercropping and border cropping for YSB suppression in irrigated rice ecosystems. This study hypothesized that appropriate crop combinations could synergistically enhance antixenosis-mediated resistance while maintaining or improving yield.

Genotypic variation in rice for resistance to YSB has been extensively reported. Balaji et al. (2025) identified several moderately resistant landraces with reduced dead-heart and white-head incidence under field conditions. Shai Prasanna and Joshi (2024a, 2024b) demonstrated that resistance is controlled by complementary gene action, suggesting potential for combining resistance and yield through breeding.

Eco-friendly management alternatives are gaining traction. Baliyan et al. (2025) reported that biopesticides and botanicals such as *Metarhizium anisopliae* and neem-based formulations were effective against YSB. Patel et al. (2025) found that seedling root-dip treatments using thiamethoxam and chlorantraniliprole significantly reduced infestation levels with minimal

residue accumulation. Intercropping and border cropping have shown success in managing other rice pests by creating habitat diversification and disrupting pest host-finding cues (Nayak & Das, 2021). However, limited empirical evidence exists for their integration with antixenosis-based mechanisms in YSB management. This study bridges that gap by evaluating border and intercropping combinations that may modify pest behavior through ecological interference. Based on that the following objectives were prepared.

1. To evaluate intercropping and border crop combinations for antixenosis-mediated resistance against YSB.
2. To quantify pest incidence, oviposition preference, and yield under various crop combinations.
3. To develop an ecological engineering-based IPM model suitable for Tamil Nadu's irrigated rice ecosystems.

Materials and Methods

Experimental Site

The experiment was conducted during the 2024–2025 *kharif* season at SRM Valliammai Engineering College, Kattankulathur, Tamil Nadu (12°44'N, 80°00'E). The site is characterized by a tropical climate with an average temperature of 30.5°C and annual rainfall of 1150 mm.

Experimental Design

A Randomized Block Design (RBD) with six treatments and three replications was adopted.

Table 1. Treatment details

Treatment Code	Cropping Pattern	Border Crop	Intercrop
T ₁	Rice (Co51) monocrop	None	None
T ₂	Rice + Sesame	None	Sesame (<i>Sesamum indicum</i>)

Treatment Code	Cropping Pattern	Border Crop	Intercrop
T ₃	Rice + Cowpea	None	Cowpea (<i>Vigna unguiculata</i>)
T ₄	Rice + Marigold	None	Marigold (<i>Tagetes erecta</i>)
T ₅	Rice + Marigold	Maize (<i>Zea mays</i>)	Marigold
T ₆	Rice + Sesame	Sorghum (<i>Sorghum bicolor</i>)	Sesame

Crop Establishment

Seedlings of rice (Co51) were transplanted at 25 × 25 cm spacing in 4 × 4 m plots. Border crops were sown two weeks prior to rice transplanting to ensure canopy establishment. Standard agronomic practices were maintained uniformly across treatments.

Data Collection

Pest parameters were recorded at 30, 60, and 90 DAT (days after transplanting):

- Number of egg masses per 10 hills
- Dead hearts (%) at vegetative stage
- White heads (%) at maturity
- Grain yield (t ha⁻¹)
- Behavioral observations included moth landing frequency and oviposition counts under field conditions using visual scoring at dusk (18:00–20:00 hrs).

Statistical Analysis

Data were subjected to ANOVA using SAS v9.4. Means were compared by Tukey's HSD ($p \leq 0.05$). Percentage data were arcsine-transformed prior to analysis.

Results

The results of the present investigation revealed significant variations among the different intercropping and border crop treatments in suppressing yellow stem borer incidence and enhancing rice yield.

Table 2 presents the influence of various intercropping and border cropping treatments on the incidence of yellow stem borer (YSB) and grain yield in irrigated rice. Significant reductions in both dead-heart and white-head percentages were observed across all diversified cropping treatments compared to the monocrop control (T₁). The rice monocrop (T₁) exhibited the highest pest incidence, with 9.8% dead hearts and 14.5% white heads, reflecting the vulnerability of a uniform cropping system to pest attack. Intercropping rice with sesame (T₂) resulted in a noticeable reduction in pest infestation, recording 7.2% dead hearts and 10.8% white heads. Similarly, the rice + cowpea treatment (T₃) further lowered infestation to 6.9% dead hearts and 9.6% white heads, suggesting moderate deterrence.

The rice + marigold combination (T₄) performed better, reducing dead hearts to 6.0% and white heads to 8.1%, which demonstrates marigold's repellency against YSB adults. Among all treatments, the marigold + maize border (T₅) was most effective, recording the lowest pest incidence of 5.6% dead hearts and 8.3% white heads. This treatment also achieved the highest grain yield (6.99 t ha⁻¹), corresponding to a 36.5% increase over the monocrop. The sesame + sorghum border combination (T₆) also provided strong protection, limiting dead hearts to 5.9% and white heads to 9.1%, with a yield of 6.72 t ha⁻¹ (31.3% gain). The observed reductions in pest infestation indicate that both visual and olfactory cues from intercrops and border crops disrupt the pest's host-finding behavior. Treatments involving non-host species such as maize and sorghum appeared to act as physical and sensory barriers to adult moth movement and oviposition. The improved yields under diversified systems further suggest that pest suppression directly translated into higher productivity without compromising crop health. Statistical analysis (ANOVA) confirmed that treatment differences were significant ($p \leq 0.05$) for all parameters recorded. The consistency of the trend across replications implies that the effect of intercropping and border cropping is reliable and reproducible under field conditions. Overall, Table 2 clearly demonstrates that strategic combinations of intercrops and border crops can enhance antixenosis-mediated resistance in rice, effectively minimizing pest damage and improving yield performance.

Table 2. Effect of intercropping and border cropping on yellow stem borer incidence and yield

Treatment	Dead hearts (%)	White heads (%)	Yield (t ha⁻¹)	Yield increase over control (%)
T ₁ – Rice monocrop	9.8	14.5	5.12	–
T ₂ – Rice + Sesame	7.2	10.8	5.78	12.9
T ₃ – Rice + Cowpea	6.9	9.6	5.85	14.3
T ₄ – Rice + Marigold	6.0	8.1	6.21	21.3
T ₅ – Rice + Marigold + Maize border	5.6	8.3	6.99	36.5
T ₆ – Rice + Sesame + Sorghum border	5.9	9.1	6.72	31.3

Figure 1. *Relative reduction (%) in dead-heart and white-head incidence in rice under different intercropping and border crop combinations compared to monocrop control.*

(Editable description: A clustered bar chart comparing each treatment (T₁–T₆) on two axes—Dead hearts % and White heads %. The highest reduction is seen in T₅ (marigold + maize border), followed by T₆ (sesame + sorghum). Error bars represent ±SE of means.)

Discussion

Intercropping and border cropping significantly reduced YSB infestation, supporting the hypothesis that habitat diversification enhances antixenosis. The marigold + maize combination (T₅) recorded the lowest oviposition and highest yield. The reduction in dead hearts (42.8%) and

increase in yield (36.5%) indicate a strong deterrence effect, likely mediated through visual contrast and volatile emissions that disrupt pest orientation.

These findings align with ecological engineering principles and previous studies demonstrating that plant diversity influences pest population dynamics (Baliyan et al., 2025). Similarly, Nayak and Das (2021) reported lower YSB incidence in farmer fields adopting non-host border crops. The study demonstrates that behavioral manipulation through plant spatial arrangement can complement genetic resistance. However, broader evaluation across seasons and sites is necessary to validate these results. Adoption of such IPM modules could reduce pesticide reliance by up to 40%, promoting biodiversity and sustainability in irrigated rice ecosystems.

References:

- Balaji, B., Vijaykumar, L., & Shivanna, B. (2025). Field evaluation of yellow stem borer (*Scirpophaga incertulas* Walker) incidence in selected rice genotypes. *Oryza – An International Journal of Rice*, 62(2), 175–181.
- Baliyan, S., Singh, H., Tiwari, M., Kumar, N., Singh, D., & Choudhary, A. (2025). Comparative efficacy of eco-friendly and chemical treatments for managing yellow stem borer (*Scirpophaga incertulas*) in Basmati rice. *Journal of Advances in Biology & Biotechnology*, 28(10), 986–995.
- Kattupalli, D., Barbadikar, K. M., Balija, V., et al. (2021). The draft genome of yellow stem borer provides insights into host specificity toward rice. *Insects*, 12(6), 563.
- Nayak, U., & Das, A. (2021). Farmer participatory assessment of pest management modules for yellow stem borer and leaf folder in summer rice. *Oryza – An International Journal of Rice*, 57(4), 350–357.
- Nyaupane, S. (2022). Evaluation of rice genotypes resistance to yellow stem borer, *Scirpophaga incertulas* (Walker) through sex pheromone trap. *Amrit Research Journal*, 3(1), 10–15.
- Patel, P. B., Patil, V. A., Mungra, K. S., & Gami, M. R. (2025). Response of seedling root dip treatments against yellow stem borer, *Scirpophaga incertulas* (Walker) infesting rice crop. *Journal of Agricultural Research and Reviews*, 18(1), 45–53.
- Shai Prasanna, G. S., & Joshi, J. L. (2024a). Studies on screening and inheritance of yellow stem borer (*Scirpophaga incertulas* Walker) resistance in F2 and F3 generations of rice (*Oryza sativa* L.). *Uttar Pradesh Journal of Zoology*, 45(10), 102–113.
- Shai Prasanna, G. S., & Joshi, J. L. (2024b). Identification of transgressive segregation for yield and resistance to yellow stem borer (*Scirpophaga incertulas* Walker). *Uttar Pradesh Journal of Zoology*, 45(11), 50–58.

**ECO-FRIENDLY PEST MANAGEMENT THROUGH DRIP IRRIGATION: A
SUSTAINABLE APPROACH**

Ramadass S^{*1}, Rajeshkumar A², Saravana Kumar M², Rajasekar M³ and Muthukumar M⁴

¹Asst. Prof. and Head, Dept. of Agronomy, SRMCAS, SRMIST, Chengalpattu-603201

²Asst. Prof., Dept. of Agronomy, SRMCAS, SRMIST, Chengalpattu-603201

³Asst. Prof., Dept. of Vegetable Science, SRMCAS, SRMIST, Chengalpattu-603201

⁴Asst. Prof. and Head, Dept. of Entomology, SRMCAS, SRMIST, Chengalpattu-603201

*Corresponding author Email: ramadass@srmist.edu.in

ABSTRACT

Pest control remains one of the most critical challenges in agriculture, often leading to significant crop losses and financial burdens for farmers. Traditional pest control methods, particularly the use of chemical pesticides, have shown limited effectiveness and have caused substantial harm to the environment, including the destruction of beneficial insect populations. This paper explores the role of drip irrigation as a modern, sustainable alternative that not only conserves water but also aids in effective pest control. Drip irrigation delivers water directly to the root zones of plants, minimizing moisture on plant surfaces and surrounding soil, thus creating an unfavorable environment for many common pests. Through a process known as fertigation, water-soluble pesticides and organic pest control agents can be delivered precisely to targeted areas, improving efficiency while reducing chemical usage. The system proves especially beneficial against soil-borne pests such as root aphids, nematodes, and fungus gnats, and also contributes to reducing populations of whiteflies, aphids, slugs, snails, and mosquitoes by altering environmental conditions necessary for their survival. Moreover, drip irrigation limits weed growth, indirectly reducing pest habitats, and contributes to healthier crops with increased pest resistance. The method, while not directly impacting canopy-level pests like caterpillars or beetles, enhances plant resilience, thereby supporting natural pest defense mechanisms. Combined with Integrated Pest Management (IPM) strategies, drip irrigation emerges as a powerful, eco-friendly solution that balances productivity with environmental responsibility. This approach offers farmers a sustainable method for pest control, addressing critical concerns such as water scarcity, environmental degradation, and long-term soil health, while promoting biodiversity and ecological balance in agricultural ecosystems.

Keywords: Drip Irrigation, Pest Control, Sustainable Agriculture

**BIOLOGY AND DEVELOPMENT OF THE BLACK SOLDIER FLY
(*Hermetia illucens* L.) REARED ON ORGANIC WASTE**

Ramazeame L

Assistant Professor, Department of Agricultural Entomology, SRM College of Agricultural Sciences, SRM Institute of Science and Technology (SRMIST), Chennai – 603203, Tamil Nadu, India.

Corresponding author Email: ramazeal@srmist.edu.in

ABSTRACT

The black soldier fly (*Hermetia illucens* L.) has emerged as a valuable insect species due to its ecological role in bioconversion of organic waste and potential as a protein source for feed industries. The present study was carried out to document the complete biology of the black soldier fly under controlled laboratory conditions in SRM College of agricultural sciences Baburayanpettai, Chennai, Tamil Nadu. Adult lifespan averaged 10–12 days, with females laying 480 ± 15 eggs per batch at 30 ± 1 °C and 75 ± 5 % RH. Egg incubation period lasted 4.8 ± 0.2 days, and larval duration averaged 19.6 ± 1.1 days, followed by a pupal stage of 15.2 ± 0.9 days. Using Dyar's rule, five larval instars were identified with mean lengths of 2.3, 6.2, 10.5, 15.6, and 17.1 mm, respectively. Adults exhibited positive phototaxis and humidity-dependent oviposition, while larvae showed strong olfactory responses toward decomposing organic matter. The entire lifecycle from egg to adult averaged 57 ± 2.5 days. Results indicate that *H. illucens* can be efficiently cultured under tropical Indian conditions, facilitating sustainable organic waste management and feed resource development.

Keywords: Black soldier fly, *Hermetia illucens*, biology, larval instars, organic waste, life cycle

INTRODUCTION

The growing global emphasis on sustainable waste management and alternative protein sources has revived interest in insects, particularly the black soldier fly (*Hermetia illucens*), for its multifaceted utility (Tomberlin and Sheppard, 2002). Waste accumulation in urban and peri-urban regions of India presents severe environmental challenges (FAO, 2023). Traditional composting and biogas production, while effective, are space- and cost-intensive.

The larvae of *H. illucens* feed voraciously on decomposing organic matter such as vegetable residues, manure, and food waste, converting it into high-value biomass rich in proteins (40–45%) and lipids (25–35%) (Zhang et al., 2010). The adult flies are non-pestiferous, non-feeding, and harmless to humans and livestock, making them suitable for controlled insect farming systems.

Previous research has mostly focused on waste reduction efficiency and larval feed utilization, while studies on complete biological parameters under tropical Indian conditions are scarce. Therefore, the present study was conducted to elucidate the life cycle, developmental stages, and morphological features of *H. illucens* under laboratory conditions at SRM College of Agricultural Sciences, Baburayanpettai .

MATERIALS AND METHODS

Study Site

The experiment was conducted during January–April 2025 in the insectary of the Department of Agricultural Entomology, SRM College of Agricultural Sciences, Baburayanpettai (12.3845° N, 79.7352° E) Ambient laboratory conditions were maintained at 30 ± 1 °C temperature, relative humidity 70–80 %, and 12L:12D photoperiod throughout the experimental period.

Culture Initiation

Adult black soldier flies were collected from compost heaps within the campus. They were maintained in mesh cages (60 × 60 × 60 cm) provided with moistened sponges as water sources and corrugated cardboard strips for oviposition. Egg clutches were transferred to plastic trays (40 × 30 × 15 cm) containing pre-fermented kitchen waste (vegetable peels, fruit residues, and bran).

Larval Rearing

Larvae were reared on 1 kg of homogenized food waste per tray, mixed with 100 g wheat bran to maintain consistency. The substrate was replaced every 3 days. Larval lengths were measured every 3 days using digital calipers.

Pupal and Adult Rearing

Mature larvae were separated at the pre-pupal stage by self-harvesting ramps and transferred to containers with moist soil for pupation. Newly emerged adults were sexed by external morphology (antennae and genitalia) and maintained for mating in cages under natural light.

Morphometric and Developmental Observations

Eggs, larvae, pupae, and adults were measured for length, width, and weight. The number of larval instars was determined using Dyar's law (Dyar, 1890). Oviposition, fecundity, incubation, and longevity were recorded through daily observations.

Statistical Analysis

All parameters were subjected to descriptive statistical analysis (mean $\pm$ SD). Environmental influences were noted qualitatively.

RESULTS AND DISCUSSION

Oviposition and Egg Morphology

Females began oviposition 6–7 days post-emergence and deposited clusters of 480 ± 15 eggs in crevices and cardboard folds near decomposing waste. Eggs were elongate, creamy white, and measured 0.90 ± 0.02 mm in length. Incubation required 4–5 days at 30 °C and 75 % RH, similar to reports by Sivanantharaja and Gnaneswaran (2018) from Sri Lanka.

Larval Development

Five distinct larval instars were recognized by mouth hook length and body segmentation (Table 1). The total larval period lasted 19.6 ± 1.1 days, and larvae displayed vigorous surface feeding behavior.

Table 1. Morphometric data of *H. illucens* larvae at different instars

Instars	Mean Length (mm) $\pm$ SD	Mean Width (mm) $\pm$ SD	Duration (days) $\pm$ SD

I	2.3 ± 0.03	0.4 ± 0.02	2.0 ± 0.1
II	6.2 ± 0.06	1.1 ± 0.03	3.0 ± 0.2
III	10.5 ± 0.08	2.1 ± 0.04	4.0 ± 0.3
IV	15.6 ± 0.10	3.1 ± 0.05	5.0 ± 0.4
V	17.1 ± 0.12	3.4 ± 0.05	5.6 ± 0.3

Larvae changed from creamy white to light brown as they matured. Feeding activity ceased at the pre-pupal stage, during which larvae turned dark brown and migrated away from moist areas seeking drier substrate for pupation.

Pupal and Adult Stages

The pupal period lasted 15.2 ± 0.9 days, with pupae measuring 17.2 ± 0.1 mm in length. Adult emergence occurred synchronously after 30–32 days from oviposition. Adults were wasp-like, black with translucent abdominal segments and non-functional mouthparts.

Mean adult lifespan was 10.6 ± 0.4 days, with females surviving slightly longer than males. Mean wingspan was 12.3 ± 0.2 mm (male) and 13.0 ± 0.2 mm (female).

Table 2. Summary of biological parameters of *H. illucens*

Parameter	Range	Mean $\pm$ SD
Egg period (days)	4–5	4.8 ± 0.2
Larval period (days)	18–22	19.6 ± 1.1
Pupal period (days)	13–17	15.2 ± 0.9
Adult longevity (days)	9–12	10.6 ± 0.4
Fecundity (eggs/female)	460–500	480 ± 15
Total life cycle (days)	55–59	57 ± 2.5

Feeding and Behavioral Observations

Larvae actively decomposed organic residues, producing nutrient-rich residue (frass). They preferred moist substrates with a moderate decomposition odor, confirming olfactory-driven feeding (Zhang et al., 2010). Adults were strongly phototactic and humidity-dependent; lack of moisture led to reduced mating activity.

Overall, the developmental duration and morphology observed under Chennai's tropical conditions were consistent with previous global reports (Sheppard et al., 2002; Tomberlin et al., 2009), validating the adaptability of *H. illucens* to Indian climates.

CONCLUSION

The black soldier fly completes its entire life cycle in about 57 days under tropical laboratory conditions in Chennai. The species shows high fecundity, rapid larval growth, and effective waste reduction potential. These findings confirm that *Hermetia illucens* can be efficiently mass-reared in Indian environments, supporting sustainable organic waste management and insect-based protein production initiatives.

ACKNOWLEDGEMENT

The author expresses sincere gratitude to the Department of Agricultural Entomology, SRM College of Agricultural Sciences, SRM institute of Science and Technology, for providing necessary facilities and technical assistance during the study.

REFERENCES

- Appelhof, M. (1982). *Worms Eat My Garbage*. Flower Press, Kalamazoo, MI.
- Briscoe, A. D., & Chittka, L. (2001). The evolution of color vision in insects. *Annual Review of Entomology*, 46:471–510.
- Dyar, H. G. (1890). The number of molts of lepidopterous larvae. *Psyche*, 5:420–422.
- Sheppard, D. C., Newton, G. L., Thompson, S. A., & Savage, S. (1994). A value-added manure management system using the black soldier fly. *Bioresource Technology*, 50:275–279.
- Sivanantharaja, A., & Gnanaswaran, R. (2018). Biology of the black soldier fly, *Hermetia illucens* (L.) in Jaffna, Sri Lanka. *Int. J. Entomol. Res.*, 3(6):18–22.
- Tomberlin, J. K., & Sheppard, D. C. (2002). Factors influencing mating and oviposition of black soldier flies. *Journal of Entomological Science*, 37:345–352.
- Zhang, J., Huang, L., He, J., et al. (2010). An artificial light source influences mating and oviposition of black soldier flies. *J. Insect Science*, 10:202.
- FAO (2023). *Circular Bioeconomy and Insect Protein Applications*. Rome: Food and Agriculture Organization.

IMAGE PROCESSING FOR HIGH-DENSITY MANGO FARMING: DETECTION OF PLANT DISEASE AND FRUIT GRADING

Arunkumar J¹* and Rajasekar P²

¹Phd (CSE) PT-INT, Department of Data Science and Business Systems, SRMIST, Kattankulathur, Tamil Nadu, India.

²Associate Professor, Department of Data Science and Business Systems, SRMIST, Kattankulathur, Tamil Nadu, India.

*Corresponding author Email: arunkumj@srmist.edu.in

ABSTRACT

High-density mango farming provides a promising pathway to boost agricultural productivity but brings new challenges in monitoring plant health and maintaining fruit quality. This study aims to develop an intelligent image processing system for automated disease detection and fruit grading in mango cultivation. The proposed approach integrates drone-based and ground-level imaging with machine learning techniques to support farmers in making informed decisions. A comprehensive image dataset will be collected, covering both healthy and diseased plants and fruits at various ripeness stages. Image pre-processing methods will be applied to improve clarity and contrast, enabling precise feature extraction. Convolutional Neural Networks (CNNs) will be trained to detect and classify plant diseases, while segmentation techniques will isolate affected regions for accurate diagnosis. Fruit grading will utilize color, size, and texture features, which are processed through machine learning classifiers to categorize mangoes into quality grades. Field experiments will assess the accuracy, precision, and recall of the models, ensuring reliable real-time performance. Additionally, a farmer-friendly application will be developed to provide instant disease alerts and grading reports. The expected benefits include early disease detection, improved grading consistency, reduced post-harvest losses, and higher market value. This research highlights the potential of integrating image processing and AI technologies to support sustainable high-density mango farming, ensuring better yield, profitability, and quality assurance.

Keywords: Image Processing, High-Density Mango Farming, Plant Disease Detection, Fruit Grading, Machine Learning, CNN, Precision Agriculture, Sustainable Farming.

ECO-INTELLIGENT TRAP CROPPING: A SUSTAINABLE APPROACH TO SMART PEST MANAGEMENT

Marimuthu Subramani* Sivakumar Kaliyanan and Jeyajothi Raman

Assistant Professor, Department of Agronomy, SRM College of Agricultural Sciences,
SRM Institute of Science and Technology, Baburayanpettai, Chengalpattu - 603 201

*Corresponding author Email: marimuts@srmist.edu.in

ABSTRACT

Eco-intelligent trap cropping, a time-tested ecological approach, is gaining renewed importance as a sustainable component of Integrated Pest Management (IPM). This strategy utilizes specific crops that are more attractive to insect pests than the main crop, thereby diverting pest populations and reducing crop damage. Beyond simple pest diversion, eco-intelligent cropping systems integrate ecological principles and precision agronomy to enhance system resilience, conserve natural enemies, and minimize pesticide dependence. The effectiveness of trap crops within an eco-intelligent framework depends on their inherent characteristics, pest behaviour, spatial arrangement, and synchronization with the main crop. Studies have shown that crops like squash around tomato or cantaloupe around cotton effectively attract pests such as whiteflies, lowering infestation and virus incidence. Moreover, eco-intelligent trap cropping provides multiple ecological services such as supporting natural enemy populations and improving agro-ecosystem stability. The success of these systems lies in their knowledge-intensive design, considering crop–pest interactions and local agro-climatic factors. By combining traditional ecological wisdom with modern precision techniques, eco-intelligent trap cropping emerges as a promising, smart, and sustainable solution to pest management, aligning with the goals of climate-resilient and environmentally responsible agriculture.

Keywords: Eco-intelligent cropping, Trap crops, Sustainable pest management, Pest diversion, Ecological resilience

**SMART FARMING FRONTIERS: PRECISION AGRICULTURE MEETS
INTEGRATED PEST MANAGEMENT**

Devi Vaishnavi V

PG Student, Department of Entomology, SRM College of Agricultural Sciences,
SRM Institute of Science and Technology, Baburayanpettai, Chengalpattu, Tamil Nadu

*Corresponding author Email: vaishnavivdevi21@gmail.com

ABSTRACT

The increasing global population has intensified the necessity for sustainable and efficient agricultural practices. One promising avenue for meeting this demand is the synergy between Precision Agriculture (PA) and Integrated Pest Management (IPM). This aims to scrutinize the multifaceted relationship between PA and IPM in augmenting crop protection and yield. Utilizing a comprehensive analysis of existing literature, the study elucidates how cutting-edge technologies in PA, such as drone imaging and soil sensor networks, can be harmoniously integrated with IPM strategies. These encompass biological, chemical, and cultural tactics to manage pest populations and mitigate damage, thereby fostering an environment conducive to optimal crop growth. The review identifies that the confluence of PA and IPM not only enhances the efficiency of resource use but also mitigates the environmental footprint of agricultural activities. Moreover, we delve into case studies that demonstrate significant yield improvements and cost reductions, underscoring the economic viability of integrating PA and IPM. The findings highlight the transformative potential of marrying these two domains, suggesting that such integration could be a cornerstone in the future of sustainable agriculture. The paper concludes by outlining research gaps and proposing avenues for future studies, emphasizing the need for multi-disciplinary approaches to fully unlock the potential of this integration.

Keywords: Agronomic crops; crop protection; integrated pest management (IPM); precision agriculture (PA); sustainable agriculture; technological innovations.

ADVANCING PEST CONTROL THROUGH RNAi GENE SILENCING

Naresh J¹ and Karpagavalli S^{2*}

¹PG Scholar, Department of Plant Pathology, SRM College of Agricultural Sciences, Vendhar Nagar, Baburayanpettai, Chengalpattu District – 603 201,

²Associate Professor (Plant Pathology), Department of Plant Pathology, SRM College of Agricultural Sciences, Vendhar Nagar, Baburayanpettai, Chengalpattu District – 603 201, Tamil Nadu, India

*Corresponding author Email: karpagas1@srmist.edu.in

ABSTRACT

The hunt for creative pest management techniques that reduce the use of chemical pesticides and save the environment has been fueled by the growing demand for sustainable agriculture practices. By precisely silencing vital genes needed for an insect's development, reproduction, or survival, RNA interference (RNAi) technology has become a potent and environmentally safe technique for controlling insect pests. A naturally occurring, highly conserved biological process called RNA interference (RNAi) uses short double-stranded RNA (dsRNA) molecules to control gene expression by causing complementary messenger RNA (mRNA) to degrade. RNA interference (RNAi) provides a tailored method to pest management, guaranteeing that beneficial organisms remain unaffected while only the targeted pest species are impacted. Through a variety of techniques, including feeding, spraying, or plant-mediated expression, dsRNA is delivered as part of the process. The RNA-induced silencing complex (RISC) recognizes the dsRNA once it is inside the cell and directs the destruction of particular mRNAs, upsetting critical physiological circuits. With little ecological disturbance, pest populations are suppressed as a result of this process. For instance, RNAi-based tactics have been effective in combating significant pests such as aphids, cotton bollworms, and western corn rootworms. The stability of dsRNA in the environment, efficient transport to target tissues, and variation in RNAi response across insect groups are some of the obstacles that RNAi technology must overcome despite its potential. Through better formulation methods, more comprehension of insect RNAi machinery, and sophisticated nanoparticle carriers, ongoing research seeks to overcome these constraints. A new generation of economically and environmentally sound precision pest control systems may result from combining RNAi with conventional pest management techniques

Keywords: RNA interference, gene silencing, sustainable pest management, dsRNA, environmentally safe control

**PLANT-DERIVED BIOFUMIGANTS: A SAFER AND SUSTAINABLE
ALTERNATIVE FOR STORED PEST MANAGEMENT**

Sharulatha S^{*1}, Muthukumar M² and Anandhi S³

¹PG Scholar, Department of Entomology, SRM College of Agricultural Sciences, SRMIST,
Baburayanpettai, Chengalpattu District, Tamil Nadu– 603201

²Asst. Professor and Head, Department of Entomology, SRMCAS, SRMIST,
Baburayanpettai, Chengalpattu District, Tamil Nadu– 603201

³Asst. Professor and Head, Department of Floriculture and Landscape Architecture, SRMCAS,
SRMIST, Baburayanpettai, Chengalpattu District, Tamil Nadu– 603201

*Corresponding author Email: ss6710@srmist.edu.in

ABSTRACT

Stored grain loss caused by insect pests is a significant issue in post-harvest storage. It is being overcome using synthetic fumigants, which cause greater risk to human health and the environment, and many insects have developed resistance. With the aim of minimising it in a more eco-friendly way, plant extracts are gaining attention. Certain medicinal and aromatic plants that have insecticidal properties can be used as a safer alternative to prevent this storage loss. For instance, Compounds like 4-nonanone, linalool, isopulegol 1, caryophyllene oxide, alpha-cadinol, 4,11,11-trimethyl-8-methylene-bicyclo[7.2.0]undec-4-ene in *Cymbopogon citratus* and isopulegal 1, methyl linolenate, aromandendrene, (-)-globulol, rosifoliol, caryophyllene-(i1), 3-octen-5-yne, 2,7-dimethyl-, (z)-, hedycaryol, 3-cyclohexen-1-ol, 4-methyl-1-(1-methylethyl)- in *Eucalyptus tereticornis* are known to possess insecticidal properties. These compounds have the potential to act as biofumigants, an eco-friendly and sustainable alternative for stored pest management.

Keywords: Stored pest management, Synthetic fumigants, Biofumigants, sustainable alternatives.

BLACK SOLDIER FLY LARVAE MEAL: ALTERNATIVE DIETARY PROTEIN SOURCES FOR THE CHICKEN

Sreyass K S* and Karthikeyan G

Assistant Professor, Section of Animal Sciences, SRM College of Agricultural Sciences, SRM IST, Chengalpattu

*Corresponding author Email: sreyassk@srmist.edu.in

ABSTRACT

India's poultry sector is witnessing robust growth, with an annual expansion rate of over 8%, driven by rising demand for affordable animal protein and increasing per capita consumption. Protein is a critical component in poultry diets, essential for muscle development, immune function, and optimal production performance. However, the escalating cost and inconsistent availability of conventional protein sources like soybean meal have prompted the search for sustainable and cost-effective alternatives. Black Soldier Fly Larvae (BSFL) meal emerges as a promising substitute, offering a crude protein content ranging from 40% to 55%, depending on processing methods and substrate. Studies suggest that BSFL meal can be safely included in poultry diets at levels ranging from 5% to 25%, depending on the species and production stage. BSFL can be mass-produced using organic waste streams such as food scraps, poultry litter, and agro-industrial by-products, making it both economical and environmentally friendly. The larvae are harvested, dried, and processed into meal, oil, or powder forms suitable for feed formulation. Incorporating BSFL in poultry diets has shown to improve feed conversion ratio (FCR), enhance growth rates in broilers, and increase egg production and shell quality in layers. Additionally, its antimicrobial peptides contribute to gut health, while its production supports circular bioeconomy through effective waste valorization. The rearing process requires minimal land and water, making it ideal for decentralized, low-cost protein production.

Keywords: Black Soldier Fly, Larvae Meal, Alternative Dietary, Protein, Chicken

APHID MECHANISM OF RESISTANT IN GROUNDNUT GENOTYPES

Vanitha J* and Mahendran R

Assistant Professor Department of Genetics and Plant Breeding, SRMCAS,
Chengalpattu– 603201.

*Corresponding author Email: vanithaj@srmist.edu.in

ABSTRACT

Aphid infestation (*Aphis craccivora* Koch) is a major constraint in groundnut (*Arachis hypogaea* L.) production, causing direct feeding damage and transmitting peanut stripe and rosette viruses. Understanding the mechanisms of resistance among groundnut genotypes is crucial for developing sustainable pest management strategies. The present study investigates the morphological, biochemical, and molecular bases of aphid resistance in selected groundnut genotypes. Resistant genotypes exhibited lower aphid populations and slower multiplication rates compared to susceptible ones. Morphological traits such as higher trichome density, thicker epidermal layers, and harder pod shells contributed to antixenosis and antibiosis effects. Biochemical analyses revealed elevated levels of phenolic compounds, tannins, and defensive enzymes (peroxidase, polyphenol oxidase, and catalase) in resistant lines, which deter aphid feeding and reproduction. Additionally, certain genotypes demonstrated tolerance through rapid recovery and maintenance of photosynthetic efficiency under infestation. The study highlights the combined action of morphological and biochemical defenses as key determinants of resistance, providing valuable insights for breeding programs aimed at developing aphid-resistant groundnut cultivars.

Keywords: Aphid, Mechanism of Resistant, Groundnut, Genotypes, *Arachis hypogaea*

**MORPHOLOGICAL AND MOLECULAR CHARACTERIZATION OF ROOT-KNOT
NEMATODE, *Meloidogyne enterolobii* INFECTING GUAVA (*Psidium guajava* L.)**

Ashokkumar Natarajan¹, Sarrvesh lakshman², Rageshwari S³ and Gopu B⁴

¹Assistant Professor (Nematology), Dept. of Plant Protection,
SRM College of Agricultural Sciences

²PG Scholar, Dept. of Plant Pathology, SRM College of Agricultural Sciences

³Assistant Professor Sr.Grade, Dept. of Plant Protection, SRM College of Agricultural Sciences

⁴Associate Professor and Head, Dept. of Fruit Science, SRM College of Agricultural Sciences

*Corresponding author Email: ashokkun@srmist.edu.in

ABSTRACT

Root-knot nematode (*Meloidogyne enterolobii*) has emerged as a serious threat to guava (*Psidium guajava* L.) cultivation in several tropical regions. The present study aimed to characterize the nematode populations associated with guava wilt–nematode disease complex through detailed morphological, morphometric and molecular approaches. Infected root samples exhibiting typical galling symptoms were collected from major guava-growing districts of Tamil Nadu. The adult females were dissected from roots and examined for perineal pattern configuration, morphological and morphometric indices. The perineal pattern was oval to high dorsal arch without lateral lines, typical of *M. enterolobii*. Second-stage juveniles (J2) possessed a distinct hemispherical head, slender tail with a rounded terminus and a stylet length of 14.5–16.2 µm. Molecular identification using species-specific primers and amplification of the ITS and COI regions confirmed the identity of *M. enterolobii*. Phylogenetic analysis clustered the isolates with reference *M. enterolobii* sequences available in GenBank. The integrative characterization confirms the presence and distribution of *M. enterolobii* in guava orchards of Tamil Nadu and provides a reliable diagnostic basis for subsequent epidemiological and management studies.

Keywords: *Meloidogyne enterolobii*, guava, morphological characterization, molecular identification, COI, ITS



SKYFOX PUBLISHING GROUP
SKYFOX PRESS
#1789, MEDICAL COLLEGE ROAD
THANJAVUR-613004 TAMIL NADU,
INDIA.

ISBN 978-93-90357-33-8



9 789390 357338

Department of Entomology
SRM College of Agricultural Sciences
SRM Institute of Science and Technology
Baburayanpettai, Chengalpattu Dist. - 603 201